

**PATENT APPLICATION**

**MODULATORS OF ANGIOGENESIS AND TUMORIGENESIS**

**Inventor(s):**

James B. Lorens, a citizen of the United States, residing at  
Kraakenesveien 10  
N-5152 Bønes NORWAY

Robert E. Atchison, a citizen of the United States, residing at  
631 O'Farrell St., #911  
San Francisco, CA 94109

Anabella Frieria, a citizen of the United States, residing at  
229 Carmelo Lane  
South San Francisco, CA 94080

Sacha Holland, a citizen of the United Kingdom, residing at  
518 Foerster Street, Apt. B  
San Francisco, CA 94127

**Assignee:**

Rigel Pharmaceuticals, Incorporated  
1180 Veterans Blvd.  
South San Francisco, CA 94080

**Entity:** Small business concern

## MODULATORS OF ANGIOGENESIS AND TUMORIGENESIS

### CROSS-REFERENCES TO RELATED APPLICATIONS

The present application claims priority to USSN 60/421,989, filed October 29, 2002, and USSN 60/\_\_\_\_\_, TTC Ref. No. 021044-005810, filed October 17, 2003, each herein incorporated by reference in their entirety.

### STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

Not applicable.

### FIELD OF THE INVENTION

The present invention relates to regulation of angiogenesis and tumorigenesis. The invention further relates to methods for identifying and using agents, including small organic molecules, antibodies, peptides, cyclic peptides, nucleic acids, antisense nucleic acids, RNAi, and ribozymes, that modulate angiogenesis and tumorigenesis via modulation of endothelial cell haptotaxis; as well as to the use of expression profiles and compositions in diagnosis and therapy of angiogenesis and cancer.

### BACKGROUND OF THE INVENTION

The migration of activated endothelial cells through a vitronectin-rich provisional matrix is critical to the formation of new blood vessels during angiogenesis and is dependent on adhesion receptors containing  $\alpha$ v integrins (such as  $\alpha$ v $\beta$ 3 which binds to vitronectin). Peptide and antibody inhibitors of  $\alpha$ v $\beta$ 3 integrin inhibit tumor growth *in vivo*.

Angiogenesis is typically limited in a normal adult to the placenta, ovary, endometrium and sites of wound healing. However, angiogenesis, or its absence, plays an important role in the maintenance of a variety of pathological states. Some of these states are characterized by neovascularization, e.g., cancer and tumorigenesis, e.g., endometriosis, diabetic retinopathy, glaucoma, glomerulonephritis, and age related macular degeneration. Others, e.g., stroke, infertility, heart disease, e.g., restenosis, ulcers, and scleroderma, are diseases of angiogenic insufficiency. Therefore, there is a

need to identify nucleic acids encoding proteins involved in the regulation of angiogenesis and tumorigenesis, to identify, e.g., modulators of angiogenesis, as well as new therapeutic and diagnostic applications.

5

## SUMMARY OF THE INVENTION

Novel targets for anti-angiogenic and anti-tumorigenic therapy have been identified using a functional genetic screen based on endothelial cell haptotaxis.

Inhibition or activation of these targets (by small molecule inhibitors; protein, antibody and peptide therapeutics; RNAi; antisense; gene therapy etc.) have therapeutic value in modulating angiogenesis and tumorigenesis,, e.g., breast, lung, colon, ovarian, liver, thyroid, stomach, bladder, and prostate cancer, basal cell carcinoma, melanoma, lymphomas, leukemias, e.g., myeloid leukemia (CML and AML), endometriosis, diabetic retinopathy, glaucoma, glomerulonephritis, age related macular degeneration, as well as, e.g., stroke, infertility, heart disease, e.g., restenosis, ulcers, and scleroderma.

15

The present invention therefore provides nucleic acids encoding proteins involved in modulation of vitronectin induced endothelial cell haptotaxis and modulation of angiogenesis and tumorigenesis. The invention therefore provides methods of screening for compounds, e.g., small organic molecules, antibodies, nucleic acids, peptides, cyclic peptides, nucleic acids, antisense molecules, RNAi, and ribozymes, that are capable of modulating angiogenesis or tumorigenesis, e.g., either activating or inhibiting angiogenesis or tumorigenesis. Therapeutic and diagnostic methods and reagents are also provided.

20

In one aspect, the present invention provides a method for identifying a compound that modulates angiogenesis and tumorigenesis, the method comprising the steps of: (i) contacting the compound with an angiogenesis or tumorigenesis modulating polypeptide or fragment thereof, the polypeptide encoded by a nucleic acid that hybridizes under stringent conditions to a reference nucleic acid encoding the polypeptide, the polypeptide selected from the group consisting of Axl, tubulin cofactor D, transglutaminase 2, cytosine deaminase, peptidase M41 (paraplegin), CD13 aminopeptidase, PRK-1, zip kinase, Gas6, SRm160, non-muscle myosin heavy chain, calmodulin 2, novel symporter, novel semaphorin, novel zinc finger helicase (FLJ22611), plexin-A2, deoxycytidylate deaminase, and a novel sugar transporter; and (ii) determining the functional effect of the compound upon the polypeptide.

30

In one embodiment, the functional effect is determined *in vitro*. In another embodiment, the functional effect is a physical effect. In another embodiment, the functional effect is determined by measuring ligand binding to the polypeptide. In another embodiment, the functional effect is a chemical effect.

5                   In one embodiment, the polypeptide is expressed in a eukaryotic host cell. In another embodiment, the functional effect is a physical effect. In another embodiment, the functional effect is determined by measuring ligand binding to the polypeptide. In another embodiment, the functional effect is a chemical or phenotypic effect. In another embodiment, the polypeptide is expressed in a eukaryotic host cell, e.g., an endothelial  
10 cell. In another embodiment, the functional effect is determined by measuring  $\alpha v \beta 3$  expression, haptotaxis, tumor cell proliferation, or tumor growth *in vivo*.

In one embodiment, modulation is inhibition of angiogenesis or tumorigenesis.

In one embodiment, the polypeptide is recombinant.

15                   In one embodiment, the compound is an antibody, a peptide, an antisense molecule, a RNAi molecule, or a small organic molecule.

In another aspect, the present invention provides a method for identifying a compound that modulates tumorigenesis or angiogenesis, the method comprising the steps of (i) contacting the compound with an angiogenesis or tumorigenesis modulating  
20 polypeptide or fragment or inactive variant thereof, the polypeptide selected from the group consisting of Ax1, tubulin cofactor D, transglutaminase 2, cytosine deaminase, peptidase M41 (paraplegin), CD13 aminopeptidase, PRK-1, zip kinase, Gas6, SRm160, non-muscle myosin heavy chain, calmodulin 2, novel symporter, novel semaphorin, novel zinc finger helicase (FLJ22611), plexin-A2, deoxycytidylate deaminase, and a novel  
25 sugar transporter (ii) determining the physical effect of the compound upon the polypeptide or fragment thereof or inactive variant thereof; and (iii) determining the chemical or phenotypic effect of the compound upon a cell comprising the polypeptide or fragment thereof or inactive variant thereof, thereby identifying a compound that modulates tumorigenesis or angiogenesis.

30                   In another aspect, the present invention provides a method of modulating angiogenesis or tumorigenesis in a subject, the method comprising the step of administering to the subject a therapeutically effective amount of a compound identified using the methods described herein.

In one embodiment, the subject is a human.

In one embodiment, the compound is an antibody, an antisense molecule, a peptide, or an RNAi molecule, or a small organic molecule.

In one embodiment, the compound inhibits angiogenesis or tumorigenesis.

5

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a screening protocol for haptotactic migration inhibitors.

Figure 2 shows a cellomics haptotaxis assay.

Figure 3 shows that a CD13/N-aminopeptidase is involved in haptotaxis.

Figure 4 shows that a transglutaminase II protein is involved in haptotaxis.

10

Figure 5 shows that a zip kinase protein is involved in haptotaxis.

Figure 6 shows that a PRK-1 protein is involved in haptotaxis.

Figure 7 shows that PRK1 mRNA is expressed in endothelial cells and

PBMCs.

Figure 8 shows that a PRK-1 protein is involved in haptotaxis and  $\alpha\beta$

15

expression.

Figure 9 shows that PRK-1 RNAi reduces PRK-1 message, haptotaxis and  $\alpha\beta$  expression.

Figure 10 shows that PRK-1 RNAi reduces tube formation in a co-culture assay.

20

Figure 11 shows that Axl and Gas6 are involved in haptotaxis.

Figure 12 shows that Axl RNAi reduces Axl mRNA expression.

Figure 13 shows that Axl RNAi inhibit haptotaxis to vitronectin.

Figure 14 shows that Axl RNAi reduces Axl protein expression.

Figure 15 shows that Axl RNAi inhibits vitronectin haptotaxis and

25

HUVEC proliferation.

Figure 16 shows that an Axl extracellular domain was isolated in a VEGF-R2 screen.

Figure 17 shows that Axl RNAi inhibits tube formation in a co-culture assay.

30

Figure 18 shows a sponge angiogenesis/xenograft model.

Figure 19 shows that a deoxycytidylate deaminase is involved in angiogenesis.

Figure 20 shows that a plexin-A2 is involved in angiogenesis.

## DETAILED DESCRIPTION OF THE INVENTION

### Introduction

We have used a functional genetic screening strategy to identify proteins involved in regulating endothelial cell migration on specific matrix components, e.g. vitronectin. Using a retroviral-based system, we have stably expressed complex libraries of various types of genetic elements (e.g. cDNAs and GFP-fusions) in human primary endothelial cells (e.g. HUVECs). Starting with early passage primary endothelial cells representing an "angiogenic state" (i.e., proliferative, highly motile), single cells that have switched to a more differentiated "angiostatic state" (i.e., quiescent, reduced motility) can be identified. In order to focus on the migration step of the angiogenic cycle, conditions were established which allowed highly efficient migration of HUVEC cells along a haptotactic gradient of specific matrix proteins (e.g. vitronectin / fibronectin) in a Boyden chamber assay. Large populations of HUVEC cells were then infected with a GFP-fused cDNA library and selected for impaired haptotaxis. Additional assays for proteins involved in angiogenesis and tumorigenesis include a VEGFR2 assay, a HUVEC/smooth muscle cell co-culture assay for endothelial tube formation, chemo-invasion assays, and tumor cell proliferation assays. Finally, human/mouse tumor xenograft assays, mouse sponge angiogenesis and tumorigenesis assays (sponge with human ECs +/- human tumor cells in SCID mice), a collagen-antibody induced arthritis model for RA, and retinal neovascularization assays can be used to identify angiogenesis and tumorigenesis proteins *in vivo* and to assay for modulators of such proteins.

The angiogenesis and tumorigenesis proteins identified using the haptotaxis assay described herein, e.g., Ax1, its ligand, Gas6, tubulin cofactor D, transglutaminase 2, cytosine deaminase, plexin-A2, peptidase M41 (paraplegin), CD13 aminopeptidase, PRK-1, zip kinase, SRm160, non-muscle myosin heavy chain, calmodulin 2, novel symporter, novel semaphorin, novel zinc finger helicase (FLJ22611), and a novel sugar transporter, therefore represent targets for the development of angiogenic drugs, preferably anti-angiogenic drugs, e.g., anti-angiogenic drugs for treatment of neovascularization, e.g., cancer, diabetic retinopathy, endometriosis, glomerulonephritis, restenosis, glaucoma, rheumatoid arthritis, and age related macular degeneration, or angiogenic drugs for treatment of angiogenic insufficiency, e.g., stroke, infertility, heart disease, ulcers, and scleroderma. Modulators include small organic molecules, nucleic acids, peptides, cyclic peptides, antibodies, antisense molecules, RNAi molecules, and ribozymes. The nucleic acids and proteins of the invention are also useful

for diagnostic applications, using, e.g., nucleic acid probes, oligonucleotides, and antibodies. These polypeptides are also involved in tumorigenesis and cellular proliferation, and are useful for the development of therapeutic molecules to treat diseases associated with angiogenesis, tumorigenesis, and cellular proliferation. Furthermore, the polypeptides described herein and the nucleic acids encoding them are useful for diagnostic assays for diseases associated with angiogenesis, tumorigenesis, and cellular proliferation.

In one embodiment, the protein Axl is involved in angiogenesis, tumorigenesis, and cellular proliferation. Axl is a receptor tyrosine kinase that possesses transforming activity *in vitro*. Axl is activated by its ligand, Gas 6. Axl, Sky, and Mer form a family of receptor tyrosine kinases that are activated by Gas6. Axl is expressed in myeloid cell types and vasculature, e.g., endothelial cells (HUVEC), VSMC, monocytes, macrophages, CD34+ cells, bronchial epithelium, breast epithelium, neurons, and chondrocytes. Axl is upregulated in VSMC after vascular injury. Gas6-Axl interactions stimulate cell survival and chemotaxis. Knockout and transgenic mice for Axl show no overt phenotypes. Axl is associated with several diseases, including rheumatoid arthritis (O'Donnell *et al.*, *Am. J. Pathol.* 154:1171-1180 (1999)), endometriosis (Sun *et al.*, *Mol. Human Reproduction* 8:552-559 (2002)), cancer, e.g. myeloid leukemias such as AML and CML, thyroid carcinomas, and breast cancer. Axl can be activated using antibody crosslinking, pervanadate, or Gas6.

In another embodiment, the protein PRK-1 is involved in angiogenesis and tumorigenesis. PRK-1 is a cytoplasmic serine/threonine kinase related to PKC. PRK-1 binds Rho GTPase through the N-terminus and is regulated by activation loop phosphorylation by PD-1. PRK activation by PDK1 is Rho-dependent and membrane localized. PRK-1 I exhibits elevated expression in endothelial cells and PBMC. PRK-1 is upregulated in breast and prostate epithelial carcinoma cells as compared to normal cells and activates androgen receptor signaling (*see, e.g., Metzger et al., EMBO J.* 22:270-280 (2003)). PRK-1 RNAi inhibits PRK-1 protein and VN and FN haptotaxis, (including a PRK-1 3' UTR RNAi and a PRK-1 RNAi vector). PRK-1 RNAi reduces tube formation in the co-culture assays, as well as SDF-1 and VEGF chemotaxis in HUVECs. PRK-1 RNAi inhibits vitronectin-induced haptotaxis in PSMCs. Expression of kinase inactive PRK-1 in HUVECs increases vitronectin haptotaxis. The PRK-1 screening hit GH1-54 co-immunoprecipitates with full-length PRK-1. *In vitro* kinase

assays with PRK-1 can be used to assay for PRK-1 modulators and thus for modulators of angiogenesis.

In another embodiment, the protein PRK-2 is involved in angiogenesis and tumorigenesis. PRK-2 is widely expressed in adherent cell types. PRK-2 siRNA inhibits vitronectin-induced haptotaxis in HUVECs and PSMCs and also inhibits PRK-2 protein expression. In another embodiment, the proteins deoxycytidylate deaminase (sense hit) (, transglutaminase II (antisense hit), plexin-A2 (sense hit), and ZIP-kinase/DAP kinase 3 (sense 3' UTR hit), are involved in angiogenesis and tumorigenesis. ZIP kinase exhibits a restricted tissue expression pattern and is highly expressed in hepatocytes. ZIP kinase RNAi inhibits VN haptotaxis.

### Definitions

By “disorder associated with angiogenesis or tumorigenesis” or “disease associated with angiogenesis or tumorigenesis” herein is meant a disease state which is marked by either an excess or a deficit of vessel development. Angiogenesis and tumorigenesis disorders associated with increased angiogenesis include, but are not limited to, breast, lung, colon, ovarian, liver, stomach, bladder, thyroid, and prostate cancer, basal cell carcinoma, melanoma, lymphomas, leukemias, e.g., myeloid leukemia (AML, CML), endometriosis, diabetic retinopathy, glaucoma, glomerulonephritis, rheumatoid arthritis, and age related macular degeneration. Pathological states for which it may be desirable to increase angiogenesis include stroke, infertility, heart disease, e.g., restenosis, ulcers, and scleroderma. An increase in angiogenesis may also be desirable in transplantation or for artificial or *in vitro* growth of organs.

By “disorder associated with cellular proliferation or tumorigenesis” or “disease associated with cellular proliferation or tumorigenesis” herein is meant a disease state which is marked by either an excess or a deficit of cellular proliferation or apoptosis. Such disorders associated with increased cellular proliferation include, but are not limited to, cancer and non-cancerous pathological proliferation.

The terms “angiogenesis and/or tumorigenesis polypeptide” or a nucleic acid encoding an “angiogenesis and/or tumorigenesis polypeptide” refer to nucleic acid and polypeptide polymorphic variants, alleles, mutants, and interspecies homologs that: (1) have an amino acid sequence that has greater than about 60% amino acid sequence identity, 65%, 70%, 75%, 80%, 85%, 90%, preferably 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or greater amino acid sequence identity, preferably over a region of

over a region of at least about 25, 50, 100, 200, 500, 1000, or more amino acids, to a polypeptide encoded by a referenced nucleic acid or an amino acid sequence described herein; (2) specifically bind to antibodies, e.g., polyclonal antibodies, raised against an immunogen comprising a referenced amino acid sequence, immunogenic fragments thereof, and conservatively modified variants thereof; (3) specifically hybridize under stringent hybridization conditions to a nucleic acid encoding a referenced amino acid sequence, and conservatively modified variants thereof; (4) have a nucleic acid sequence that has greater than about 95%, preferably greater than about 96%, 97%, 98%, 99%, or higher nucleotide sequence identity, preferably over a region of at least about 25, 50, 100, 200, 500, 1000, or more nucleotides, to a reference nucleic acid sequence. A polynucleotide or polypeptide sequence is typically from a mammal including, but not limited to, primate, e.g., human; rodent, e.g., rat, mouse, hamster; cow, pig, horse, sheep, or any mammal. The nucleic acids and proteins of the invention include both naturally occurring or recombinant molecules.

The phrase “functional effects” in the context of assays for testing compounds that modulate activity of an angiogenesis and tumorigenesis protein includes the determination of a parameter that is indirectly or directly under the influence of an angiogenesis polypeptide, e.g., a chemical or phenotypic effect such as loss-of angiogenesis or tumorigenesis phenotype represented by a change in expression of a cell surface marker  $\alpha v \beta 3$  integrin, changes in cellular migration, changes in endothelial tube formation, and changes in tumor growth, or changes in cellular proliferation, especially endothelial cell proliferation; or enzymatic activity; or, e.g., a physical effect such as ligand binding or inhibition of ligand binding. A functional effect therefore includes ligand binding activity, the ability of cells to proliferate, expression in cells undergoing angiogenesis or tumorigenesis, and other characteristics of angiogenic and tumorigenic cells. “Functional effects” include *in vitro*, *in vivo*, and *ex vivo* activities.

By “determining the functional effect” is meant assaying for a compound that increases or decreases a parameter that is indirectly or directly under the influence of an angiogenesis protein, e.g., measuring physical and chemical or phenotypic effects. Such functional effects can be measured by any means known to those skilled in the art, e.g., changes in spectroscopic characteristics (e.g., fluorescence, absorbance, refractive index); hydrodynamic (e.g., shape), chromatographic; or solubility properties for the protein; ligand binding assays, e.g., binding to antibodies; measuring inducible markers or transcriptional activation of the angiogenesis protein; measuring changes in enzymatic

activity; the ability to increase or decrease cellular proliferation, apoptosis, cell cycle arrest, measuring changes in cell surface markers, e.g.,  $\alpha v \beta 3$  integrin; and measuring cellular proliferation, particularly endothelial cell proliferation. Determination of the functional effect of a compound on angiogenesis or tumorigenesis can also be performed using assays known to those of skill in the art such as endothelial cell tube formation assays; haptotaxis assays; the chick CAM assay; the mouse corneal assay; VEGF receptor assays, co-culture tube formation assays, and assays that assess vascularization of an implanted tumor. Tumorigenesis can be measured using *in vivo* mouse models such as a xenograft model. The functional effects can be evaluated by many means known to those skilled in the art, e.g., microscopy for quantitative or qualitative measures of alterations in morphological features, e.g., tube or blood vessel formation, measurement of changes in RNA or protein levels for angiogenesis-associated sequences, measurement of RNA stability, identification of downstream or reporter gene expression (CAT, luciferase,  $\beta$ -gal, GFP and the like), e.g., via chemiluminescence, fluorescence, colorimetric reactions, antibody binding, inducible markers, etc.

“Inhibitors,” “activators,” and “modulators” of angiogenesis and tumorigenesis polynucleotide and polypeptide sequences are used to refer to activating, inhibitory, or modulating molecules identified using *in vitro* and *in vivo* assays of angiogenesis and tumorigenesis polynucleotide and polypeptide sequences. Inhibitors are compounds that, e.g., bind to, partially or totally block activity, decrease, prevent, delay activation, inactivate, desensitize, or down regulate the activity or expression of angiogenesis and tumorigenesis proteins, e.g., antagonists. “Activators” are compounds that increase, open, activate, facilitate, enhance activation, sensitize, agonize, or up regulate angiogenesis and tumorigenesis protein activity, agonists. Inhibitors, activators, or modulators also include genetically modified versions of angiogenesis and tumorigenesis proteins, e.g., versions with altered activity, as well as naturally occurring and synthetic ligands, antagonists, agonists, antibodies, peptides, cyclic peptides, nucleic acids, antisense molecules, ribozymes, RNAi molecules, small organic molecules and the like. Such assays for inhibitors and activators include, e.g., expressing angiogenesis or tumorigenesis protein *in vitro*, in cells, or cell extracts, applying putative modulator compounds, and then determining the functional effects on activity, as described above.

Samples or assays comprising angiogenesis or tumorigenesis proteins that are treated with a potential activator, inhibitor, or modulator are compared to control samples without the inhibitor, activator, or modulator to examine the extent of inhibition.

Control samples (untreated with inhibitors) are assigned a relative protein activity value of 100%. Inhibition of an angiogenesis or tumorigenesis protein is achieved when the activity value relative to the control is about 80%, preferably 50%, more preferably 25-0%. Activation of an angiogenesis or tumorigenesis protein is achieved when the activity value relative to the control (untreated with activators) is 110%, more preferably 150%, more preferably 200-500% (i.e., two to five fold higher relative to the control), more preferably 1000-3000% higher.

The term “test compound” or “drug candidate” or “modulator” or grammatical equivalents as used herein describes any molecule, either naturally occurring or synthetic, e.g., protein, oligopeptide (e.g., from about 5 to about 25 amino acids in length, preferably from about 10 to 20 or 12 to 18 amino acids in length, preferably 12, 15, or 18 amino acids in length), small organic molecule, polysaccharide, peptide, circular peptide, lipid, fatty acid, siRNA, polynucleotide, oligonucleotide, etc., to be tested for the capacity to directly or indirectly modulate angiogenesis and tumorigenesis. The test compound can be in the form of a library of test compounds, such as a combinatorial or randomized library that provides a sufficient range of diversity. Test compounds are optionally linked to a fusion partner, e.g., targeting compounds, rescue compounds, dimerization compounds, stabilizing compounds, addressable compounds, and other functional moieties. Conventionally, new chemical entities with useful properties are generated by identifying a test compound (called a “lead compound”) with some desirable property or activity, e.g., inhibiting activity, creating variants of the lead compound, and evaluating the property and activity of those variant compounds. Often, high throughput screening (HTS) methods are employed for such an analysis.

A “small organic molecule” refers to an organic molecule, either naturally occurring or synthetic, that has a molecular weight of more than about 50 daltons and less than about 2500 daltons, preferably less than about 2000 daltons, preferably between about 100 to about 1000 daltons, more preferably between about 200 to about 500 daltons.

“RNAi molecule” or an “siRNA” refers to a nucleic acid that forms a double stranded RNA, which double stranded RNA has the ability to reduce or inhibit expression of a gene or target gene when the siRNA expressed in the same cell as the gene or target gene. “siRNA” thus refers to the double stranded RNA formed by the complementary strands. The complementary portions of the siRNA that hybridize to form the double stranded molecule typically have substantial or complete identity. In one

embodiment, an siRNA refers to a nucleic acid that has substantial or complete identity to a target gene and forms a double stranded siRNA. The sequence of the siRNA can correspond to the full length target gene, or a subsequence thereof. Typically, the siRNA is at least about 15-50 nucleotides in length (e.g., each complementary sequence of the double stranded siRNA is 15-50 nucleotides in length, and the double stranded siRNA is about 15-50 base pairs in length, preferable about preferably about 20-30 base nucleotides, preferably about 20-25 nucleotides in length, e.g., 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleotides in length.

“Biological sample” include sections of tissues such as biopsy and autopsy samples, and frozen sections taken for histologic purposes. Such samples include blood, sputum, tissue, cultured cells, e.g., primary cultures, explants, and transformed cells, stool, urine, etc. A biological sample is typically obtained from a eukaryotic organism, most preferably a mammal such as a primate, e.g., chimpanzee or human; cow; dog; cat; a rodent, e.g., guinea pig, rat, mouse; rabbit; or a bird; reptile; or fish.

The terms “identical” or percent “identity,” in the context of two or more nucleic acids or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same (i.e., about 70% identity, preferably 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or higher identity over a specified region (e.g., SEQ ID NO:1 or 2), when compared and aligned for maximum correspondence over a comparison window or designated region) as measured using a BLAST or BLAST 2.0 sequence comparison algorithms with default parameters described below, or by manual alignment and visual inspection (*see, e.g.*, NCBI web site or the like). Such sequences are then said to be “substantially identical.” This definition also refers to, or may be applied to, the compliment of a test sequence. The definition also includes sequences that have deletions and/or additions, as well as those that have substitutions. As described below, the preferred algorithms can account for gaps and the like. Preferably, identity exists over a region that is at least about 25 amino acids or nucleotides in length, or more preferably over a region that is 50-100 amino acids or nucleotides in length.

For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. Preferably, default program parameters can be used, or alternative parameters

can be designated. The sequence comparison algorithm then calculates the percent sequence identities for the test sequences relative to the reference sequence, based on the program parameters.

A "comparison window", as used herein, includes reference to a segment  
 5 of any one of the number of contiguous positions selected from the group consisting of from 20 to 600, usually about 50 to about 200, more usually about 100 to about 150 in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. Methods of alignment of sequences for comparison are well-known in the art. Optimal alignment of  
 10 sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and  
 15 TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by manual alignment and visual inspection (*see, e.g., Current Protocols in Molecular Biology* (Ausubel *et al.*, eds. 1995 supplement)).

A preferred example of algorithm that is suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms,  
 20 which are described in Altschul *et al.*, *Nuc. Acids Res.* 25:3389-3402 (1997) and Altschul *et al.*, *J. Mol. Biol.* 215:403-410 (1990), respectively. BLAST and BLAST 2.0 are used, with the parameters described herein, to determine percent sequence identity for the nucleic acids and proteins of the invention. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. This  
 25 algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al., supra*). These initial neighborhood word hits act as seeds for initiating searches to  
 30 find longer HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the

cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The

5 BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, M=5, N=-4 and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength of 3, and expectation (E) of 10, and the BLOSUM62 scoring matrix (*see* Henikoff &  
10 Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915 (1989)) alignments (B) of 50, expectation (E) of 10, M=5, N=-4, and a comparison of both strands.

“Nucleic acid” refers to deoxyribonucleotides or ribonucleotides and polymers thereof in single- or double-stranded form. The term encompasses nucleic acids containing known nucleotide analogs or modified backbone residues or linkages, which  
15 are synthetic, naturally occurring, and non-naturally occurring, which have similar binding properties as the reference nucleic acid, and which are metabolized in a manner similar to the reference nucleotides. Examples of such analogs include, without limitation, phosphorothioates, phosphoramidates, methyl phosphonates, chiral-methyl phosphonates, 2-O-methyl ribonucleotides, peptide-nucleic acids (PNAs).

20 Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (e.g., degenerate codon substitutions) and complementary sequences, as well as the sequence explicitly indicated. Specifically, degenerate codon substitutions may be achieved by generating sequences in which the third position of one or more selected (or all) codons is substituted with mixed-  
25 base and/or deoxyinosine residues (Batzer *et al.*, *Nucleic Acid Res.* 19:5081 (1991); Ohtsuka *et al.*, *J. Biol. Chem.* 260:2605-2608 (1985); Rossolini *et al.*, *Mol. Cell. Probes* 8:91-98 (1994)). The term nucleic acid is used interchangeably with gene, cDNA, mRNA, oligonucleotide, and polynucleotide.

A particular nucleic acid sequence also implicitly encompasses “splice  
30 variants.” Similarly, a particular protein encoded by a nucleic acid implicitly encompasses any protein encoded by a splice variant of that nucleic acid. “Splice variants,” as the name suggests, are products of alternative splicing of a gene. After transcription, an initial nucleic acid transcript may be spliced such that different (alternate) nucleic acid splice products encode different polypeptides. Mechanisms for

the production of splice variants vary, but include alternate splicing of exons. Alternate polypeptides derived from the same nucleic acid by read-through transcription are also encompassed by this definition. Any products of a splicing reaction, including recombinant forms of the splice products, are included in this definition.

5           The terms “polypeptide,” “peptide” and “protein” are used interchangeably herein to refer to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers and non-naturally occurring amino acid polymer.

10           The term “amino acid” refers to naturally occurring and synthetic amino acids, as well as amino acid analogs and amino acid mimetics that function in a manner similar to the naturally occurring amino acids. Naturally occurring amino acids are those encoded by the genetic code, as well as those amino acids that are later modified, e.g., hydroxyproline,  $\gamma$ -carboxyglutamate, and O-phosphoserine. Amino acid analogs refers to  
15 compounds that have the same basic chemical structure as a naturally occurring amino acid, i.e., an  $\alpha$  carbon that is bound to a hydrogen, a carboxyl group, an amino group, and an R group, e.g., homoserine, norleucine, methionine sulfoxide, methionine methyl sulfonium. Such analogs have modified R groups (e.g., norleucine) or modified peptide backbones, but retain the same basic chemical structure as a naturally occurring amino  
20 acid. Amino acid mimetics refers to chemical compounds that have a structure that is different from the general chemical structure of an amino acid, but that functions in a manner similar to a naturally occurring amino acid.

Amino acids may be referred to herein by either their commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB  
25 Biochemical Nomenclature Commission. Nucleotides, likewise, may be referred to by their commonly accepted single-letter codes.

“Conservatively modified variants” applies to both amino acid and nucleic acid sequences. With respect to particular nucleic acid sequences, conservatively modified variants refers to those nucleic acids which encode identical or essentially  
30 identical amino acid sequences, or where the nucleic acid does not encode an amino acid sequence, to essentially identical sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given protein. For instance, the codons GCA, GCC, GCG and GCU all encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon, the codon can be altered

to any of the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are “silent variations,” which are one species of conservatively modified variations. Every nucleic acid sequence herein which encodes a polypeptide also describes every possible silent variation of the nucleic acid. One of skill  
 5 will recognize that each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine, and TGG, which is ordinarily the only codon for tryptophan) can be modified to yield a functionally identical molecule. Accordingly, each silent variation of a nucleic acid which encodes a polypeptide is implicit in each described sequence with respect to the expression product, but not with respect to actual probe  
 10 sequences.

As to amino acid sequences, one of skill will recognize that individual substitutions, deletions or additions to a nucleic acid, peptide, polypeptide, or protein sequence which alters, adds or deletes a single amino acid or a small percentage of amino acids in the encoded sequence is a “conservatively modified variant” where the alteration  
 15 results in the substitution of an amino acid with a chemically similar amino acid. Conservative substitution tables providing functionally similar amino acids are well known in the art. Such conservatively modified variants are in addition to and do not exclude polymorphic variants, interspecies homologs, and alleles of the invention.

The following eight groups each contain amino acids that are conservative  
 20 substitutions for one another: 1) Alanine (A), Glycine (G); 2) Aspartic acid (D), Glutamic acid (E); 3) Asparagine (N), Glutamine (Q); 4) Arginine (R), Lysine (K); 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W); 7) Serine (S), Threonine (T); and 8) Cysteine (C), Methionine (M) (*see, e.g., Creighton, Proteins* (1984)).

Macromolecular structures such as polypeptide structures can be described  
 25 in terms of various levels of organization. For a general discussion of this organization, *see, e.g., Alberts et al., Molecular Biology of the Cell* (3<sup>rd</sup> ed., 1994) and Cantor and Schimmel, *Biophysical Chemistry Part I: The Conformation of Biological Macromolecules* (1980). “Primary structure” refers to the amino acid sequence of a  
 30 particular peptide. “Secondary structure” refers to locally ordered, three dimensional structures within a polypeptide. These structures are commonly known as domains, e.g., transmembrane domains, pore domains, and cytoplasmic tail domains. Domains are portions of a polypeptide that form a compact unit of the polypeptide and are typically 15 to 350 amino acids long. Exemplary domains include domains with enzymatic activity,

e.g., ligand binding domains, etc. Typical domains are made up of sections of lesser organization such as stretches of  $\beta$ -sheet and  $\alpha$ -helices. "Tertiary structure" refers to the complete three dimensional structure of a polypeptide monomer. "Quaternary structure" refers to the three dimensional structure formed by the noncovalent association of independent tertiary units. Anisotropic terms are also known as energy terms.

A "label" or a "detectable moiety" is a composition detectable by spectroscopic, photochemical, biochemical, immunochemical, chemical, or other physical means. For example, useful labels include  $^{32}\text{P}$ , fluorescent dyes, electron-dense reagents, enzymes (e.g., as commonly used in an ELISA), biotin, digoxigenin, or haptens and proteins which can be made detectable, e.g., by incorporating a radiolabel into the peptide or used to detect antibodies specifically reactive with the peptide.

The term "recombinant" when used with reference, e.g., to a cell, or nucleic acid, protein, or vector, indicates that the cell, nucleic acid, protein or vector, has been modified by the introduction of a heterologous nucleic acid or protein or the alteration of a native nucleic acid or protein, or that the cell is derived from a cell so modified. Thus, for example, recombinant cells express genes that are not found within the native (non-recombinant) form of the cell or express native genes that are otherwise abnormally expressed, under expressed or not expressed at all.

The term "heterologous" when used with reference to portions of a nucleic acid indicates that the nucleic acid comprises two or more subsequences that are not found in the same relationship to each other in nature. For instance, the nucleic acid is typically recombinantly produced, having two or more sequences from unrelated genes arranged to make a new functional nucleic acid, e.g., a promoter from one source and a coding region from another source. Similarly, a heterologous protein indicates that the protein comprises two or more subsequences that are not found in the same relationship to each other in nature (e.g., a fusion protein).

The phrase "stringent hybridization conditions" refers to conditions under which a probe will hybridize to its target subsequence, typically in a complex mixture of nucleic acids, but to no other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures. An extensive guide to the hybridization of nucleic acids is found in Tijssen, *Techniques in Biochemistry and Molecular Biology--Hybridization with Nucleic Probes*, "Overview of principles of hybridization and the strategy of nucleic acid assays" (1993). Generally, stringent conditions are selected to be about 5-10°C lower

than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength pH. The  $T_m$  is the temperature (under defined ionic strength, pH, and nucleic concentration) at which 50% of the probes complementary to the target hybridize to the target sequence at equilibrium (as the target sequences are present in excess, at  $T_m$ , 50% of the probes are occupied at equilibrium). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. For selective or specific hybridization, a positive signal is at least two times background, preferably 10 times background hybridization. Exemplary stringent hybridization conditions can be as following: 50% formamide, 5x SSC, and 1% SDS, incubating at 42°C, or, 5x SSC, 1% SDS, incubating at 65°C, with wash in 0.2x SSC, and 0.1% SDS at 65°C.

Nucleic acids that do not hybridize to each other under stringent conditions are still substantially identical if the polypeptides which they encode are substantially identical. This occurs, for example, when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code. In such cases, the nucleic acids typically hybridize under moderately stringent hybridization conditions. Exemplary “moderately stringent hybridization conditions” include a hybridization in a buffer of 40% formamide, 1 M NaCl, 1% SDS at 37°C, and a wash in 1X SSC at 45°C. A positive hybridization is at least twice background. Those of ordinary skill will readily recognize that alternative hybridization and wash conditions can be utilized to provide conditions of similar stringency. Additional guidelines for determining hybridization parameters are provided in numerous reference, e.g., and *Current Protocols in Molecular Biology*, ed. Ausubel, *et al.*

For PCR, a temperature of about 36°C is typical for low stringency amplification, although annealing temperatures may vary between about 32°C and 48°C depending on primer length. For high stringency PCR amplification, a temperature of about 62°C is typical, although high stringency annealing temperatures can range from about 50°C to about 65°C, depending on the primer length and specificity. Typical cycle conditions for both high and low stringency amplifications include a denaturation phase of 90°C - 95°C for 30 sec - 2 min., an annealing phase lasting 30 sec. - 2 min., and an extension phase of about 72°C for 1 - 2 min. Protocols and guidelines for low and high stringency amplification reactions are provided, e.g., in Innis *et al.* (1990) *PCR Protocols, A Guide to Methods and Applications*, Academic Press, Inc. N.Y.).

“Antibody” refers to a polypeptide comprising a framework region from an immunoglobulin gene or fragments thereof that specifically binds and recognizes an antigen. The recognized immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon, and mu constant region genes, as well as the myriad  
 5 immunoglobulin variable region genes. Light chains are classified as either kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD and IgE, respectively. Typically, the antigen-binding region of an antibody will be most critical in specificity and affinity of binding.

10 An exemplary immunoglobulin (antibody) structural unit comprises a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one “light” (about 25 kD) and one “heavy” chain (about 50-70 kD). The N-terminus of each chain defines a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The terms variable light chain ( $V_L$ ) and  
 15 variable heavy chain ( $V_H$ ) refer to these light and heavy chains respectively.

Antibodies exist, e.g., as intact immunoglobulins or as a number of well-characterized fragments produced by digestion with various peptidases. Thus, for example, pepsin digests an antibody below the disulfide linkages in the hinge region to produce  $F(ab)'_2$ , a dimer of Fab which itself is a light chain joined to  $V_H$ - $C_{H1}$  by a  
 20 disulfide bond. The  $F(ab)'_2$  may be reduced under mild conditions to break the disulfide linkage in the hinge region, thereby converting the  $F(ab)'_2$  dimer into an Fab' monomer. The Fab' monomer is essentially Fab with part of the hinge region (*see Fundamental Immunology* (Paul ed., 3d ed. 1993). While various antibody fragments are defined in terms of the digestion of an intact antibody, one of skill will appreciate that such  
 25 fragments may be synthesized *de novo* either chemically or by using recombinant DNA methodology. Thus, the term antibody, as used herein, also includes antibody fragments either produced by the modification of whole antibodies, or those synthesized *de novo* using recombinant DNA methodologies (e.g., single chain Fv) or those identified using phage display libraries (*see, e.g., McCafferty et al., Nature* 348:552-554 (1990))

30 For preparation of antibodies, e.g., recombinant, monoclonal, or polyclonal antibodies, many technique known in the art can be used (*see, e.g., Kohler & Milstein, Nature* 256:495-497 (1975); Kozbor *et al., Immunology Today* 4: 72 (1983); Cole *et al.*, pp. 77-96 in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc. (1985); Coligan, *Current Protocols in Immunology* (1991); Harlow & Lane, *Antibodies, A*

*Laboratory Manual* (1988); and Goding, *Monoclonal Antibodies: Principles and Practice* (2d ed. 1986)). The genes encoding the heavy and light chains of an antibody of interest can be cloned from a cell, e.g., the genes encoding a monoclonal antibody can be cloned from a hybridoma and used to produce a recombinant monoclonal antibody. Gene libraries encoding heavy and light chains of monoclonal antibodies can also be made from hybridoma or plasma cells. Random combinations of the heavy and light chain gene products generate a large pool of antibodies with different antigenic specificity (*see, e.g.,* Kuby, *Immunology* (3<sup>rd</sup> ed. 1997)). Techniques for the production of single chain antibodies or recombinant antibodies (U.S. Patent 4,946,778, U.S. Patent No. 4,816,567) can be adapted to produce antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms such as other mammals, may be used to express humanized or human antibodies (*see, e.g.,* U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, Marks *et al.*, *Bio/Technology* 10:779-783 (1992); Lonberg *et al.*, *Nature* 368:856-859 (1994); Morrison, *Nature* 368:812-13 (1994); Fishwild *et al.*, *Nature Biotechnology* 14:845-51 (1996); Neuberger, *Nature Biotechnology* 14:826 (1996); and Lonberg & Huszar, *Intern. Rev. Immunol.* 13:65-93 (1995)). Alternatively, phage display technology can be used to identify antibodies and heteromeric Fab fragments that specifically bind to selected antigens (*see, e.g.,* McCafferty *et al.*, *Nature* 348:552-554 (1990); Marks *et al.*, *Biotechnology* 10:779-783 (1992)). Antibodies can also be made bispecific, i.e., able to recognize two different antigens (*see, e.g.,* WO 93/08829, Traunecker *et al.*, *EMBO J.* 10:3655-3659 (1991); and Suresh *et al.*, *Methods in Enzymology* 121:210 (1986)). Antibodies can also be heteroconjugates, e.g., two covalently joined antibodies, or immunotoxins (*see, e.g.,* U.S. Patent No. 4,676,980, WO 91/00360; WO 92/200373; and EP 03089).

Methods for humanizing or primatizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as import residues, which are typically taken from an import variable domain. Humanization can be essentially performed following the method of Winter and co-workers (*see, e.g.,* Jones *et al.*, *Nature* 321:522-525 (1986); Riechmann *et al.*, *Nature* 332:323-327 (1988); Verhoeven *et al.*, *Science* 239:1534-1536 (1988) and Presta, *Curr. Op. Struct. Biol.* 2:593-596 (1992)), by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such humanized antibodies are chimeric antibodies (U.S. Patent No. 4,816,567), wherein

substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

5           A “chimeric antibody” is an antibody molecule in which (a) the constant region, or a portion thereof, is altered, replaced or exchanged so that the antigen binding site (variable region) is linked to a constant region of a different or altered class, effector function and/or species, or an entirely different molecule which confers new properties to the chimeric antibody, e.g., an enzyme, toxin, hormone, growth factor, drug, etc.; or (b)  
10   the variable region, or a portion thereof, is altered, replaced or exchanged with a variable region having a different or altered antigen specificity.

          In one embodiment, the antibody is conjugated to an “effector” moiety. The effector moiety can be any number of molecules, including labeling moieties such as radioactive labels or fluorescent labels, or can be a therapeutic moiety. In one aspect the  
15   antibody modulates the activity of the protein.

          The phrase “specifically (or selectively) binds” to an antibody or “specifically (or selectively) immunoreactive with,” when referring to a protein or peptide, refers to a binding reaction that is determinative of the presence of the protein, often in a heterogeneous population of proteins and other biologics. Thus, under  
20   designated immunoassay conditions, the specified antibodies bind to a particular protein at least two times the background and more typically more than 10 to 100 times background. Specific binding to an antibody under such conditions requires an antibody that is selected for its specificity for a particular protein. For example, polyclonal antibodies, polymorphic variants, alleles, orthologs, and conservatively modified variants,  
25   or splice variants, or portions thereof, can be selected to obtain only those polyclonal antibodies that are specifically immunoreactive with angiogenesis proteins and not with other proteins. This selection may be achieved by subtracting out antibodies that cross-react with other molecules. A variety of immunoassay formats may be used to select antibodies specifically immunoreactive with a particular protein. For example, solid-  
30   phase ELISA immunoassays are routinely used to select antibodies specifically immunoreactive with a protein (*see, e.g., Harlow & Lane, Antibodies, A Laboratory Manual* (1988) for a description of immunoassay formats and conditions that can be used to determine specific immunoreactivity).

## Assays for proteins that modulate angiogenesis, tumorigenesis, and cellular proliferation

High throughput functional genomics assays can be used to identify modulators of angiogenesis and tumorigenesis. Such assays can monitor changes in cell surface marker expression,  $\alpha\beta$  integrin production, proliferation, and differentiation using either cell lines or primary cells. Typically, early passage or primary endothelial cells are contacted with a cDNA or a random peptide library (encoded by nucleic acids). The cDNA library can comprise sense, antisense, full length, and truncated cDNAs. The peptide library is encoded by nucleic acids. The effect of the cDNA or peptide library on the endothelial cells is then monitored, using an assay such as cell surface marker expression (e.g.,  $\alpha\beta$  integrin) or a phenotypic assay for angiogenesis such as migration towards an ECM (extracellular matrix) component (*see, e.g., Klemke et al., J. Cell Biol.* 4:961-972 (1998)) or endothelial cell tube formation assays, as well as other bioassays such as the chick CAM assay, the mouse corneal assay, haptotaxis assays, VEGF-R assays, co-culture tube formation assays, and assays measuring the effect of administering potential modulators on implanted tumors. The effect of the cDNA or peptide can be validated and distinguished from somatic mutations, using, e.g., regulatable expression of the nucleic acid such as expression from a tetracycline promoter. cDNAs and nucleic acids encoding peptides can be rescued using techniques known to those of skill in the art, e.g., using a sequence tags. *In vivo* assays for tumor growth, such as mouse xenograft models, can also be used.

Proteins interacting with the peptide or with the protein encoded by the cDNA can be isolated using a yeast two-hybrid system, mammalian two hybrid system, or phage display screen, etc. Targets so identified can be further used as bait in these assays to identify additional members of the angiogenesis pathway, which members are also targets for drug development (*see, e.g., Fields et al., Nature* 340:245 (1989); Vasavada et al., *Proc. Nat'l Acad. Sci. USA* 88:10686 (1991); Fearon et al., *Proc. Nat'l Acad. Sci. USA* 89:7958 (1992); Dang et al., *Mol. Cell. Biol.* 11:954 (1991); Chien et al., *Proc. Nat'l Acad. Sci. USA* 9578 (1991); and U.S. Patent Nos. 5,283,173, 5,667,973, 5,468,614, 5,525,490, and 5,637,463).

Suitable endothelial cell lines include human umbilical vein cells (*see, e.g., Jaffe et al., J. Clin. Invest.* 52:2745-2754 (1973)); human adult dermal capillary-derived cells (*see, e.g., Davison et al., In Vitro* 19:937-945 (1983)); human adipose capillary derived cells (*see, e.g., Kern et al., J. Clin Invest.* 71:1822-1829 (1983); bovine aorta (*see,*

*e.g.*, Booyse *et al.*, *Thromb. Diathes. Aemorrh.* 34:825-839 (1975); and rat brain capillary derived cells (*see, e.g.*, Bowman *et al.*, *In Vitro* 17:353-362 (1981)). For culture of endothelial cell lines, explants, and primary cells, *see* Freshney *et al.*, *Culture of Animal Cells* (3<sup>rd</sup> ed. 1994). Suitable angiogenesis cell surface markers include

5     $\alpha$ v $\beta$ 3 integrin (*see, e.g.*, Elicerir & Cheresch, *Cancer J. Sci. Am.* 6 Supp. 3:S245-249 (2000), Maeshima *et al.*, *J. Biol. Chem.* (June 8, 2001)).

Cell surface markers such as  $\alpha$ v $\beta$ 3 can be assayed using fluorescently labeled antibodies and FACS. Cell proliferation can be measured using <sup>3</sup>H-thymidine or dye inclusion. Angiogenesis or tumorigenesis phenotype is measured by loss of

10    phenotype observation.

cDNA libraries are made from any suitable source, preferably from endothelial cells. Libraries encoding random peptides are made according to techniques well known to those of skill in the art (*see, e.g.*, U.S. Patent No. 6,153,380, 6,114,111, and 6,180,343). Any suitable vector can be used for the cDNA and peptide libraries,

15    including, *e.g.*, retroviral vectors.

In a preferred embodiment, target proteins that modulate angiogenesis or tumorigenesis are identified using a high throughput cell based assay (using a microtiter plate format) and FACS screening for  $\alpha$ v $\beta$ 3 cell surface expression. cDNA libraries are made which include, *e.g.*, sense, antisense, full length, and truncated cDNAs. The

20    cDNAs are cloned into a retroviral vector. Endothelial cells are infected with the library, cultured for a short effector phase and then the cells with reduced  $\alpha$ v $\beta$ 3 surface levels are enriched by antibody staining and magnetic cell sorting. The enriched cell population is then sorted into microtiter plates using fluorescent antibodies and FACS. Resultant cell colonies are analyzed by immunofluorescence for reduced  $\alpha$ v $\beta$ 3 surface levels. Selected

25    colonies are infected with wild type MMLV virus to rescue the proviral vector. The infectious supernatant is used to infect endothelial cells, which are subsequently analyzed for  $\alpha$ v $\beta$ 3 levels by FACS. The cDNA is isolated and sequenced to determine if it represents a wild type or mutated cDNA, *e.g.*, whether the cDNA represents a negative transdominant mutant. Optionally, a marker such as GFP can be used to select for

30    retrovirally infected cells.

### **Isolation of nucleic acids**

This invention relies on routine techniques in the field of recombinant genetics. Basic texts disclosing the general methods of use in this invention include

Sambrook *et al.*, *Molecular Cloning, A Laboratory Manual* (2nd ed. 1989); Kriegler, *Gene Transfer and Expression: A Laboratory Manual* (1990); and *Current Protocols in Molecular Biology* (Ausubel *et al.*, eds., 1994)).

Nucleic acids, polymorphic variants, orthologs, and alleles can be isolated using nucleic acid probes and oligonucleotides under stringent hybridization conditions, by screening libraries. Alternatively, expression libraries can be used to clone angiogenesis proteins, polymorphic variants, orthologs, and alleles by detecting expressed homologs immunologically with antisera or purified antibodies made against human angiogenesis proteins or portions thereof.

To make a cDNA library, one should choose a source that is rich in the desired RNA, e.g., endothelial cells. The mRNA is then made into cDNA using reverse transcriptase, ligated into a recombinant vector, and transfected into a recombinant host for propagation, screening and cloning. Methods for making and screening cDNA libraries are well known (*see, e.g.*, Gubler & Hoffman, *Gene* 25:263-269 (1983); Sambrook *et al.*, *supra*; Ausubel *et al.*, *supra*).

For a genomic library, the DNA is extracted from the tissue and either mechanically sheared or enzymatically digested to yield fragments of about 12-20 kb. The fragments are then separated by gradient centrifugation from undesired sizes and are constructed in bacteriophage lambda vectors. These vectors and phage are packaged *in vitro*. Recombinant phage are analyzed by plaque hybridization as described in Benton & Davis, *Science* 196:180-182 (1977). Colony hybridization is carried out as generally described in Grunstein *et al.*, *Proc. Natl. Acad. Sci. USA.*, 72:3961-3965 (1975).

An alternative method of isolating nucleic acids and orthologs, alleles, mutants, polymorphic variants, and conservatively modified variants combines the use of synthetic oligonucleotide primers and amplification of an RNA or DNA template (*see* U.S. Patents 4,683,195 and 4,683,202; *PCR Protocols: A Guide to Methods and Applications* (Innis *et al.*, eds, 1990)). Methods such as polymerase chain reaction (PCR) and ligase chain reaction (LCR) can be used to amplify nucleic acid sequences directly from mRNA, from cDNA, from genomic libraries or cDNA libraries. Degenerate oligonucleotides can be designed to amplify homologs using the sequences provided herein. Restriction endonuclease sites can be incorporated into the primers. Polymerase chain reaction or other *in vitro* amplification methods may also be useful, for example, to clone nucleic acid sequences that code for proteins to be expressed, to make nucleic acids to use as probes for detecting the presence of angiogenesis protein encoding mRNA in

physiological samples, for nucleic acid sequencing, or for other purposes. Genes amplified by the PCR reaction can be purified from agarose gels and cloned into an appropriate vector.

Gene expression can also be analyzed by techniques known in the art, e.g., reverse transcription and amplification of mRNA, isolation of total RNA or poly A<sup>+</sup> RNA, northern blotting, dot blotting, *in situ* hybridization, RNase protection, high density polynucleotide array technology, e.g., and the like.

Nucleic acids can be used with high density oligonucleotide array technology (e.g., GeneChip<sup>TM</sup>) to identify angiogenesis protein, orthologs, alleles, conservatively modified variants, and polymorphic variants in this invention. In the case where the homologs being identified are linked to a known disease state, they can be used with GeneChip<sup>TM</sup> as a diagnostic tool in detecting the disease in a biological sample, *see, e.g.,* Gunthand *et al.*, *AIDS Res. Hum. Retroviruses* 14: 869-876 (1998); Kozal *et al.*, *Nat. Med.* 2:753-759 (1996); Matson *et al.*, *Anal. Biochem.* 224:110-106 (1995); Lockhart *et al.*, *Nat. Biotechnol.* 14:1675-1680 (1996); Gingeras *et al.*, *Genome Res.* 8:435-448 (1998); Hacia *et al.*, *Nucleic Acids Res.* 26:3865-3866 (1998).

The gene is typically cloned into intermediate vectors before transformation into prokaryotic or eukaryotic cells for replication and/or expression. These intermediate vectors are typically prokaryote vectors, e.g., plasmids, or shuttle vectors.

### **Expression in prokaryotes and eukaryotes**

To obtain high level expression of a cloned gene, such as those cDNAs encoding an angiogenesis protein, one typically subclones the desired nucleic acid into an expression vector that contains a strong promoter to direct transcription, a transcription/translation terminator, and if for a nucleic acid encoding a protein, a ribosome binding site for translational initiation. Suitable bacterial promoters are well known in the art and described, e.g., in Sambrook *et al.*, and Ausubel *et al.*, *supra*. Bacterial expression systems for expressing the protein are available in, e.g., *E. coli*, *Bacillus sp.*, and *Salmonella* (Palva *et al.*, *Gene* 22:229-235 (1983); Mosbach *et al.*, *Nature* 302:543-545 (1983). Kits for such expression systems are commercially available. Eukaryotic expression systems for mammalian cells, yeast, and insect cells are well known in the art and are also commercially available.

Selection of the promoter used to direct expression of a heterologous nucleic acid depends on the particular application. The promoter is preferably positioned about the same distance from the heterologous transcription start site as it is from the transcription start site in its natural setting. As is known in the art, however, some variation in this distance can be accommodated without loss of promoter function.

In addition to the promoter, the expression vector typically contains a transcription unit or expression cassette that contains all the additional elements required for the expression of the nucleic acid in host cells. A typical expression cassette thus contains a promoter operably linked to the nucleic acid sequence and signals required for efficient polyadenylation of the transcript, ribosome binding sites, and translation termination. Additional elements of the cassette may include enhancers and, if genomic DNA is used as the structural gene, introns with functional splice donor and acceptor sites.

In addition to a promoter sequence, the expression cassette should also contain a transcription termination region downstream of the structural gene to provide for efficient termination. The termination region may be obtained from the same gene as the promoter sequence or may be obtained from different genes.

The particular expression vector used to transport the genetic information into the cell is not particularly critical. Any of the conventional vectors used for expression in eukaryotic or prokaryotic cells may be used. Standard bacterial expression vectors include plasmids such as pBR322 based plasmids, pSKF, pET23D, and fusion expression systems such as MBP, GST, and LacZ. Epitope tags can also be added to recombinant proteins to provide convenient methods of isolation, e.g., c-myc.

Expression vectors containing regulatory elements from eukaryotic viruses are typically used in eukaryotic expression vectors, e.g., SV40 vectors, papilloma virus vectors, retroviral vectors, and vectors derived from Epstein-Barr virus. Other exemplary eukaryotic vectors include pMSG, pAV009/A<sup>+</sup>, pMTO10/A<sup>+</sup>, pMAMneo-5, baculovirus pDSVE, and any other vector allowing expression of proteins under the direction of the CMV promoter, SV40 early promoter, SV40 later promoter, metallothionein promoter, murine mammary tumor virus promoter, Rous sarcoma virus promoter, polyhedrin promoter, or other promoters shown effective for expression in eukaryotic cells.

Expression of proteins from eukaryotic vectors can be also be regulated using inducible promoters. With inducible promoters, expression levels are tied to the concentration of inducing agents, such as tetracycline or ecdysone, by the incorporation

of response elements for these agents into the promoter. Generally, high level expression is obtained from inducible promoters only in the presence of the inducing agent; basal expression levels are minimal. Inducible expression vectors are often chosen if expression of the protein of interest is detrimental to eukaryotic cells.

5                   In one embodiment, the vectors of the invention have a regulatable promoter, e.g., tet-regulated systems and the RU-486 system (*see, e.g.,* Gossen & Bujard, *Proc. Nat'l Acad. Sci. USA* 89:5547 (1992); Oligino *et al.*, *Gene Ther.* 5:491-496 (1998); Wang *et al.*, *Gene Ther.* 4:432-441 (1997); Neering *et al.*, *Blood* 88:1147-1155 (1996); and Rendahl *et al.*, *Nat. Biotechnol.* 16:757-761 (1998)). These impart small molecule  
10 control on the expression of the candidate target nucleic acids. This beneficial feature can be used to determine that a desired phenotype is caused by a transfected cDNA rather than a somatic mutation.

                  Some expression systems have markers that provide gene amplification such as thymidine kinase and dihydrofolate reductase. Alternatively, high yield  
15 expression systems not involving gene amplification are also suitable, such as using a baculovirus vector in insect cells, with a protein encoding sequence under the direction of the polyhedrin promoter or other strong baculovirus promoters.

                  The elements that are typically included in expression vectors also include a replicon that functions in *E. coli*, a gene encoding antibiotic resistance to permit  
20 selection of bacteria that harbor recombinant plasmids, and unique restriction sites in nonessential regions of the plasmid to allow insertion of eukaryotic sequences. The particular antibiotic resistance gene chosen is not critical, any of the many resistance genes known in the art are suitable. The prokaryotic sequences are preferably chosen such that they do not interfere with the replication of the DNA in eukaryotic cells, if  
25 necessary.

                  Standard transfection methods are used to produce bacterial, mammalian, yeast or insect cell lines that express large quantities of protein, which are then purified using standard techniques (*see, e.g.,* Colley *et al.*, *J. Biol. Chem.* 264:17619-17622 (1989); *Guide to Protein Purification*, in *Methods in Enzymology*, vol. 182 (Deutscher, ed., 1990)). Transformation of eukaryotic and prokaryotic cells are performed according  
30 to standard techniques (*see, e.g.,* Morrison, *J. Bact.* 132:349-351 (1977); Clark-Curtiss & Curtiss, *Methods in Enzymology* 101:347-362 (Wu *et al.*, eds, 1983).

                  Any of the well-known procedures for introducing foreign nucleotide sequences into host cells may be used. These include the use of calcium phosphate

transfection, polybrene, protoplast fusion, electroporation, biolistics, liposomes, microinjection, plasma vectors, viral vectors and any of the other well known methods for introducing cloned genomic DNA, cDNA, synthetic DNA or other foreign genetic material into a host cell (*see, e.g., Sambrook et al., supra*). It is only necessary that the particular genetic engineering procedure used be capable of successfully introducing at least one gene into the host cell capable of expressing the protein of choice.

After the expression vector is introduced into the cells, the transfected cells are cultured under conditions favoring expression of the protein of choice, which is recovered from the culture using standard techniques identified below.

### **Purification of polypeptides**

Either naturally occurring or recombinant protein can be purified for use in functional assays. Naturally occurring protein can be purified, e.g., from human tissue. Recombinant protein can be purified from any suitable expression system.

The protein may be purified to substantial purity by standard techniques, including selective precipitation with such substances as ammonium sulfate; column chromatography, immunopurification methods, and others (*see, e.g., Scopes, Protein Purification: Principles and Practice* (1982); U.S. Patent No. 4,673,641; Ausubel *et al., supra*; and Sambrook *et al., supra*).

A number of procedures can be employed when recombinant protein is being purified. For example, proteins having established molecular adhesion properties can be reversibly fused to the protein. With the appropriate ligand, angiogenesis protein can be selectively adsorbed to a purification column and then freed from the column in a relatively pure form. The fused protein is then removed by enzymatic activity. Finally, protein could be purified using immunoaffinity columns.

#### *A. Purification of protein from recombinant bacteria*

Recombinant proteins are expressed by transformed bacteria in large amounts, typically after promoter induction; but expression can be constitutive. Promoter induction with IPTG is one example of an inducible promoter system. Bacteria are grown according to standard procedures in the art. Fresh or frozen bacteria cells are used for isolation of protein.

Proteins expressed in bacteria may form insoluble aggregates ("inclusion bodies"). Several protocols are suitable for purification of protein inclusion bodies. For

example, purification of inclusion bodies typically involves the extraction, separation and/or purification of inclusion bodies by disruption of bacterial cells, e.g., by incubation in a buffer of 50 mM TRIS/HCL pH 7.5, 50 mM NaCl, 5 mM MgCl<sub>2</sub>, 1 mM DTT, 0.1 mM ATP, and 1 mM PMSF. The cell suspension can be lysed using 2-3 passages  
5 through a French Press, homogenized using a Polytron (Brinkman Instruments) or sonicated on ice. Alternate methods of lysing bacteria are apparent to those of skill in the art (*see, e.g., Sambrook et al., supra; Ausubel et al., supra*).

If necessary, the inclusion bodies are solubilized, and the lysed cell suspension is typically centrifuged to remove unwanted insoluble matter. Proteins that  
10 formed the inclusion bodies may be renatured by dilution or dialysis with a compatible buffer. Suitable solvents include, but are not limited to urea (from about 4 M to about 8 M), formamide (at least about 80%, volume/volume basis), and guanidine hydrochloride (from about 4 M to about 8 M). Some solvents which are capable of solubilizing aggregate-forming proteins, for example SDS (sodium dodecyl sulfate), 70% formic acid,  
15 are inappropriate for use in this procedure due to the possibility of irreversible denaturation of the proteins, accompanied by a lack of immunogenicity and/or activity. Although guanidine hydrochloride and similar agents are denaturants, this denaturation is not irreversible and renaturation may occur upon removal (by dialysis, for example) or dilution of the denaturant, allowing re-formation of immunologically and/or biologically  
20 active protein. Other suitable buffers are known to those skilled in the art. Human proteins are separated from other bacterial proteins by standard separation techniques, e.g., with Ni-NTA agarose resin.

Alternatively, it is possible to purify protein from bacteria periplasm. After lysis of the bacteria, the periplasmic fraction of the bacteria can be isolated by cold  
25 osmotic shock in addition to other methods known to skill in the art. To isolate recombinant proteins from the periplasm, the bacterial cells are centrifuged to form a pellet. The pellet is resuspended in a buffer containing 20% sucrose. To lyse the cells, the bacteria are centrifuged and the pellet is resuspended in ice-cold 5 mM MgSO<sub>4</sub> and kept in an ice bath for approximately 10 minutes. The cell suspension is centrifuged and  
30 the supernatant decanted and saved. The recombinant proteins present in the supernatant can be separated from the host proteins by standard separation techniques well known to those of skill in the art.

*B. Standard protein separation techniques for purifying proteins*

Solubility fractionation

Often as an initial step, particularly if the protein mixture is complex, an initial salt fractionation can separate many of the unwanted host cell proteins (or proteins derived from the cell culture media) from the recombinant protein of interest. The preferred salt is ammonium sulfate. Ammonium sulfate precipitates proteins by effectively reducing the amount of water in the protein mixture. Proteins then precipitate on the basis of their solubility. The more hydrophobic a protein is, the more likely it is to precipitate at lower ammonium sulfate concentrations. A typical protocol includes adding saturated ammonium sulfate to a protein solution so that the resultant ammonium sulfate concentration is between 20-30%. This concentration will precipitate the most hydrophobic of proteins. The precipitate is then discarded (unless the protein of interest is hydrophobic) and ammonium sulfate is added to the supernatant to a concentration known to precipitate the protein of interest. The precipitate is then solubilized in buffer and the excess salt removed if necessary, either through dialysis or diafiltration. Other methods that rely on solubility of proteins, such as cold ethanol precipitation, are well known to those of skill in the art and can be used to fractionate complex protein mixtures.

Size differential filtration

The molecular weight of the protein can be used to isolate it from proteins of greater and lesser size using ultrafiltration through membranes of different pore size (for example, Amicon or Millipore membranes). As a first step, the protein mixture is ultrafiltered through a membrane with a pore size that has a lower molecular weight cut-off than the molecular weight of the protein of interest. The retentate of the ultrafiltration is then ultrafiltered against a membrane with a molecular cut off greater than the molecular weight of the protein of interest. The recombinant protein will pass through the membrane into the filtrate. The filtrate can then be chromatographed as described below.

Column chromatography

The protein can also be separated from other proteins on the basis of its size, net surface charge, hydrophobicity, and affinity for ligands. In addition, antibodies raised against proteins can be conjugated to column matrices and the proteins immunopurified. All of these methods are well known in the art. It will be apparent to

one of skill that chromatographic techniques can be performed at any scale and using equipment from many different manufacturers (e.g., Pharmacia Biotech).

### Assays for modulators of angiogenesis or tumorigenesis proteins

#### 5           A.     Assays

Modulation of an angiogenesis protein, and corresponding modulation of angiogenesis or tumorigenesis, can be assessed using a variety of *in vitro* and *in vivo* assays, including high throughput ligand binding and cell based assays, as described herein. Such assays can be used to test for inhibitors and activators of the angiogenesis protein, and, consequently, inhibitors and activators of angiogenesis. Such modulators of the angiogenesis protein are useful for treating angiogenesis and tumorigenesis disorders. Modulators of the angiogenesis protein are tested using either recombinant or naturally occurring protein, preferably human protein.

Measurement of an angiogenic or tumorigenic or loss-of-angiogenesis or tumorigenesis phenotype on the protein or cell expressing the protein, either recombinant or naturally occurring, can be performed using a variety of assays, *in vitro*, *in vivo*, and *ex vivo*. For example, recombinant or naturally occurring protein can be used *in vitro*, in a ligand binding or enzymatic function assay. Protein present in a cellular extract can also be used in *in vitro* assays. Cell- and animal-based *in vivo* assays can also be used to assay for angiogenesis modulators. Any suitable physical, chemical, or phenotypic change that affects activity or binding can be used to assess the influence of a test compound on the polypeptide of this invention. When the functional effects are determined using intact cells or animals, one can also measure a variety of effects such as, in the case of angiogenesis associated with tumors, tumor growth, neovascularization, endothelial tube formation, cell surface markers such as  $\alpha\beta3$ , hormone release, transcriptional changes to both known and uncharacterized genetic markers (e.g., northern blots), changes in cell metabolism such as cell growth or pH changes, and changes in intracellular second messengers such as cGMP. In one embodiment, measurement of  $\alpha\beta3$  integrin cell surface expression and FACS sorting is used to identify modulators of angiogenesis.

30

#### *In vitro* assays

Assays to identify compounds with angiogenesis or tumorigenesis modulating activity, e.g., anti-angiogenic or anti-tumorigenic activity, can be performed *in vitro*, e.g., binding assays. Purified recombinant or naturally occurring protein can be

used in the *in vitro* methods of the invention. In addition to purified protein, the recombinant or naturally occurring protein can be part of a cellular lysate. As described below, the assay can be either solid state or soluble. Preferably, the protein is bound to a solid support, either covalently or non-covalently. Often, the *in vitro* assays of the invention are ligand binding or ligand affinity assays, either non-competitive or competitive. Other *in vitro* assays include measuring changes in spectroscopic (e.g., fluorescence, absorbance, refractive index), hydrodynamic (e.g., shape), chromatographic, or solubility properties for the protein.

In one embodiment, a high throughput binding assay is performed in which the protein or chimera comprising a fragment thereof is contacted with a potential modulator and incubated for a suitable amount of time. In one embodiment, the potential modulator is bound to a solid support, and the protein is added. In another embodiment, the protein is bound to a solid support. A wide variety of modulators can be used, as described below, including small organic molecules, peptides, and antibodies. A wide variety of assays can be used to identify angiogenesis-modulator binding, including labeled protein-protein binding assays, electrophoretic mobility shifts, immunoassays, and the like. In some cases, the binding of the candidate modulator is determined through the use of competitive binding assays, where interference with binding of a known ligand is measured in the presence of a potential modulator. Often, either the potential modulator or the known ligand is labeled.

#### Cell-based *in vivo* assays

In another embodiment, the protein is expressed in a cell, and functional, e.g., physical and chemical or phenotypic, changes are assayed to identify angiogenesis or tumorigenesis modulators, preferably anti-angiogenesis or anti-tumorigenesis compounds. Cells expressing angiogenesis proteins can also be used in binding assays or enzymatic assays. Any suitable functional effect can be measured, as described herein. For example, ligand binding, cell surface marker expression, cellular proliferation, VEGF-R assays, co-culture assays for tube formation, and cell migration assays are all suitable assays to identify potential modulators using a cell based system. Suitable cells for such cell based assays include both primary endothelial cells and cell lines, as described herein. The protein can be naturally occurring or recombinant.

As described above, in one embodiment, loss-of angiogenesis or tumorigenesis phenotype is measured by contacting endothelial cells comprising an

angiogenesis target with a potential modulator. Modulation of angiogenesis or tumorigenesis is identified by screening for cell surface marker expression, e.g.,  $\alpha v \beta 3$  integrin expression levels, using fluorescent antibodies and FACS sorting.

5 In another embodiment, cellular proliferation can be measured using  $^3\text{H}$ -thymidine incorporation or dye inclusion.

In another embodiment, cellular polypeptide levels are determined by measuring the level of protein or mRNA. The level of protein or proteins are measured using immunoassays such as western blotting, ELISA and the like with an antibody that selectively binds to the polypeptide or a fragment thereof. For measurement of mRNA,  
10 amplification, e.g., using PCR, LCR, or hybridization assays, e.g., northern hybridization, RNase protection, dot blotting, are preferred. The level of protein or mRNA is detected using directly or indirectly labeled detection agents, e.g., fluorescently or radioactively labeled nucleic acids, radioactively or enzymatically labeled antibodies, and the like, as described herein.

15 Alternatively, protein expression can be measured using a reporter gene system. Such a system can be devised using an angiogenesis protein promoter operably linked to a reporter gene such as chloramphenicol acetyltransferase, firefly luciferase, bacterial luciferase,  $\beta$ -galactosidase and alkaline phosphatase. Furthermore, the protein of interest can be used as an indirect reporter via attachment to a second reporter such as  
20 red or green fluorescent protein (*see, e.g.,* Mistili & Spector, *Nature Biotechnology* 15:961-964 (1997)). The reporter construct is typically transfected into a cell. After treatment with a potential modulator, the amount of reporter gene transcription, translation, or activity is measured according to standard techniques known to those of skill in the art.

25 A variety of phenotypic angiogenesis or tumorigenesis assays are known to those of skill in the art. Various models have been employed to evaluate angiogenesis (e.g., Croix *et al.*, *Science* 289:1197-1202 (2000) and Kahn *et al.*, *Amer. J. Pathol.* 156:1887-1900). Assessment of angiogenesis or tumorigenesis in the presence of a potential modulator can be performed using cell-culture-based assays, e.g., endothelial  
30 cell tube formation assays and haptotaxis assays, as well as other animal based bioassays such as the chick CAM assay, the mouse corneal assay, and assays measuring the effect of administering potential modulators on implanted tumors.

For determination of cellular proliferation, any suitable functional effect can be measured, as described herein. For example, cellular morphology (e.g., cell volume, nuclear volume, cell perimeter, and nuclear perimeter), ligand binding, kinase activity, apoptosis, cell surface marker expression, cellular proliferation, GFP positivity and dye dilution assays (e.g., cell tracker assays with dyes that bind to cell membranes), DNA synthesis assays (e.g.,  $^3\text{H}$ -thymidine and fluorescent DNA-binding dyes such as BrdU or Hoescht dye with FACS analysis),  $G_0/G_1$  cell cycle arrest, are all suitable assays to identify potential modulators using a cell based system. Suitable cells for such cell based assays include both primary cancer or tumor cells and cell lines, as described herein, e.g., A549 (lung), MCF7 (breast, p53 wild-type), H1299 (lung, p53 null), HeLa (cervical), PC3 (prostate, p53 mutant), MDA-MB-231 (breast, p53 wild-type). Cancer cell lines can be p53 mutant, p53 null, or express wild type p53.

#### Animal models

A number of animal based assays for angiogenesis or tumorigenesis phenotypes are known to those of skill in the art and can be used to assay for modulators of angiogenesis. For example, the chick CAM assay is described by O'Reilly, *et al.* Cell 79: 315-328 (1994). Briefly, 3 day old chicken embryos with intact yolks are separated from the egg and placed in a petri dish. After 3 days of incubation, a methylcellulose disc containing the protein to be tested is applied to the CAM of individual embryos. After about 48 hours of incubation, the embryos and CAMs are observed to determine whether endothelial growth has been inhibited.

The mouse corneal assay involves implanting a growth factor-containing pellet, along with another pellet containing the suspected endothelial growth inhibitor, in the cornea of a mouse and observing the pattern of capillaries that are elaborated in the cornea.

Angiogenesis can also be measured by determining the extent of neovascularization of a tumor. For example, carcinoma cells can be subcutaneously inoculated into athymic or nude mice or SCID mice and tumor growth then monitored. Immunoassays using endothelial cell-specific antibodies are typically used to stain for vascularization of tumor and the number of vessels in the tumor.

As described above, animal models of angiogenesis find use in screening for modulators of angiogenesis and tumorigenesis. Similarly, transgenic animal technology including gene knockout technology, for example as a result of homologous

recombination with an appropriate gene targeting vector, or gene overexpression, will result in the absence or increased expression of the protein. The same technology can also be applied to make knock-out cells. When desired, tissue-specific expression or knockout of the protein may be necessary. Transgenic animals generated by such methods find use as animal models of angiogenesis and are additionally useful in screening for modulators of angiogenesis and tumorigenesis.

Knock-out cells and transgenic mice can be made by insertion of a marker gene or other heterologous gene into the endogenous gene site in the mouse genome via homologous recombination. Such mice can also be made by substituting the endogenous gene with a mutated version of the gene, or by mutating the endogenous gene, e.g., by exposure to carcinogens.

A DNA construct is introduced into the nuclei of embryonic stem cells. Cells containing the newly engineered genetic lesion are injected into a host mouse embryo, which is re-implanted into a recipient female. Some of these embryos develop into chimeric mice that possess germ cells partially derived from the mutant cell line. Therefore, by breeding the chimeric mice it is possible to obtain a new line of mice containing the introduced genetic lesion (*see, e.g., Capecchi et al., Science* 244:1288 (1989)). Chimeric targeted mice can be derived according to Hogan *et al., Manipulating the Mouse Embryo: A Laboratory Manual*, Cold Spring Harbor Laboratory (1988) and *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, Robertson, ed., IRL Press, Washington, D.C., (1987).

### B. Modulators

The compounds tested as modulators of the angiogenesis protein can be any small organic molecule, or a biological entity, such as a protein, e.g., an antibody or peptide, a sugar, a nucleic acid, e.g., an antisense oligonucleotide, RNAi molecule, or a ribozyme, or a lipid. Alternatively, modulators can be genetically altered versions of an angiogenesis protein. Typically, test compounds will be small organic molecules, peptides, lipids, and lipid analogs.

Essentially any chemical compound can be used as a potential modulator or ligand in the assays of the invention, although most often compounds can be dissolved in aqueous or organic (especially DMSO-based) solutions are used. The assays are designed to screen large chemical libraries by automating the assay steps and providing compounds from any convenient source to assays, which are typically run in parallel

(e.g., in microtiter formats on microtiter plates in robotic assays). It will be appreciated that there are many suppliers of chemical compounds, including Sigma (St. Louis, MO), Aldrich (St. Louis, MO), Sigma-Aldrich (St. Louis, MO), Fluka Chemika-Biochemica Analytika (Buchs Switzerland) and the like.

5                   In one preferred embodiment, high throughput screening methods involve providing a combinatorial small organic molecule or peptide library containing a large number of potential therapeutic compounds (potential modulator or ligand compounds). Such "combinatorial chemical libraries" or "ligand libraries" are then screened in one or more assays, as described herein, to identify those library members (particular chemical  
10 species or subclasses) that display a desired characteristic activity. The compounds thus identified can serve as conventional "lead compounds" or can themselves be used as potential or actual therapeutics.

                  A combinatorial chemical library is a collection of diverse chemical compounds generated by either chemical synthesis or biological synthesis, by combining  
15 a number of chemical "building blocks" such as reagents. For example, a linear combinatorial chemical library such as a polypeptide library is formed by combining a set of chemical building blocks (amino acids) in every possible way for a given compound length (i.e., the number of amino acids in a polypeptide compound). Millions of chemical compounds can be synthesized through such combinatorial mixing of chemical building  
20 blocks.

                  Preparation and screening of combinatorial chemical libraries is well known to those of skill in the art. Such combinatorial chemical libraries include, but are not limited to, peptide libraries (*see, e.g.*, U.S. Patent 5,010,175, Furka, *Int. J. Pept. Prot. Res.* 37:487-493 (1991) and Houghton *et al.*, *Nature* 354:84-88 (1991)). Other  
25 chemistries for generating chemical diversity libraries can also be used. Such chemistries include, but are not limited to: peptoids (e.g., PCT Publication No. WO 91/19735), encoded peptides (e.g., PCT Publication No. WO 93/20242), random bio-oligomers (e.g., PCT Publication No. WO 92/00091), benzodiazepines (e.g., U.S. Pat. No. 5,288,514), diversomers such as hydantoins, benzodiazepines and dipeptides (Hobbs *et al.*, *Proc. Nat.*  
30 *Acad. Sci. USA* 90:6909-6913 (1993)), vinylogous polypeptides (Hagihara *et al.*, *J. Amer. Chem. Soc.* 114:6568 (1992)), nonpeptidal peptidomimetics with glucose scaffolding (Hirschmann *et al.*, *J. Amer. Chem. Soc.* 114:9217-9218 (1992)), analogous organic syntheses of small compound libraries (Chen *et al.*, *J. Amer. Chem. Soc.* 116:2661 (1994)), oligocarbamates (Cho *et al.*, *Science* 261:1303 (1993)), and/or peptidyl

phosphonates (Campbell *et al.*, *J. Org. Chem.* 59:658 (1994)), nucleic acid libraries (*see* Ausubel, Berger and Sambrook, all *supra*), peptide nucleic acid libraries (*see, e.g.*, U.S. Patent 5,539,083), antibody libraries (*see, e.g.*, Vaughn *et al.*, *Nature Biotechnology*, 14(3):309-314 (1996) and PCT/US96/10287), carbohydrate libraries (*see, e.g.*, Liang *et al.*, *Science*, 274:1520-1522 (1996) and U.S. Patent 5,593,853), small organic molecule libraries (*see, e.g.*, benzodiazepines, Baum C&EN, Jan 18, page 33 (1993); isoprenoids, U.S. Patent 5,569,588; thiazolidinones and metathiazanones, U.S. Patent 5,549,974; pyrrolidines, U.S. Patents 5,525,735 and 5,519,134; morpholino compounds, U.S. Patent 5,506,337; benzodiazepines, 5,288,514, and the like).

Devices for the preparation of combinatorial libraries are commercially available (*see, e.g.*, 357 MPS, 390 MPS, Advanced Chem Tech, Louisville KY, Symphony, Rainin, Woburn, MA, 433A Applied Biosystems, Foster City, CA, 9050 Plus, Millipore, Bedford, MA). In addition, numerous combinatorial libraries are themselves commercially available (*see, e.g.*, ComGenex, Princeton, N.J., Asinex, Moscow, Ru, Tripos, Inc., St. Louis, MO, ChemStar, Ltd, Moscow, RU, 3D Pharmaceuticals, Exton, PA, Martek Biosciences, Columbia, MD, etc.).

### C. Solid State and soluble high throughput assays

In one embodiment the invention provides soluble assays using an angiogenesis protein, or a cell or tissue expressing an angiogenesis protein, either naturally occurring or recombinant. In another embodiment, the invention provides solid phase based *in vitro* assays in a high throughput format, where the angiogenesis protein is attached to a solid phase substrate. Any one of the assays described herein can be adapted for high throughput screening, e.g., ligand binding, cellular proliferation, cell surface marker flux, e.g.,  $\alpha\beta3$  integrin, etc. In one preferred embodiment, the cell-based system using  $\alpha\beta3$  integrin modulation and FACS assays is used in a high throughput format for identifying modulators of angiogenesis proteins, and therefore modulators of T cell angiogenesis.

In the high throughput assays of the invention, either soluble or solid state, it is possible to screen up to several thousand different modulators or ligands in a single day. This methodology can be used for angiogenesis proteins *in vitro*, or for cell-based assays comprising an angiogenesis protein. In particular, each well of a microtiter plate can be used to run a separate assay against a selected potential modulator, or, if concentration or incubation time effects are to be observed, every 5-10 wells can test a

single modulator. Thus, a single standard microtiter plate can assay about 100 (e.g., 96) modulators. If 1536 well plates are used, then a single plate can easily assay from about 100- about 1500 different compounds. It is possible to assay many plates per day; assay screens for up to about 6,000, 20,000, 50,000, or more than 100,000 different compounds are possible using the integrated systems of the invention.

For a solid state reaction, the protein of interest or a fragment thereof, e.g., an extracellular domain, or a cell comprising the protein of interest or a fragment thereof as part of a fusion protein can be bound to the solid state component, directly or indirectly, via covalent or non covalent linkage e.g., via a tag. The tag can be any of a variety of components. In general, a molecule which binds the tag (a tag binder) is fixed to a solid support, and the tagged molecule of interest is attached to the solid support by interaction of the tag and the tag binder.

A number of tags and tag binders can be used, based upon known molecular interactions well described in the literature. For example, where a tag has a natural binder, for example, biotin, protein A, or protein G, it can be used in conjunction with appropriate tag binders (avidin, streptavidin, neutravidin, the Fc region of an immunoglobulin, etc.) Antibodies to molecules with natural binders such as biotin are also widely available and appropriate tag binders; *see*, SIGMA Immunochemicals 1998 catalogue SIGMA, St. Louis MO).

Similarly, any haptenic or antigenic compound can be used in combination with an appropriate antibody to form a tag/tag binder pair. Thousands of specific antibodies are commercially available and many additional antibodies are described in the literature. For example, in one common configuration, the tag is a first antibody and the tag binder is a second antibody which recognizes the first antibody. In addition to antibody-antigen interactions, receptor-ligand interactions are also appropriate as tag and tag-binder pairs. For example, agonists and antagonists of cell membrane receptors (e.g., cell receptor-ligand interactions such as transferrin, c-kit, viral receptor ligands, cytokine receptors, chemokine receptors, interleukin receptors, immunoglobulin receptors and antibodies, the cadherein family, the integrin family, the selectin family, and the like; *see*, e.g., Pigott & Power, *The Adhesion Molecule Facts Book I* (1993). Similarly, toxins and venoms, viral epitopes, hormones (e.g., opiates, steroids, etc.), intracellular receptors (e.g. which mediate the effects of various small ligands, including steroids, thyroid hormone, retinoids and vitamin D; peptides), drugs, lectins, sugars, nucleic acids (both linear and

cyclic polymer configurations), oligosaccharides, proteins, phospholipids and antibodies can all interact with various cell receptors.

Synthetic polymers, such as polyurethanes, polyesters, polycarbonates, polyureas, polyamides, polyethyleneimines, polyarylene sulfides, polysiloxanes, polyimides, and polyacetates can also form an appropriate tag or tag binder. Many other tag/tag binder pairs are also useful in assay systems described herein, as would be apparent to one of skill upon review of this disclosure.

Common linkers such as peptides, polyethers, and the like can also serve as tags, and include polypeptide sequences, such as poly gly sequences of between about 5 and 200 amino acids. Such flexible linkers are known to persons of skill in the art. For example, poly(ethylene glycol) linkers are available from Shearwater Polymers, Inc. Huntsville, Alabama. These linkers optionally have amide linkages, sulfhydryl linkages, or heterofunctional linkages.

Tag binders are fixed to solid substrates using any of a variety of methods currently available. Solid substrates are commonly derivatized or functionalized by exposing all or a portion of the substrate to a chemical reagent which fixes a chemical group to the surface which is reactive with a portion of the tag binder. For example, groups which are suitable for attachment to a longer chain portion would include amines, hydroxyl, thiol, and carboxyl groups. Aminoalkylsilanes and hydroxyalkylsilanes can be used to functionalize a variety of surfaces, such as glass surfaces. The construction of such solid phase biopolymer arrays is well described in the literature. *See, e.g.,* Merrifield, *J. Am. Chem. Soc.* 85:2149-2154 (1963) (describing solid phase synthesis of, e.g., peptides); Geysen *et al.*, *J. Immun. Meth.* 102:259-274 (1987) (describing synthesis of solid phase components on pins); Frank & Doring, *Tetrahedron* 44:6031-6040 (1988) (describing synthesis of various peptide sequences on cellulose disks); Fodor *et al.*, *Science*, 251:767-777 (1991); Sheldon *et al.*, *Clinical Chemistry* 39(4):718-719 (1993); and Kozal *et al.*, *Nature Medicine* 2(7):753-759 (1996) (all describing arrays of biopolymers fixed to solid substrates). Non-chemical approaches for fixing tag binders to substrates include other common methods, such as heat, cross-linking by UV radiation, and the like.

### **Antibodies to angiogenesis and tumorigenesis polypeptides**

In addition to the detection of gene and gene expression using nucleic acid hybridization technology, one can also use immunoassays to detect proteins of the

invention. Such assays are useful for screening for modulators of angiogenesis, as well as for therapeutic and diagnostic applications. Immunoassays can be used to qualitatively or quantitatively analyze angiogenesis protein. A general overview of the applicable technology can be found in Harlow & Lane, *Antibodies: A Laboratory Manual* (1988).

5

#### *A. Production of antibodies*

Methods of producing polyclonal and monoclonal antibodies that react specifically with the angiogenesis proteins are known to those of skill in the art (*see, e.g.,* Coligan, *Current Protocols in Immunology* (1991); Harlow & Lane, *supra*; Goding, *Monoclonal Antibodies: Principles and Practice* (2d ed. 1986); and Kohler & Milstein, *Nature* 256:495-497 (1975)). Such techniques include antibody preparation by selection of antibodies from libraries of recombinant antibodies in phage or similar vectors, as well as preparation of polyclonal and monoclonal antibodies by immunizing rabbits or mice (*see, e.g.,* Huse *et al.*, *Science* 246:1275-1281 (1989); Ward *et al.*, *Nature* 341:544-546 (1989)).

15

A number of immunogens comprising portions of an angiogenesis protein may be used to produce antibodies specifically reactive with protein. For example, recombinant protein or an antigenic fragment thereof, can be isolated as described herein. Recombinant protein can be expressed in eukaryotic or prokaryotic cells as described above, and purified as generally described above. Recombinant protein is the preferred immunogen for the production of monoclonal or polyclonal antibodies. Alternatively, a synthetic peptide derived from the sequences disclosed herein and conjugated to a carrier protein can be used as an immunogen. Naturally occurring protein may also be used either in pure or impure form. The product is then injected into an animal capable of producing antibodies. Either monoclonal or polyclonal antibodies may be generated, for subsequent use in immunoassays to measure the protein.

20

25

Methods of production of polyclonal antibodies are known to those of skill in the art. An inbred strain of mice (*e.g.,* BALB/C mice) or rabbits is immunized with the protein using a standard adjuvant, such as Freund's adjuvant, and a standard immunization protocol. The animal's immune response to the immunogen preparation is monitored by taking test bleeds and determining the titer of reactivity to the beta subunits. When appropriately high titers of antibody to the immunogen are obtained, blood is collected from the animal and antisera are prepared. Further fractionation of the antisera

30

to enrich for antibodies reactive to the protein can be done if desired (*see*, Harlow & Lane, *supra*).

Monoclonal antibodies may be obtained by various techniques familiar to those skilled in the art. Briefly, spleen cells from an animal immunized with a desired antigen are immortalized, commonly by fusion with a myeloma cell (*see*, Kohler & Milstein, *Eur. J. Immunol.* 6:511-519 (1976)). Alternative methods of immortalization include transformation with Epstein Barr Virus, oncogenes, or retroviruses, or other methods well known in the art. Colonies arising from single immortalized cells are screened for production of antibodies of the desired specificity and affinity for the antigen, and yield of the monoclonal antibodies produced by such cells may be enhanced by various techniques, including injection into the peritoneal cavity of a vertebrate host. Alternatively, one may isolate DNA sequences which encode a monoclonal antibody or a binding fragment thereof by screening a DNA library from human B cells according to the general protocol outlined by Huse, *et al.*, *Science* 246:1275-1281 (1989).

Monoclonal antibodies and polyclonal sera are collected and titered against the immunogen protein in an immunoassay, for example, a solid phase immunoassay with the immunogen immobilized on a solid support. Typically, polyclonal antisera with a titer of  $10^4$  or greater are selected and tested for their cross reactivity against non- angiogenesis proteins, using a competitive binding immunoassay. Specific polyclonal antisera and monoclonal antibodies will usually bind with a  $K_d$  of at least about 0.1 mM, more usually at least about 1  $\mu$ M, preferably at least about 0.1  $\mu$ M or better, and most preferably, 0.01  $\mu$ M or better. Antibodies specific only for a particular ortholog, such as a human ortholog, can also be made, by subtracting out other cross-reacting orthologs from a species such as a non-human mammal. In this manner, antibodies that bind only to a desired protein may be obtained.

Once the specific antibodies against the protein are available, the protein can be detected by a variety of immunoassay methods. In addition, the antibody can be used therapeutically as modulators. For a review of immunological and immunoassay procedures, *see Basic and Clinical Immunology* (Stites & Terr eds., 7<sup>th</sup> ed. 1991).

Moreover, the immunoassays of the present invention can be performed in any of several configurations, which are reviewed extensively in *Enzyme Immunoassay* (Maggio, ed., 1980); and Harlow & Lane, *supra*.

*B. Immunological binding assays*

Protein can be detected and/or quantified using any of a number of well recognized immunological binding assays (*see, e.g.*, U.S. Patents 4,366,241; 4,376,110; 4,517,288; and 4,837,168). For a review of the general immunoassays, see also *Methods in Cell Biology: Antibodies in Cell Biology*, volume 37 (Asai, ed. 1993); *Basic and Clinical Immunology* (Stites & Terr, eds., 7th ed. 1991). Immunological binding assays (or immunoassays) typically use an antibody that specifically binds to a protein or antigen of choice (in this case the protein or antigenic subsequence thereof). The antibody may be produced by any of a number of means well known to those of skill in the art and as described above.

Immunoassays also often use a labeling agent to specifically bind to and label the complex formed by the antibody and antigen. The labeling agent may itself be one of the moieties comprising the antibody/antigen complex. Thus, the labeling agent may be a labeled protein or a labeled antibody. Alternatively, the labeling agent may be a third moiety, such a secondary antibody, that specifically binds to the antibody/protein complex (a secondary antibody is typically specific to antibodies of the species from which the first antibody is derived). Other proteins capable of specifically binding immunoglobulin constant regions, such as protein A or protein G may also be used as the label agent. These proteins exhibit a strong non-immunogenic reactivity with immunoglobulin constant regions from a variety of species (*see, e.g.*, Kronval *et al.*, *J. Immunol.* 111:1401-1406 (1973); Akerstrom *et al.*, *J. Immunol.* 135:2589-2542 (1985)). The labeling agent can be modified with a detectable moiety, such as biotin, to which another molecule can specifically bind, such as streptavidin. A variety of detectable moieties are well known to those skilled in the art.

Throughout the assays, incubation and/or washing steps may be required after each combination of reagents. Incubation steps can vary from about 5 seconds to several hours, optionally from about 5 minutes to about 24 hours. However, the incubation time will depend upon the assay format, antigen, volume of solution, concentrations, and the like. Usually, the assays will be carried out at ambient temperature, although they can be conducted over a range of temperatures, such as 10°C to 40°C.

### Non-competitive assay formats

Immunoassays for detecting protein in samples may be either competitive or noncompetitive. Noncompetitive immunoassays are assays in which the amount of antigen is directly measured. In one preferred “sandwich” assay, for example, the antibodies can be bound directly to a solid substrate on which they are immobilized. These immobilized antibodies then capture protein present in the test sample. Proteins thus immobilized are then bound by a labeling agent, such as a second antibody bearing a label. Alternatively, the second antibody may lack a label, but it may, in turn, be bound by a labeled third antibody specific to antibodies of the species from which the second antibody is derived. The second or third antibody is typically modified with a detectable moiety, such as biotin, to which another molecule specifically binds, e.g., streptavidin, to provide a detectable moiety.

### Competitive assay formats

In competitive assays, the amount of protein present in the sample is measured indirectly by measuring the amount of a known, added (exogenous) protein displaced (competed away) from an antibody by the unknown protein present in a sample. In one competitive assay, a known amount of protein is added to a sample and the sample is then contacted with an antibody that specifically binds to protein. The amount of exogenous protein bound to the antibody is inversely proportional to the concentration of protein present in the sample. In a particularly preferred embodiment, the antibody is immobilized on a solid substrate. The amount of protein bound to the antibody may be determined either by measuring the amount of protein present in protein/antibody complex, or alternatively by measuring the amount of remaining uncomplexed protein. The amount of protein may be detected by providing a labeled molecule.

A hapten inhibition assay is another preferred competitive assay. In this assay the known protein is immobilized on a solid substrate. A known amount of antibody is added to the sample, and the sample is then contacted with the immobilized protein. The amount of antibody bound to the known immobilized protein is inversely proportional to the amount of protein present in the sample. Again, the amount of immobilized antibody may be detected by detecting either the immobilized fraction of antibody or the fraction of the antibody that remains in solution. Detection may be direct where the antibody is labeled or indirect by the subsequent addition of a labeled moiety that specifically binds to the antibody as described above.

### Cross-reactivity determinations

Immunoassays in the competitive binding format can also be used for crossreactivity determinations. For example, a protein can be immobilized to a solid support. Proteins are added to the assay that compete for binding of the antisera to the immobilized antigen. The ability of the added proteins to compete for binding of the antisera to the immobilized protein is compared to the ability of the protein to compete with itself. The percent crossreactivity for the above proteins is calculated, using standard calculations. Those antisera with less than 10% crossreactivity with each of the added proteins listed above are selected and pooled. The cross-reacting antibodies are optionally removed from the pooled antisera by immunoabsorption with the added considered proteins, e.g., distantly related homologs.

The immunoabsorbed and pooled antisera are then used in a competitive binding immunoassay as described above to compare a second protein, thought to be perhaps an allele or polymorphic variant of a protein, to the immunogen protein. In order to make this comparison, the two proteins are each assayed at a wide range of concentrations and the amount of each protein required to inhibit 50% of the binding of the antisera to the immobilized protein is determined. If the amount of the second protein required to inhibit 50% of binding is less than 10 times the amount of the protein that is required to inhibit 50% of binding, then the second protein is said to specifically bind to the polyclonal antibodies generated to the immunogen.

### Other assay formats

Western blot (immunoblot) analysis is used to detect and quantify the presence of protein in the sample. The technique generally comprises separating sample proteins by gel electrophoresis on the basis of molecular weight, transferring the separated proteins to a suitable solid support, (such as a nitrocellulose filter, a nylon filter, or derivatized nylon filter), and incubating the sample with the antibodies that specifically bind the protein. The antibodies specifically bind to the protein on the solid support. These antibodies may be directly labeled or alternatively may be subsequently detected using labeled antibodies (e.g., labeled sheep anti-mouse antibodies) that specifically bind to the antibodies.

Other assay formats include liposome immunoassays (LIA), which use liposomes designed to bind specific molecules (e.g., antibodies) and release encapsulated

reagents or markers. The released chemicals are then detected according to standard techniques (*see Monroe et al., Amer. Clin. Prod. Rev.* 5:34-41 (1986)).

#### Reduction of non-specific binding

5           One of skill in the art will appreciate that it is often desirable to minimize non-specific binding in immunoassays. Particularly, where the assay involves an antigen or antibody immobilized on a solid substrate it is desirable to minimize the amount of non-specific binding to the substrate. Means of reducing such non-specific binding are well known to those of skill in the art. Typically, this technique involves coating the  
10       substrate with a proteinaceous composition. In particular, protein compositions such as bovine serum albumin (BSA), nonfat powdered milk, and gelatin are widely used with powdered milk being most preferred.

#### Labels

15           The particular label or detectable group used in the assay is not a critical aspect of the invention, as long as it does not significantly interfere with the specific binding of the antibody used in the assay. The detectable group can be any material having a detectable physical or chemical property. Such detectable labels have been well-developed in the field of immunoassays and, in general, most any label useful in such  
20       methods can be applied to the present invention. Thus, a label is any composition detectable by spectroscopic, photochemical, biochemical, immunochemical, electrical, optical or chemical means. Useful labels in the present invention include magnetic beads (e.g., DYNABEADS<sup>TM</sup>), fluorescent dyes (e.g., fluorescein isothiocyanate, Texas red, rhodamine, and the like), radiolabels (e.g., <sup>3</sup>H, <sup>125</sup>I, <sup>35</sup>S, <sup>14</sup>C, or <sup>32</sup>P), enzymes (e.g., horse  
25       radish peroxidase, alkaline phosphatase and others commonly used in an ELISA), and colorimetric labels such as colloidal gold or colored glass or plastic beads (e.g., polystyrene, polypropylene, latex, etc.).

          The label may be coupled directly or indirectly to the desired component of the assay according to methods well known in the art. As indicated above, a wide  
30       variety of labels may be used, with the choice of label depending on sensitivity required, ease of conjugation with the compound, stability requirements, available instrumentation, and disposal provisions.

          Non-radioactive labels are often attached by indirect means. Generally, a ligand molecule (e.g., biotin) is covalently bound to the molecule. The ligand then binds

to another molecules (e.g., streptavidin) molecule, which is either inherently detectable or covalently bound to a signal system, such as a detectable enzyme, a fluorescent compound, or a chemiluminescent compound. The ligands and their targets can be used in any suitable combination with antibodies that recognize the protein, or secondary antibodies.

The molecules can also be conjugated directly to signal generating compounds, e.g., by conjugation with an enzyme or fluorophore. Enzymes of interest as labels will primarily be hydrolases, particularly phosphatases, esterases and glycosidases, or oxidotases, particularly peroxidases. Fluorescent compounds include fluorescein and its derivatives, rhodamine and its derivatives, dansyl, umbelliferone, etc. Chemiluminescent compounds include luciferin, and 2,3-dihydrophthalazinediones, e.g., luminol. For a review of various labeling or signal producing systems that may be used, see U.S. Patent No. 4,391,904.

Means of detecting labels are well known to those of skill in the art. Thus, for example, where the label is a radioactive label, means for detection include a scintillation counter or photographic film as in autoradiography. Where the label is a fluorescent label, it may be detected by exciting the fluorochrome with the appropriate wavelength of light and detecting the resulting fluorescence. The fluorescence may be detected visually, by means of photographic film, by the use of electronic detectors such as charge coupled devices (CCDs) or photomultipliers and the like. Similarly, enzymatic labels may be detected by providing the appropriate substrates for the enzyme and detecting the resulting reaction product. Finally simple colorimetric labels may be detected simply by observing the color associated with the label. Thus, in various dipstick assays, conjugated gold often appears pink, while various conjugated beads appear the color of the bead.

Some assay formats do not require the use of labeled components. For instance, agglutination assays can be used to detect the presence of the target antibodies. In this case, antigen-coated particles are agglutinated by samples comprising the target antibodies. In this format, none of the components need be labeled and the presence of the target antibody is detected by simple visual inspection.

### **Gene therapy**

The present invention provides the nucleic acids of angiogenesis or tumorigenesis associated protein for the transfection of cells *in vitro* and *in vivo*. These

nucleic acids can be inserted into any of a number of well-known vectors for the transfection of target cells and organisms as described below. The nucleic acids are transfected into cells, *ex vivo* or *in vivo*, through the interaction of the vector and the target cell. The nucleic acid, under the control of a promoter, then expresses a protein of the present invention, thereby mitigating the effects of absent, partial inactivation, or abnormal expression of the angiogenesis or tumorigenesis gene, particularly as it relates to angiogenesis. The compositions are administered to a patient in an amount sufficient to elicit a therapeutic response in the patient. An amount adequate to accomplish this is defined as "therapeutically effective dose or amount."

Such gene therapy procedures have been used to correct acquired and inherited genetic defects, cancer, and other diseases in a number of contexts. The ability to express artificial genes in humans facilitates the prevention and/or cure of many important human diseases, including many diseases which are not amenable to treatment by other therapies (for a review of gene therapy procedures, *see* Anderson, *Science* 256:808-813 (1992); Nabel & Felgner, *TIBTECH* 11:211-217 (1993); Mitani & Caskey, *TIBTECH* 11:162-166 (1993); Mulligan, *Science* 926-932 (1993); Dillon, *TIBTECH* 11:167-175 (1993); Miller, *Nature* 357:455-460 (1992); Van Brunt, *Biotechnology* 6(10):1149-1154 (1998); Vigne, *Restorative Neurology and Neuroscience* 8:35-36 (1995); Kremer & Perricaudet, *British Medical Bulletin* 51(1):31-44 (1995); Haddada *et al.*, in *Current Topics in Microbiology and Immunology* (Doerfler & Böhm eds., 1995); and Yu *et al.*, *Gene Therapy* 1:13-26 (1994)).

The nucleic acids of the invention can also be used to make transgenic animals, such as transgenic mice, either by knock-out or overexpression. Such animals are useful, e.g., for testing modulators of angiogenesis and tumorigenesis.

## **Pharmaceutical compositions and administration**

Pharmaceutically acceptable carriers are determined in part by the particular composition being administered (e.g., nucleic acid, protein, modulatory compounds or transduced cell), as well as by the particular method used to administer the composition. Accordingly, there are a wide variety of suitable formulations of pharmaceutical compositions of the present invention (*see, e.g., Remington's Pharmaceutical Sciences*, 17<sup>th</sup> ed., 1989). Administration can be in any convenient manner, e.g., by injection, oral administration, inhalation, transdermal application, or rectal administration.

Formulations suitable for oral administration can consist of (a) liquid solutions, such as an effective amount of the packaged nucleic acid suspended in diluents, such as water, saline or PEG 400; (b) capsules, sachets or tablets, each containing a predetermined amount of the active ingredient, as liquids, solids, granules or gelatin; (c) 5 suspensions in an appropriate liquid; and (d) suitable emulsions. Tablet forms can include one or more of lactose, sucrose, mannitol, sorbitol, calcium phosphates, corn starch, potato starch, microcrystalline cellulose, gelatin, colloidal silicon dioxide, talc, magnesium stearate, stearic acid, and other excipients, colorants, fillers, binders, diluents, buffering agents, moistening agents, preservatives, flavoring agents, dyes, disintegrating 10 agents, and pharmaceutically compatible carriers. Lozenge forms can comprise the active ingredient in a flavor, e.g., sucrose, as well as pastilles comprising the active ingredient in an inert base, such as gelatin and glycerin or sucrose and acacia emulsions, gels, and the like containing, in addition to the active ingredient, carriers known in the art.

The compound of choice, alone or in combination with other suitable 15 components, can be made into aerosol formulations (i.e., they can be “nebulized”) to be administered via inhalation. Aerosol formulations can be placed into pressurized acceptable propellants, such as dichlorodifluoromethane, propane, nitrogen, and the like.

Formulations suitable for parenteral administration, such as, for example, by intraarticular (in the joints), intravenous, intramuscular, intradermal, intraperitoneal, 20 and subcutaneous routes, include aqueous and non-aqueous, isotonic sterile injection solutions, which can contain antioxidants, buffers, bacteriostats, and solutes that render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions that can include suspending agents, solubilizers, thickening agents, stabilizers, and preservatives. In the practice of this invention, compositions can 25 be administered, for example, by intravenous infusion, orally, topically, intraperitoneally, intravesically or intrathecally. Parenteral administration and intravenous administration are the preferred methods of administration. The formulations of compounds can be presented in unit-dose or multi-dose sealed containers, such as ampules and vials.

Injection solutions and suspensions can be prepared from sterile powders, 30 granules, and tablets of the kind previously described. Cells transduced by nucleic acids for *ex vivo* therapy can also be administered intravenously or parenterally as described above.

The dose administered to a patient, in the context of the present invention should be sufficient to effect a beneficial therapeutic response in the patient over time.

The dose will be determined by the efficacy of the particular vector employed and the condition of the patient, as well as the body weight or surface area of the patient to be treated. The size of the dose also will be determined by the existence, nature, and extent of any adverse side-effects that accompany the administration of a particular vector, or  
 5 transduced cell type in a particular patient.

In determining the effective amount of the vector to be administered in the treatment or prophylaxis of conditions owing to diminished or aberrant expression of the angiogenesis protein, the physician evaluates circulating plasma levels of the vector, vector toxicities, progression of the disease, and the production of anti-vector antibodies.

10 In general, the dose equivalent of a naked nucleic acid from a vector is from about 1  $\mu\text{g}$  to 100  $\mu\text{g}$  for a typical 70 kilogram patient, and doses of vectors which include a retroviral particle are calculated to yield an equivalent amount of therapeutic nucleic acid.

For administration, compounds and transduced cells of the present invention can be administered at a rate determined by the LD-50 of the inhibitor, vector,  
 15 or transduced cell type, and the side-effects of the inhibitor, vector or cell type at various concentrations, as applied to the mass and overall health of the patient. Administration can be accomplished via single or divided doses.

## EXAMPLES

20 The following examples are offered to illustrate, but not to limit the claimed invention.

### Example 1: Identification of genes involved in modulation of angiogenesis and tumorigenesis *in vitro*

25 Using a retroviral base system, , complex libraries of cDNAs and GFP-fusions have been expressed in human primary endothelial cells (e.g., HUVECS). Cells that switch to a differentiated angiostatic state can be identified. The cells are then assayed for migration along a haptotactic gradient of specific matrix proteins such as vitronectin and fibronectin in a Boyden chamber assay. Cells were selected for impaired  
 30 haptotaxis (*see, e.g., Klemke et al., J. Cell Biol.* 4:961-972 (1998)).

Example 2: Identification of genes involved in modulation of angiogenesis and tumorigenesis *in vivo*

This study in mice was designed to provide information on the growth and angiogenic characteristics of MDA-MB-231 infected with either EFS-U6TO-Ax14 vector or EFS-U6TO-FF1 (isogenic negative control) and implanted subcutaneously in mice (Strain:CB-17 scid). MDA-MB-231 (wt) cells will also be implanted and measured. Tumor Cell Injection Density:  $1 \times 10^7$  per injection site (shaved right hind flank).

When tumors reach a diameter of approximately 1 cm, animals are randomly divided into subgroups according to tumor size using a stratified randomization method (Day 0). Tumors in selected subgroups are resected by survival surgery and fixed in formalin and the animals will be observed for signs of metastasis via changes in body weight and/or clinical signs. Animals showing any of these signs in severity are terminated and the lungs, liver, sternum, femur, and lymph nodes (axial and mesenteric) are harvested and fixed in 10% buffered formalin for histological examination. Tumors in remaining subgroups will continue to be measured until they reach 2,000mm<sup>3</sup>.

The mean tumor volume for each group was calculated for each time point. Comparisons between groups at specific times are made using an unpaired, two-tailed t-test, and the results analyzed using analysis of variance (ANOVA) or the student's t-test. Differentials in tumor growth between groups are determined by calculating volume-doubling time (VDT), or by determining the survival time.

It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference in their entirety for all purposes.

## SEQUENCE LISTING

Ax1

5 GH2\_420\_G3F1  
CTCCAGGGGTTTCAGGATAACCTCCACCCTCATCCATGTTGACATAGAGGATTTTCGTCAGGCTCCTGGGCAGG  
AGGCAAGG

GH2\_420\_G3R1

10 ATCTATCTAACCCTGTGCTTGGGTTCTGCGGCCTTGCCCTCCTGCCAGGAGCCTGACGAAATCCTCTATGT  
CAACATGGATGAGGGTGGAGGTTATCCTGAACCCCTGGAG

>gi|21536465|ref|NM\_021913.2| Homo sapiens AXL receptor tyrosine kinase  
(AXL), transcript variant 1, mRNA

15 GAGTGGAGTTCTGGAGGAATGTTTACCAGACACAGAGCCCAGAGGGACAGCGCCCAGAGCCCAGATAGAG  
AGACACGGCCTCACTGGCTCAGCACCAGGGTCCCTTCCCTCCCTCAGCTCCCTCCCTGGCCCCCTTTAA  
GAAAGAGCTGATCCTCTCCTCTCTTGAGTTAACCCTGATTGTCCAGGTGGCCCCCTGGCTCTGGCCTGGT  
GGGCGGAGGCAAAGGGGGAGCCAGGGGCGGAGAAAGGGTTGCCCAAGTCTGGGAGTGAGGGAAGGAGGCA  
GGGGTGCTGAGAAGGCGGCTGCTGGGCAGAGCCGGTGGCAAGGGCCTCCCTGCGCTGTGCCAGGCAGG  
20 CAGTGCCAAATCCGGGGAGCCTGGAGCTGGGGGAGGGCCGGGACAGCCCGGCCCTGCCCTGTGCCAGGCAGG  
CTGGGAGCCCAGCAACTTCTGAGGAAAGTTTGGCACCCATGGCGTGGCGGTGCCCCAGGATGGGCAGGGT  
CCCGCTGGCCTGGTGTGGCGCTGTGCGGCTGGGCGTGCATGGCCCCCAGGGGCACGCAGGCTGAAGAA  
AGTCCCTTCGTGGGCAACCCAGGGAATATCACAGGTGCCCGGGGACTCACGGGCACCCTTCGGTGTTCAGC  
TCCAGGTTTCAGGGAGAGCCCCCGAGGTACATTGGCTTCGGGATGGACAGATCCTGGAGCTCGCGGACAG  
25 CACCCAGACCCAGGTGCCCTGGGTGAGGATGAACAGGATGACTGGATAGTGGTCAGCCAGCTCAGAATC  
ACCTCCCTGCAGCTTTCGACACGGGACAGTACCAGTGTTCGGTGTTCCTGGGACATCAGACCTTCGTGT  
CCCAGCCTGGCTATGTTGGGCTGGAGGGCTTGCTTACTTCCTGGAGGAGCCCGAAGACAGGACTGTGGC  
CGCCAACACCCCTTCAACCTGAGCTGCCAAGCTCAGGGACCCCAAGAGCCCGTGGACCTACTCTGGCTC  
CAGGATGCTGTCCCCCTGGCCACGGCTCCAGGTACGGCCCCCAGCGCAGCCTGCATGTTCCAGGGCTGA  
30 ACAAGACACTCCTCTTCTCCTGCGAAGCCCATACGCCAAGGGGGTCAACACATCCCGCACAGCCACCAT  
CACAGTGCTCCCCCAGCAGCCCCGTAACTCCACCTGGTCTCCCGCCAACCCACGGAGCTGGAGGTGGCT  
TGGACTCCAGGCCTGAGCGGCATCTACCCCTGACCCACTGCACCCTGCAGGCTGTGCTGTTCAGACGATG  
GGATGGGCATCCAGGCGGGAGAACAGACCCCCCAGAGGAGCCCTCACCTCGCAAGCATCCGTGCCCCCC  
CCATCAGCTTCGGCTAGGCAGCCTCCATCCTCACACCCCTTATCACATCCGCGTGGCATGCACCAGCAGC  
35 CAGGGCCCCCTCATCCTGGACCCACTGGCTTCCTGTGGAGACGCCGGAGGGAGTGCCCCCTGGGCCCCCTG  
AGAACATTAGTGCTACGCGGAATGGGAGCCAGGCCTTCGTGCATTGGCAAGAGCCCCGGGCGCCCCCTGCA  
GGGTACCCTGTTAGGGTACCGGCTGGCGTATCAAGGCCAGGACACCCAGAGGTGCTAATGGACATAGGG  
CTAAGGCAAGAGGTGACCCCTGGAGCTGCAGGGGACGGGTCTGTGTCCAATCTGACAGTGTGTGTGGCAG  
CCTACACTGCTGTGGGGATGGACCTGGAGCCTCCAGTACCCCTGGAGGCCTGGCGCCCAGGGCAAGC  
40 ACAGCCAGTCCACCAGCTGGTGAAGGAACCTTCAACTCCTGCCTTCTCGTGGCCCTGGTGGTATGTACTG  
CTAGGAGCAGTCGTGGCGCTGCCTGTGTCTCTTGGCTCTCTTCTTGTCCACCGGCGAAAGAAGG  
AGACCCGTTATGGAGAAGTGTGTTGAACCAACAGTGGAAAGAGGTGAACTGGTAGTCAGGTACCGCGTGC  
CAAGTCTACAGTCGTGCGACCACTGAAGCTACCTTGAACAGCCTGGGCATCAGTGAAGAGCTGAAGGAG  
AAGCTGCGGGATGTGATGGTGGACCGGCACAAGGTGGCCCTGGGGAAGACTCTGGGAGAGGGAGAGTTTG  
45 GAGCTGTGATGGAAGGCCAGCTCAACCAGGACGACTCCATCCTCAAGGTGGCTGTGAAGACGATGAAGAT  
TGCCATCTGCACGAGGTGAGAGCTGGAGGATTTCTGAGTGAAGCGGTCTGCATGAAGGAATTTGACCAT  
CCCAACGTCATGAGGCTCATCGGTGTCTGTTTCCAGGGTTCTGAACGAGAGAGCTTCCAGCACCTGTGG  
TCATCTTACCTTTTCATGAAACATGGAGACCTACACAGCTTCCCTCCTCTATTCCCGGCTCGGGGACCAGCC  
AGTGTAACCTGCCACTCAGATGCTAGTGAAGTTTCATGGCAGACATCGCCAGTGGCATGGAGTATCTGAGT  
50 ACCAAGAGATTTCATACACCGGGACCTGGCGGCCAGGAAGTGCATGCTGAATGAGAACATGTCCGTGTGTG  
TGGCGGACTTCGGGCTCTCCAAGAAGATCTACAATGGGGACTACTACCGCCAGGGACGTATCGCCAAGAT  
GCCAGTCAAGTGGATTGCCATTGAGAGTCTAGCTGACCGTGTCTACACCAGCAAGAGCGATGTGTGGTCC  
TTCGGGGTGACAATGTGGGAGATTGCCACAAGAGGCCAAACCCCATATCCGGGCGTGGAGAACAGCGAGA  
TTTATGACTATCTGCGCCAGGGAAATCGCCTGAAGCAGCCTGCGGACTGTCTGGATGGACTGTATGCCTT  
55 GATGTGCGGGTGTGGGAGCTAAATCCCCAGGACCGGCCAAGTTTTACAGAGCTGCGGGAAGATTTGGAG  
AACACACTGAAGGCCTTGCTCCTGCCAGGAGCCTGACGAAATCCTCTATGTCAACATGGATGAGGGTG  
GAGGTTATCTGAACCCCTGGAGCTGCAGGAGGAGCTGACCCCAACCCAGCCAGACCTAAGGATTTC  
CTGTAGCTGCCTCACTGCGGCTGAGGTCCATCCTGCTGGACGCTATGTCCTCTGCCCTTCCACAACCCCT  
AGCCCCGCTCAGCCTGCTGATAGGGGCTCCCCAGCAGCCCCAGGGCAGGAGGATGGTGCCTGAGACAACC  
60 CTCCACCTGGTACTCCCTCTCAGGATCCAAGCTAAGCACTGCCACTGGGGAAAACCTCCACCTTCCCACTT  
TCCCACCCACGCCTTATCCCCACTTGCAGCCCTGTCTTCTACCTATCCCACCTCCATCCAGACAGGT

CCCTCCCCCTTCTCTGTGCAGTAGCATCACCTTGAAAGCAGTAGCATCACCATCTGTAAAAGGAAGGGGTT  
 GGATTGCAATATCTGAAGCCCTCCCAGGTGTTAACATTCCAAGACTCTAGAGTCCAAGGTTTAAAGAGTC  
 TAGATTCAAAGGTTCTAGGTTTCAAAGATGCTGTGAGTCTTTGGTTCTAAGGACCTGAAATTCCAAAGTC  
 TCTAATTCTATTAAAGTGCTAAGGTTCTAAGGCCCTACTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTT  
 5 TGCGATAGAGTCTCACTGTGTCAACCAGGCTGGAGTGCAATCTCGCCTCACTGCAACCTTCA  
 CCTACCGAGTTCAAGTGATTTTCTGCCTTGGCCTCCCAAGTAGCTGGGATTACAGGTGTGTGCCACCAC  
 ACCCGGCTAATTTTTATATTTTATAGTAGAGACAGGTTTCACCATGTTGGCCAGGCTGGTCTAAAACCTC  
 TGACCTCAAGTGATCTGCCCACCTCAGCCTCCCAAAGTGCTGAGATTACAGGCATGAGCCACTGCACTCA  
 ACCTTAAGACCTACTGTTCTAAAGCTCTGACATTATGTGGTTTTAGATTTTCTGGTTCTAACATTTTGA  
 10 TAAAGCCTCAAGGTTTTAGGTTCTAAAGTTCTAAGATTCTGATTTTAGGAGCTAAGGCTCTATGAGTCTA  
 GATGTTTATTCTTCTAGAGTTCAGAGTCTTAAATGTAAGATTATAGATTCTAAAGATTCTATAGTTCT  
 AGACATGGAGGTTCTAAGGCCTAGGATTCTAAATGTGATGTTCTAAGGCTCTGAGAGCTAGATTCTCT  
 GGCTGTAAGGCTCTAGATCATAAGGCTTCAAAATGTTATCTTCTCAAGTTCTAAGATTCTAATGATGATC  
 AATTATAGTTTCTGAGGCTTTATGATAATAGATTCTCTTGATAAGATCCTAGATCCTAAGGGTCGAAAG  
 15 CTCTAGAATCTGCAATTCAAAGTTCCAAGAGTCTAAAGATGGAGTTTCTAAGGTCCGGTGTCTAAGAT  
 GTGATATTCTAAGACTTACTCTAAGATCTTAGATTCTCTGTGTCTAAGATTCTAGATCAGATGCTCCAAG  
 ATTCTAGATGATTAAATAAGATTCTAACGGTCTGTTCTGTTTCAAGGCACTCTAGATTCCATTGGTCCAA  
 GATTCGGATCCTAAGCATCTAAGTTATAAGACTCTCACACTCAGTTGTGACTAACTAGACACCAAAGTT  
 CTAATAATTTCTAATGTTGGACACCTTTAGGTTCTTTGCTGCATTCTGCCTCTCTAGGACCATGGTTAAG  
 20 AGTCCAAGATCCACATTTCTAAAATCTTAGTTCTAGGCACTGTAGTTCTAAGACTCAAATGTTCTAA  
 GTTCTAAGATTCTAAGGTCCACAGGTCTAGACTATTAGGTGCAATTTCAAGGTTCTAACCTTACTATG  
 TAGTATTCTTTGGGGTGCCCTCTCCTTCTTAGCTATCATTGCTTCTCCTCCCAACTGTGGGGGTGTG  
 CCCCCTTCAAGCCTGTGCAATGCATTAGGGATGCCTCCTTTCCCGCAGGGGATGGACGATCTCCACCTT  
 TCGGGCCATGTTGCCCCGTGAGCCAATCCCTCACCTTCTGAGTACAGAGTGTGGACTCTGGTGCCTCCA  
 25 GAGGGGCTCAGGTCACATAAACTTTGTATATCAACGAAAAAA

>gi|21536466|ref|NP\_068713.2| AXL receptor tyrosine kinase isoform 1;  
 AXL transforming sequence/gene; oncogene AXL [Homo sapiens]  
 MAWRCPMRGRVPLAWCLALCGWACMAPRGTAQESPFVGNPGNITGARGLTGTLRCQLQVQGEPPPEVHWL  
 30 RDGQILELADSTQTQVPLGEDEQDDWIVVSQLRITSLQLSDTGQYQCLVFLGHQTFVSPQGYVGLGLPY  
 FLEPEDRTVAANTPFNLSCQAQGPPEPVDLLWLQDAVPLATAPGHGPQRSLLHVPGLNKTSSFSCEAHNA  
 KGVTTSTRATITVLPQQPRNLHLVSRQPTELEVAWTPGLSGIYPLTHCTLQAVLSDDGMGIQAGEPDPE  
 EPLTSQASVPPHQLRLGSLHPHTPYHIRVACTSSQGPSSWTHWLPVETPEGVPLGPPENISATRNGSQAF  
 VHWQEPRAPLQGTLLGYRLAYQGQDTPEVLMIDIGLRQEVTLQLQDGSVSNLTVCAAYTAAGDGPWSLP  
 35 VPLEAWRPGQAQPVHQLVKEPSTPAFSWPWWYVLLGAVVAAACVLILALFLVHRRKKETRYGEVFEPVE  
 RGELVVRYRVRKYSRRTTEATLNSLGISEELKEKL RDVMVDRHKVALGKTLGEGEFGAVMEGQLNQDDS  
 ILKVAVKTMKIAICTRSELEDFLSEAVCMKEFDHPNVMRLIGVCFQGSERESFPAPVVILPFMKHGDLS  
 FLLYSRLGDQPVYLPQMLVKFMADIASGMEYLSKRFIHRDLAARNCMLNENMSVCVADFGLSKKIYNG  
 DYYRQGRIAKMPVKWIAIESLADRVYTSKSDVWSFGVTMWEIATRGQTPYPGVENSEIYDYLRLQGNRLKQ  
 40 PADCLDGLYALMSRCWELNPQDRPSFTELREDLNTLKPAPQEPDEILYVNMDEGGGYPEPPGAAGGA  
 DPPTQPDPKDSCSLTAAEVHPAGRYVLCPTSTPSPAQPADRGSPAAPGQEDGA

>gi|21536467|ref|NM\_001699.3| Homo sapiens AXL receptor tyrosine kinase  
 (AXL), transcript variant 2, mRNA  
 45 GAGTGGAGTTCTGGAGGAATGTTTACCAGACACAGAGCCCAGAGGGACAGCGCCCAGAGCCCAGATAGAG  
 AGACACGGCCTCACTGGCTCAGCACCAGGGTCCCCCTCCCCCTCCTCAGCTCCCTCCCTGGCCCCCTTAA  
 GAAAGAGCTGATCCTCTCCTCTCTTGAGTTAACCCCTGATTGTCCAGGTGGCCCCCTGGCTCTGGCCTGGT  
 GGGCGGAGGCAAAGGGGGAGCCAGGGCGGAGAAAGGTTGCCCAAGTCTGGGAGTGAGGGAAGGAGGCA  
 GGGGTGCTGAGAAGGCGGCTGCTGGGCAGAGCCGGTGGCAAGGGCCTCCCCCTGCCGCTGTGCCAGGCAGG  
 50 CAGTGCCAAATCCGGGGAGCCTGGAGCTGGGGGGAGGGCCGGGGACAGCCCGGCCCTGCCCCCTCCCCCG  
 CTGGGAGCCCAGCAACTTCTGAGGAAAGTTTGGCACCCATGGCGTGGCGGTGCCCCAGGATGGGCAGGGT  
 CCCGCTGGCCTGGTGTCTGGCGCTGTGCGGCTGGGCGTGCATGGCCCCCAGGGGCACGCAGGCTGAAGAA  
 AGTCCCTTCGTGGGCAACCCAGGGAATATCACAGGTGCCCGGGGACTCACGGGCACCCCTTCGGTGTGAGC  
 TCCAGGTTCCAGGGAGAGCCCCCGAGGTACATTGGCTTCGGGATGGACAGATCCTGGAGCTCGCGGACAG  
 55 CACCCAGACCCAGGTGCCCCCTGGGTGAGGATGAACAGGATGACTGGATAGTGGTCAGCCAGCTCAGAATC  
 ACCTCCCTGCAGCTTTCCGACACGGGACAGTACCAGTGTGTTGGTGTCTTGGGACATCAGACCTTCGTGT  
 CCCAGCTGGCTATGTTGGGCTGGAGGGCTGACCTTACTTCTGGAGGAGCCCGAAGCAGGACTGTGGC  
 CGCCAAACCCCCCTTCAACCTGAGCTGCCAAGCTCAGGGACCCCCAGAGCCCGTGGACCTACTCTGGCTC  
 CAGGATGCTGTCCCCCTGGCCACGGCTCCAGGTACAGGCCCCCAGCGCAGCCTGCATGTTCCAGGGCTGA  
 60 ACAAGACATCCTCTTCTCCTGCGAAGCCCATAACGCCAAGGGGGTACCACATCCCGCACAGCCACCAT  
 CACAGTGCTCCCCCAGCAGCCCCGTAACTCCACCTGGTCTCCCGCCAACCCACGGAGCTGGAGGTGGCT

TGGACTCCAGGCCTGAGCGGCATCTACCCCCTGACCCACTGCACCCTGCAGGCTGTGCTGTGACAGCATG  
 GGATGGGCATCCAGGCGGGAGAACCAGACCCCCCAGAGGAGCCCCTCACCTCGCAAGCATCCGTGCCCCC  
 CCATCAGCTTCGGCTAGGCAGCCTCCATCCTCACACCCCTTATCACATCCGCGTGGCATGCACCAGCAGC  
 CAGGGCCCCCTCATCCTGGACCCACTGGCTTCTGTGGAGACGCCGAGGGAGTGCCCCCTGGGCCCCCTG  
 5 AGAACATTAGTGCTACGCGGAATGGGAGCCAGGCCTTCGTGCATTGGCAAGAGCCCCGGGCGCCCCCTGCA  
 GGGTACCCTGTTAGGGTACCGGCTGGCGTATCAAGGCCAGGACACCCAGAGGTGCTAATGGACATAGGG  
 CTAAGGCAAGAGGTGACCCTGGAGCTGCAGGGGGACGGGTCTGTGTCCAATCTGACAGTGTGTGTGGCAG  
 CCTACACTGCTGCTGGGGATGGACCCTGGAGCCTCCCAGTACCCCTGGAGGCCTGGCGCCCAGTGAAGGA  
 ACCTTCAACTCCTGCCTTCTCGTGGCCCTGGTGGTATGTACTGCTAGGAGCAGTCTGTGGCCGCTGCCTGT  
 10 GTCCTCATCTTGGCTCTCTTCTTGTCCACCGGCGAAAGAAGGAGACCCGTTATGGAGAAGTGTGTTGAAC  
 CAACAGTGGAAAGAGGTGAAGTGGTAGTCAGGTACCGCGTGCAGTCTACAGTCTGCGGACCACTGA  
 AGCTACCTTGAACAGCCTGGGCATCAGTGAAGAGCTGAAGGAGAAGCTGCGGGATGTGATGGTGGACCGG  
 CACAAGGTGGCCCTGGGGAAGACTCTGGGAGAGGGAGAGTTTGGAGCTGTGATGGAAGGCCAGCTCAACC  
 AGGACGACTCCATCCTCAAGGTGGCTGTGAAGACGATGAAGATTGCCATCTGCACGAGGTGAGAGCTGGA  
 15 GGATTTCTGAGTGAAGCGGTCTGCATGAAGGAATTTGACCATCCCAACGTGATGAGGCTCATCGGTGTC  
 TGTTTCCAGGGTTCTGAACGAGAGAGCTTCCAGCACCTGTGGTTCATCTTACCTTTTCATGAAACATGGAG  
 ACCTACACAGCTTCTCTCTATTCCCGGCTCGGGGACCAGCCAGTGTACCTGCCCCACTCAGATGCTAGT  
 GAAGTTCATGGCAGACATCGCCAGTGGCATGGAGTATCTGAGTACCAAGAGATTTCATACACCGGGACCTG  
 GCGGCCAGGAAGTGCATGCTGAATGAGAATGTCCGTGTGTGTGGCGGACTTCGGGCTCTCCAAGAAGA  
 20 TCTACAATGGGGACTACTACCGCCAGGGACGTATCGCAAGATGCCAGTCAAGTGGATTGCCATTGAGAG  
 TCTAGCTGACCGTGTCTACACCAGCAAGAGCGATGTGTGGTCTTCGGGGTGACAATGTGGGAGATTGCC  
 ACAAGAGGCCAAACCCCATATCCGGGCGTGGAGAACAGCGAGATTTATGACTATCTGCGCCAGGGAAATC  
 GCCTGAAGCAGCCTGCGGACTGTCTGGATGGACTGTATGCCCTTGATGTGCGGGTGTGGGAGCTAAATCC  
 CCAGGACCGGCCAAGTTTTACAGAGCTGCGGGAAGATTTGGAGAACACACTGAAGGCCTTGCCCTCTGCC  
 25 CAGGAGCCTGACGAAATCCTCTATGTCAACATGGATGAGGGTGGAGGTTATCCTGAACCCCCCTGGAGCTG  
 CAGGAGGAGCTGACCCCCCAACCCAGCCAGACCTAAGGATTCCCTGTAGCTGCCTCACTGCGGCTGAGGT  
 CCATCCTGCTGGACGCTATGTCTCTGCCCTTCCACAACCCCTAGCCCCGCTCAGCCTGCTGATAGGGGC  
 TCCCCAGCAGCCCCAGGGCAGGAGGATGGTGCCTGAGACAACCCCTCACCTGGTACTCCCTCTCAGGATC  
 CAAGCTAAGCACTGCCACTGGGGAAAACTCCACCTTCCCACTTTCCCAACCCAGCCTTATCCCCACTTG  
 30 CAGCCCTGTCTTCTCCTATCCCACTCCAGTCCCAAGTCCCTCCCCTTCTCTGTGCTAGTAGCATC  
 ACCTTGAAGCAGTAGCATCACCATCTGTAAAGGAAGGGGTTGGATTGCAATATCTGAAGCCCTCCAG  
 GTGTTAACAATTCCAAGACTCTAGAGTCCAAGGTTTAAAGAGTCTAGATTCAAAGGTTCTAGGTTTCAAAG  
 ATGCTGTGAGTCTTTGGTTCTAAGGACCTGAAATTCCAAAGTCTCTAATTCTATTAAAGTGCTAAGGTTT  
 TAAGGCCTACTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTGGGATAGAGTCTCACTGTGTCAACCA  
 35 GGCTGGAGTGCAGTGGTGCAATCTCGCCTCACTGCAACCTTCACCTACCGAGTTCAAGTGATTTTCTGTC  
 CTTGGCCTCCCAAGTAGCTGGGATTACAGGTGTGTGCCACCACACCCGGCTAATTTTTATATTTTATAGTA  
 GAGACAGGGTTTACCATGTTGGCCAGGCTGGTCTAAAACCTCTGACCTCAAGTGATCTGCCCACCTCAG  
 CCTCCCAAAGTGCTGAGATTACAGGCATGAGCCACTGCACTCAACCTTAAGACCTACTGTTCTAAAGCTC  
 TGACATTATGTGTTTTAGATTTTCTGGTTCTAACATTTTGTGATAAAGCCTCAAGGTTTAGGTTCTAAA  
 40 GTTCTAAGATTCTGATTTTAGGAGCTAAGGCTCTATGAGTCTAGATGTTTATTCTTCTAGAGTTTCAAGT  
 CCTTAAATGTAAGATTATAGATTCTAAAGATTCTATAGTTCTAGACATGGAGGTTCTAAGGCCTAGGAT  
 TCTAAATGTGATGTTCTAAGGCTCTGAGAGTCTAGATTCTCTGGCTGTAAGGCTCTAGATCATAAGGCT  
 TCAAATGTTATCTTCTCAAGTTCTAAGATTCTAATGATGATCAATTATAGTTTCTGAGGCTTTATGATA  
 ATAGATTCTCTTGATATAAGATCCTAGATCCTAAGGGTCGAAAGCTCTAGAATCTGCAATTCAAAGTTCC  
 45 AAGAGTCTAAGATGGAGTTTCTAAGGTCCGGTGTCTAAGATGTGATATTCTAAGACTTACTCTAAGAT  
 CTTAGATTCTCTGTGTCTAAGATTCTAGATCAGATGCTCCAAGATTCTAGATGATTAAATAAGATTCTAA  
 CGGTCTGTTCTGTTTCAAGGCACTCTAGATTCCATTGGTCCAAGATTCCGGATCCTAAGCATCTAAGTTA  
 TAAGACTCTCACACTCAGTTGTGACTAAGTACACCAAGTTCTAATAATTTCTAATGTTGGACACCTT  
 TAGGTTCTTTGCTGCATTCTGCCTCTCTAGGACCATGGTTAAGAGTCCAAGAATCCACATTTCTAAAAATC  
 50 TTATAGTTCTAGGCACTGTAGTTCTAAGACTCAAATGTTCTAAGTTTCTAAGATTCTAAGGTCCACAGG  
 TCTAGACTATTAGGTGCAATTTCAAGGTCTAACCCTATACTGTAGTATTCTTTGGGGTGCCCCCTCTCCT  
 TCTTAGCTATCATTGCTTCTCTCTCCCACTGTGGGGGTGTGCCCCCTTCAAGCCTGTGCAATGCATTA  
 GGGATGCCTCCTTTCCCGCAGGGGATGGACGATCTCCACCTTTCGGGCCATGTTGCCCCCGTGAGCCAA  
 TCCCTCACCTTCTGAGTACAGAGTGTGGACTCTGGTGCCTCCAGAGGGGCTCAGGTACATAAAAACTTTG  
 55 TATATCAACGAAAAAA

>gi|21536468|ref|NP\_001690.2| AXL receptor tyrosine kinase isoform 2;  
 AXL transforming sequence/gene; oncogene AXL [Homo sapiens]  
 MAWRCPRMGRVPLAWCLALCGWACMAPRGTAQAEESPFVGNPGNITGARGLTGTLRCLQVQGEPEVHVHL  
 60 RDGQILELADSTQTQVPLGEDEQDDWIVVSQLRITSLQLSDTGQYQCLVFLGHQTFVSPQGYVGLLEGLPY  
 FLEPEDRTVAANTPFNLSCQAQGPPEPVDLLWLQDAVPLATAPGHGPQRS LHVPGLNKTSSFSCEAHNA

KGVTTSTRATITIVLPQQPRNLHLVSRQPTELEVAVTPGLSGIYPLTHCTLQAVLSDDGMGIQAGEPDPPE  
 EPLTSQASVPPHQLRLGSLHPHTPYHIRVACTSSQGPSSWTHWLPVETPEGVPLGPPENISATRNGSQAF  
 VHWQEPRAPLQGTLLGYRLAYQQQDTPPEVLMDIGLRQEVTLLEQDGSVSNLTVCAAYTAAGDGWWSLP  
 VPLEAWRPVKEPSTPAFSWPWWYVLLGAVVAAACVLILALFLVHRRKKETRYGEVFEPTVERGELVVRYR  
 5 VRKSYSRRTEATLNSLGISEELKEKLRDVMVDRHKVALGKTLGEGEFGAVMEGQLNQDDSIKVAVKTM  
 KIAICTRSELEDFLSEAVCMKEFDHPNVMRLIGVCFQGSERESFPAPVVILPFMKHGDLSFLLYSRLGD  
 QPVYLPQTQMLVKFMADIASGMEYLSTKRFIHRDLAARNCMLNENMSVCVADFGLSKKIYNGDYRQGRIA  
 KMPVKWIAIESLADRVYTSKSDVWSFGVTMWEIATRGQTPYPGVENSEIYDYLRLQGNRLKQPADCLDGLY  
 10 ALMSRCWELNPQDRPSFTELREDLENTLKALEPPAQEPDEILYVNMDEGGGYPEPPGAAGGADPPTQPDPK  
 DSCSCLTAAEVHPAGRYVLCPSSTPSPAQPADRGSPAAPGQEDGA

# Tubulin cofactor D

15 GH1-13-PCR-G3F1  
 CTTCCGCAGCAGGGTCTGGTTGCTCTCAGGGAGTCTGCAGCCATCGAGGCACCTGAGGACAGTGGCAGCATA  
 GGGCAAACAGTCTTCACGTTTTCCATGTTTAAATATTTGTGCCAGGGCCTGCAGCGTCCCATCCATGGTGTAT  
 GACCCCTGCATGGTCTGGAAGGAGGAACGGGCCAGATTGCACAGGCTCCAGTCCAGGAACCTCAGCCATCTT  
 20 GCTTTGCTTGACATCAGGACGTGTGATAAATCTGGACACAAGGACAGCAG  
 >gi|8400735|ref|NM\_005993.2| Homo sapiens tubulin-specific chaperone d  
 (TBCD), mRNA  
 TAGCGGGCGCCTCCTTTTCATCCCTCATCCTTCATCCCTGGCTTTTCGCGCTCTAGCGGAGTGGGATCTGC  
 GAACACGTGAGGCGGGGCGCGGTCCCAGGCTGCCGAGATGGCCCTGAGCGACGAACCGGCCGCGGGTG  
 25 GCCCGAGGAGGAGGCGGAGGACGAGACACTGGCCTTTGGCGCGGCGCTGGAAGCGTTCCGCGAGAGCGC  
 GGAGACCCGGGCGCTGCTGGGCCGCTGCGGGAGGTGCACGGCGGCGGCGCGGAGCGCGAGGTGGCCCTG  
 GAGCGGTTCCGCGTAATAATGGACAAATACCAGGAGCAGCCTCATCTGTTGGACCCGCACCTTGAATGGA  
 TGATGAACCTGTTGTTGGACATAGTGCAAGATCAGACATCTCCAGCTTCCCTTGTACATCTGGCTTTTAA  
 ATTTCTTTACATCATCACAAGGTTTCGAGGCTATAAAACATTTCTTCGTTTATTTCCTCATGAAGTTGCC  
 30 GATGTAGAGCCTGTTTTAGATTTGGTCAACAATTCAGAAATCCCAAGGACCATGAAGCTTGGGAAACCCGCT  
 ACATGCTTTTGCTCTGGCTCTCCGTGACCTGCCTGATCCCTTTTGATTTTCTCGCCTTGACGGGAACCT  
 CCTCACCCAGCCTGGGCAAGCACGAATGTCCATAATGGACCGTATTCTCAAATAGCAGAGTCCTACTTG  
 ATTGTCACTGACAAGGCCCGAGATGCAGCTGCTGTCCTTGTGTCCAGATTTATCACACGTCCTGATGTCA  
 AGCAAAGCAAGATGGCTGAGTTCCTGGACTGGAGCCTGTGCAATCTGGCCCGTTTCTCCTTCCAGACCAT  
 35 GCAGGGGGTCATCACCATGGATGGGACGCTGCAGGCCCTGGCACAAATATTTAAACATGGAAAAACGTGAA  
 GACTGTTTGCCCTATGCTGCCACTGTCTCAGGTGCCTCGATGGCTGCAGACTCCCTGAGAGCAACCAGA  
 CCCTGCTGCGGAAGCTGGGGGTGAAGCTTGTGCAGCGACTGGGGCTGACATTCTGAAGCCGAAGGTGGC  
 AGCATGGAGGTACCAGCGTGGCTGCCGATCTTTGGTGCAAAATCTGCAGCTCCTCACTCAGGGTCAGAGT  
 GAGCAGAAGCCACTCATCCTGACCGAAGATGACGAGCAGAAAGATGACGACGTCCAGAGGGGGTGGAGCGTG  
 40 TGATAGAGCAGCTGCTGGTGGGCTGAAGGACAAGGACACGGTCGTGCGGTGGTCTGCAGCCAAGGGCAT  
 CGGTAGGATGGCTGGCAGGCTTCCCAGAGCCCTGGCGGATGATGTGGTGGGTCTGTGCTGGACTGCTTC  
 AGTTTCCAGGAGACTGACAAGGCGTGGCATGGGGGATGTCTGGCGCTGGCAGAGCTGGGCAGGAGAGGCC  
 TGTTGCTGCCGTCTCGACTCGTGGATGTTGTGCGCGTGATCCTGAAGGCGCTGACCTACGACGAGAAGCG  
 GGGTGCTGACGCGTGGGCACCAACGTCAGGGACGCCGCTGCTACGTGTGCTGTGCCTTCGCGCGTGCC  
 45 TATGAGCCTCAGGAGCTGAAGCCCTTTGTGACTGCAATCTCGAGTGCAGTGGTGATTGCTGCGGTGTTTG  
 ACCGAGACATAAACTGCAGAAGAGCAGCCTCTGCCGCCCTTCCAGGAGAATGTGGGGAGACAGGGCACTTT  
 CCCTCATGGTATTGATATTTTGACCACAGCTGACTATTTTGGCGTCGGTAACAGATCCAAGTGTTCCTG  
 GTTATAAGTGTGTTTATTGCGCGCTTTCCTGAGTACACGCAGCCAATGATAGACCACCTGGTTACCATGA  
 AGATCAGCCACTGGGATGGGGTCATCCGAGAGTTGGCTGCGAGGGCGCTGCACAACCTGGCCCAGCAGGC  
 50 ACCCGAGTTCAGCGCCACGCAAGTCTTCCCAGAGGCTGCTGTCCATGACACTGAGTCCAGATCTTCACATG  
 AGGCATGGGTGATTCTCGCCTGCGCAGAAGTTGCTTACGCCTTGTACAACTTGCAGCCCCAAGAGAACA  
 GGCCCGTCACGGACCATCTGGACGAGCAGGCAGTGCAGGGCCTGAAGCAGATTACCAGCAGCTCTATGA  
 TCGTCAGTTATACAGGGGTCTGGGAGGACAGCTCATGAGACAAGCAGTGTGTGTTTAAATAGAAAAGTTG  
 TCACTTTCCAAAATGCCCTTTAGAGGTGACACCGTAATTGATGGTTGGCAATGGCTGATAAATGACACTT  
 55 TGAGACATCTCCATCTCATCTCAAGTCACTCCCGCCAGCAGATGAAGGATGCAGCAGTCTCGGCCCTGGC  
 TGCTCTATGCAGTGAATATTACATGAAGGAGCCGGGGGAGGCAGATCCCGCAATTCAGGAGGAGCTGATC  
 ACGCAGTACCTGGCTGAGCTTCGGAACCCGAGGAGATGACTCGCTGTGGCTTCTCGTTGGCCTTGGCGC  
 CCCTTCCAGGCTTCTTCTGAAAGGCCGGCTCCAGCAGGTTCTCACAGGTTTAAGAGCAGTTACCCACAC  
 TCCCCCGAGGACGTAAGTTTGTGAGTCCAGGAGAGACGGCTTGAAGGCCATTGCGAGGATTGCGCAG  
 60 ACTGTTGGTGTGAAAGCAGGAGCCCCAGACGAAGCTGTGTGCGGAGAGAATGTTTCCAGATTTACTGTG  
 CGCTGCTGGGCTGCATGGACGACTACACCACGGACAGCAGAGGGGACGTGGGCACCTGGGTCCGCAAGGC

CGCCATGACCAGTCTGATGGATCTGACACTTCTGCTGGCTCGGAGCCAGCCTGAGCTGATCGAGGCCCAT  
 ACCTGTGAGCGCATCATGTGCTGTGTGGCCCAGCAGGCCAGTGAGAAGATTGACCGTTTCCGTGCTCACG  
 CCGCCAGCGTGTTCCTGACGCTCCTGCACTTTGACAGCCCTCCCATCCCCACGTGCCCCACCGAGGAGA  
 ACTGGAAAAGCTGTTTCCAGGTCCGATGTGGCCTCCGTGAAGTGGAGTGCACCTTCCCAGGCCTTCCCA  
 5 CGCATACCCAGCTCCTTGGGCTGCCACCTACCGCTACCACGTCCTGCTGGGGCTAGTCGTGTCCCTGG  
 GCGGCTTGACGGAGTCGACGATCCGGCACTCCACCCAGAGCCTCTTTGAGTACATGAAGGGCATTGAGAG  
 CGACCCGCGAGGCCCTGGGCAGCTTCAGCGGGACCCTTCTGCAGATCTTTGAGGACAACCTTCTGAATGAG  
 AGGGTGTCCGTGCCGCTGCTGAAGACGCTGGACCACGTGCTCACCCACGGCTGCTTCGACATCTTCACCA  
 CGGAGGAGGACCACCCCTTTGCTGTGAAGTTGCTTGCGCTCTGTAAGAAAGAAATCAAGAATTCAAAAGA  
 10 TATCCAGAAGCTCCTGTGAGGCATCGCAGTGTTCTGCGGGATGGTGAGTTCCCCGGCGACGTGAGGAGG  
 CAGGCCCTCCTGCAGCTGTGTCTGCTCCTGCTGCCACCGTTTCCCGCTGATCCGGAAGACCACGGCCAGCC  
 AGGTGTACGAGACATTGCTCACCTACAGTCAGCTCGTGGGCGCGGATGTGCTGGACGAGGTGGTGACTGT  
 GCTCAGTGACACTGCGTGGGACGCGGAGCTTGCACTGGTGAGAGAGCAGCGCAACCGTCTGTGTGACCTT  
 CTGGGCGTACCCAGGCCCCAGCTGGTGCCCCAGCCTGGTGCTGCTGAAGCCAGTCTTGAGGCCCATACC  
 15 TCACCCCTGCCTGGTGAGGATGTCTTGTTCCTGAGGGAGGCCGGTGTGGAAAGCCTTGCACAGTGGTGCC  
 TCCAGCTGTTGAAGGGTAGCGCTGGCCCTTGGAGGCTGGCACTAGCTGACAGCTTTTCTCTCTGCACCT  
 GCGCTCTGGTGACTTGGGGTGGACGCCTCTGCCTTCACTTGAACACAAATGTGCTTCTATAAAATCATG  
 TACCAAG

20 >gi|8400736|ref|NP\_005984.2| beta-tubulin cofactor D [Homo sapiens]  
 MALSDPEAAGGPEEEAEDETAFGAALAEAFGESAEETRALLGRLREVHGGGAEREVALERFRVIMDKYQEQ  
 PHLLDPHLEWMMNLLLDIVQDQTSPASLVHLAFKFLYIITKVRGYKTFLRFPHEVADVEPVLDLVTIQN  
 PKDHEAWETRYMLLLWLSVTCLIPDFSRDLGNLLTQPGQARMSIMDRILQIAESYLIVSDKARDAAVL  
 VSRFITRPDVKQSKMAEFLDWSLCNLARSSFQTMQGVITMDGTLQALAQIFKHGKREDCLPYAATVLRCL  
 25 DGCRLPESNQTLRLKLVQRLGLTFLKPKVAAWRYQRCRSLAANLQLLTQGGSEQKPLILTEDDDE  
 DDDVPEGVERVIEQLLVGLKDKDTVVRWSAAKGIGRMAGRLPRALADDVGVSLDCFSFQETDKAWHGGC  
 LALAEELGRRGLLLPSRLVDVVAVILKALTYDEKRGACSVGTNVRDAACYVCCAFARAYEPQELKPFVTAI  
 SSALVIAAVFDRDINCRRAASAAEQENVGRQGTFFPHGIDILTTADYFAVGNRSNCFVLVISVFIAGFPEYT  
 QPMIDHLVTMKISHWDGVIRELAARALHNLAAQAPESATQVFPRLLSMTLSPDLHMRHGSILACAEVAY  
 30 ALYKLAQENRPVTDHLDEQAVQGLKQIHQQLYDRQLYRGLGGQLMRQAVCVLIEKLSLSKMPFRGDTV  
 DGWQWLINDTLRHLHLISSHRQQMKDAAVSALAALCSEYYMKEPGEADPAIQEELITQYLAELRNPEEM  
 TRCGFSLALGALPGFLLKGRLLQVLTGLRAVTHTSPEVDVSFAESRRDGLKAIARICQTVGVKAGAPDEAV  
 CGENVSQIYCALLGCMDDYTTDSRGDVGTVWRKAAMTSLMDLTLLLARSQPELIEAHTCERIMCCVAQQA  
 SEKIDRFRAHAASVFLTLHLHFDSPPIPHVPHRGELEKLFPRSDVASVNWSAPSQAFPRITQLLGLPTYRY  
 35 HVLLGLVSVLGGTTESTIRHSTQSLFEYMKGIQSDPQALGSFSGTLLQIFEDNLLNERVSVPLLKTLHDV  
 LTHGCFDIFTTEEDHPFAVKLLALCKKEIKNSKDIQKLLSGIAVFCGMVQFPQGDVRRQALLQLCLLLCHR  
 FPLIRKTTASQVYETLLTYSDVVGADVLDEVTVLSDTAWDAELAVVREQRNRLCDLLGVPRPQLVPQPG  
 AC

40 Transglutaminase 2

GH1-173-PCR-G3F1  
 CCAGTGTGCTTGGGTTCTGCGGCACCCCTGGATCTCCCCAAACTCATTGCGGAAGTACTCGATGAGAAGGTTG  
 45 CTGTTCTGGTCATGGGCCGAGTTGTAGTTGGTCACGACGCGGGTAGGGATGCCAGGCACCTCAGCACTGTG  
 CAGGCCACGGCGGCAAGACCCAGCACTGGCCATACTTGACGCGCTGGCAGCCGTGGTTCTTCCAGCGCCGCA  
 GGATGTCCACGCTGCCGATCCAGGACATGGGGCTGACCCGCAGACCCAGCACAGTGGGTAGATGATAAAGCG  
 GCCGCTCGACTAGTCTGAGGTCTGATACTCACTGACTGTCGTA

50 gi|20141877|sp|P21980|TGM2\_HUMAN Protein-glutamine gamma-  
 glutamyltransferase (Tissue transglutaminase) (TGase C) (TGC) (TG(C))  
 (Transglutaminase 2) (TGase-H)  
 MAEELVLERCDELETNGRDHHTADLCREKLTVRRGQPFWLTLHFEGRNYESVDSLTFSSVVTGPAPSQE  
 AGTKARFPLRDAVEEGDWTATVVDQDQCTLQSLTPANAPIGLYRLSLEASTGYQGSSFVLGHFILLFN  
 55 AWC PADAVYLDSEERQEYVLTQQGFIYQGSAKFIKNIPWNFGQFEDGILDICLILLDVNPKFLKNAGR  
 CSRSSPVYVGRVSGMVNNDQGVLLGRWDNNYGDGVSMSWIGSVLILRRWKNHGCQRVKYQGQWVF  
 AAVACTVLRCLGIPTRVVTNYSNLDQNSNLLIEYFRNEFGEIQGDKSEMIWNFNHCVESWMTRPDLQPG  
 YEGWQALDPTPQEKSEGTYYCCGPVPVRAIKEGDLSTKYDAPFVFAEVNADVVDWIQQDDGSVHKSINRSL  
 IVGLKISTKSVGRDEREDITHYKYPEGSSEEREAFTRANHLNKLAEKEETGMAMRIRVGQSMNMGSDFD  
 60 VFAHITNNTAEYVCRLLLCARTVSYNGILGPECCTKYLLNLNLEPFSEKSVPLCILYEKYRDCLTESNL  
 IKVRALLVEPVINSYLLAERDLYLENPEIKIRILGEPKQKRKLVAEVSLLQNPLPVALEGCTFTVEGAGLT

EEQKTVEIPDPVEAGEEVKVRMDLLPLHMGHLKLVNPFESDKLKAVKGFRNVIIGPA

>gi|4759227|ref|NM\_004613.1| Homo sapiens transglutaminase 2 (C polypeptide, protein-glutamine-gamma-glutamyltransferase) (TGM2), mRNA

5 AACAGGCGTGACGCCAGTTCTAAACTTGAAACAAAACAACTTCAAAGTACACCAAAATAGAACCTCCT  
TAAAGCATAAATCTCACGGAGGGTCTCGGCCGCCAGTGGAGAGGCCACCGCCCCCGCCCCGACCATGGC  
CGAGGAGCTGGTCTTAGAGAGGTGTGATCTGGAGCTGGAGACCAATGGCCGAGACCACCACACGGCCGAC  
CTGTGCCGGGAGAAGCTGGTGGTGCAGCGGGCCAGCCCTTCTGGCTGACCCTGCACTTTGAGGGCCGCA  
10 ACTACCAGGCCAGTGTAGACAGTCTCACCTTCAGTGTCTGTGACCGGCCAGCCCCCTAGCCAGGAGGCCGG  
GACCAAGGCCCGTTTTCCACTAAGAGATGCTGTGGAGGAGGGTGAAGTGGACAGCCACCGTGGTGGACCAG  
CAAGACTGCACCCCTCTCGCTGCAGCTCACCCCGGCCAACGCCCCCATCGGCCTGTATCGCCTCAGCC  
TGGAGGCCTCCACTGGCTACCAGGGATCCAGCTTTGTGCTGGGCCACTTCATTTTGCTCTTCAACGCCTG  
GTGCCAGCGGATGCTGTGTACCTGGACTCGGAAGAGGAGCGGCAGGAGTATGTCCTCACCCAGCAGGGC  
15 TTTATCTACCAGGGCTCGGCCAAGTTCATCAAGAACATACCTTGGAAATTTTGGGCAGTTTCAAGATGGGA  
TCCTAGACATCTGCCTGATCCTTCTAGATGTCAACCCCAAGTTCCTGAAGAAGCCGGCCGTGACTGCTC  
CCGGCGCAGCAGCCCCGTCTACGTGGGCCGGGTGGGTAGTGGCATGGTCAACTGCAACGATGACCAGGGT  
GTGCTGCTGGGACGCTGGGACAACAACCTACGGGGACGGCGTCAGCCCCATGTCTGGATCGGCAGCGTGG  
ACATCCTGCGGCGCTGGAAGAACCACGGCTGCCAGCGCGTCAAGTATGGCCAGTGTGGGTCTTCGCCGC  
CGTGGCCTGCACAGTGTGAGGTGCCTAGGCATCCCTACCCGCGTCTGTGACCAACTACAACCTCGGCCAT  
20 GACCAGAACAGCAACCTTCTCATCGAGTACTCCGCAATGAGTTTGGGGAGATCCAGGGTGACAAGAGCG  
AGATGATCTGGAACCTTCCACTGCTGGGTGGAGTCTGTGGATGACCAGGCCGGACCTGCAGCCGGGTACGA  
GGGCTGGCAGGCCCTGGACCCAACGCCCCAGGAGAAGAGCGAAGGAACGTAAGTGTGTGGCCAGTTCCA  
GTTCTGTGCCATCAAGGAGGGCGACCTGAGCACCAGTACGATGCGCCCTTTGTCTTTGCGGAGGTCAATG  
CCGACGTGGTAGACTGGATCCAGCAGGACGATGGGTCTGTGCACAAATCCATCAACCGTTCCCTGATCGT  
25 TGGGCTGAAGATCAGCACTAAGAGCGTGGGCCGAGACGAGCGGGAGGATATCACCCACACCTACAAATAC  
CCAGAGGGGTCTCAGAGGAGAGGGAGGCCCTTCCAAAGGGCGAACCACCTGAACAAACTGGCCGAGAAGG  
AGGAGACAGGGATGGCCATGCGGATCCGTGTGGGCCAGAGCATGAACATGGGCAGTGAAGTTCGCGTCTT  
TGCCACATCACCAACAACACCGCTGAGGAGTACGTCTGCCGCCCTCCTGCTCTGTGCCCGCACCGTCAAG  
30 TACAATGGGATCTTGGGGCCCCGAGTGTGGCACCAGTACCTGTCTCAACCTAACCTGGAGCCTTTCTCTG  
AGAAGAGCGTTTCTCTTTGTCATCCTCTATGAGAAATACCGTGAAGTGCCTTACGGAGTCCAACCTCATCAA  
GGTGCAGGCCCTCCTCGTGGAGCCAGTTATCAACAGCTACCTGCTGGCTGAGAGGGACCTCTACCTGGAG  
AATCCAGAAATCAAGATCCGGATCCTTGGGGAGCCCAAGCAGAAACGCAAGCTGGTGGCTGAGGTGTCCC  
TGCAGAACCCGCTCCCTGTGGCCCTGGAAGGCTGCACCTTCACTGTGGAGGGGGCCGGCCTGACTGAGGA  
GCAGAAAGACGGTGGAGATCCAGACCCCGTGGAGGCAGGGGAGGAAGTTAAGGTGAGAATGGACCTCGTG  
35 CCGCTCCACATGGGCCTCCACAAGCTGGTGGTGAACCTTCAGAGCGACAAGCTGAAGGCTGTGAAGGGCT  
TCCGGAATGTCATCATTGGCCCCCGCTAAGGGACCCCTGCTCCAGCCTGCTGAGAGCCCCACCTTGAT  
CCCAATCCTTATCCCAAGCTAGTGAGCAAAATATGCCCTTATTGGGCCCCAGACCCAGGGCAGGGTGG  
GCAGCCTATGGGGCTCTCGGAAATGGAATGTGCCCCTGCCCCATCTCAGCCTCCTGAGCCTGTGGGTCC  
CCACTCACCCCTTTGCTGTGAGGAATGCTCTGTGCCAGAAACAGTGGGAGCCCTGACCTGTGCTGACTG  
40 GGGCTGGGGTGGAGAGGAAAGACCTACATTCCCTCTCCTGCCAGATGCCCTTTGGAAAGCCATTGACC  
ACCCACCATATTGTTTGTATCTACTTCATAGCTCCTTGGAGCAGGCAAAAAGGGACAGCATGCCCTTGGC  
TGGATCAGGAATCCAGCTCCCTAGACTGCATCCCGTACCTCTTCCCATGACTGCACCCAGCTCCAGGGC  
CCTTGGGACACCCAGAGCTGGGTGGGGACAGTGATAGGCCAAGGTCCCCTCCACATCCAGCAGCCCAA  
GCTTAATAGCCCTCCCCCTCAACCTCACCATTGTGAAGCACCTACTATGTGCTGGGTGCCTCCACACTT  
45 GCTGGGGCTCACGGGGCTCCAACCCATTTAATCACCATGGGAAACTGTTGTGGGCGCTGCTTCCAGGAT  
AAGGAGACTGAGGCTTAGAGAGAGGAGGCAGCCCCCTCCACACCAGTGGCCTCGTGGTTATAAGCAAGGC  
TGGGTAATGTGAAGGCCAAGAGCAGAGTCTGGGCCTCTGACTCTGAGTCCACTGCTCCATTATAACCC  
CAGCCTGACCTGAGACTGTGCGCAGAGGCTGTCTGGGGCTTTATCAAAAAAAGACTCAGCCAAGACAAGG  
AGGTAGAGAGGGGACTGGGGGACTGGGAGTCAGAGCCCTGGCTGGGTTCAGGTCCACGTCTGGCCAGCG  
50 ACTGCCTTCTCCTCTCTGGGCCTTTGTTTCTTGTGGTTCAGAGGAGTGATTGAACCTGCTCATCTCCAA  
GGATCCTCTCCACTCCATGTTTGAATACACAATTC

>gi|4759228|ref|NP\_004604.1| transglutaminase 2 (C polypeptide, protein-glutamine-gamma-glutamyltransferase) [Homo sapiens]

55 MAEELVLERCDLELETNGRDHHTADLCREKLVVRRGQFPWLTLLHFEGRNYSQASVDSLTFVSVTGPAPSQE  
AGTKARFPLRDAVEEGDWATVVDQDCTLSLQLTTPANAPIGLYRLSLEASTGYQGSSFVLGHFILLFN  
AWCPADAVYLDSEERQEYVLTQQGFYQGSAPFIKNIPWNFGQFQDGIIDICLILLDVNPKFLKNAGRD  
CSRSSPVYVGRVSGMVNNDQGVLLGRWDNNYGDGVSMSWIGSVDILRRWKNHGCQRVKYQGCWVF  
AAVACTVLRCLGIPTRVVTNYSNHDQNSNLLIEYFRNEFGEIQDKSEMIWNFHCWVESWMTRPDLQPG  
60 YEGWQALDPTPQEKSEGTYCCGPVPVRAIKEGDLSTKYDAPFVFAEVNADVVDWIQDDGSVHKSINRSL  
IVGLKISTKSVGRDEREDITHYKYPEGSSSEEREAFTRANHLNKLAEKEETGMAMRIRVQSMNMGSDFD

VFAHITNNTAEYVCRLLLCARTVSYNGILGPECGTKYLLNLTLEPFSEKSVPLCILYEKYRDCLTESNL  
IKVRALLVEPVINSYLLAERDLYLENPEIKIRILGEPKQKRKLVAEVS LQNPLPVALEGCTFTVEGAGLT  
EEQKTVEIPDPVEAGEEVKVRMDLVPLHMLHKLKLVNFESDKLKAVKGFNRNVIIGPA

- 5 >gi|13097680|gb|BC003551.1|BC003551 Homo sapiens, Similar to  
transglutaminase 2 (C polypeptide, protein-glutamine-gamma-  
glutamyltransferase), clone MGC:1193 IMAGE:3544757, mRNA, complete cds  
CTCCGCCTCGGCAGTGCCAGCCGCCAGTGGTCGCACCTTGGAGGGTCTCGCCGCCAGTGGAAAGGAGCCACC  
GCCCCCGCCCCGACCATGGCCGAGGAGCTGGTCTTAGAGAGGTGTGATCTGGAGCTGGAGACCAATGGCCG  
10 AGACCACCACACGGCCGACCTGTGCCGGGAGAAGCTGGTGGTGGTGGACGGGGCCAGCCCTTCTGGCTGACC  
CTGCACCTTTGAGGGCCGCAACTACGAGGCCAGTGTAGACAGTCTCACCTTCAGTGTCTGTCGACCGGCCAG  
CCCCTAGCCAGGAGGCCGGGACCAAGGCCCGTTTCCACTAAGAGATGCTGTGGAGGAGGGTGACTGGAC  
AGCCACCGTGGTGGACCAGCAAGACTGCACCCCTCTCGCTGCAGCTCACCAACCCCGGCCAACGCCCCCATC  
GGCCTGTATCGCCTCAGCCTGGAGGCCTCCACTGGCTACCAGGGATCCAGCTTTGTGCTGGGCCACTTCA  
15 TTTTGTCTTTCAACGCCTGGTGCCAGCGATGCTGTGTACCTGGACTCGGAAGAGGAGCGGCAGGAGTA  
TGTCTCACCCAGCAGGGCTTTATCTACCAGGGCTCGGCCAAGTTCATCAAGAACATACCTTGGAATTTT  
GGGCAGTTTGAAGATGGGATCCTAGACATCTGCCTGATCCTTCTAGATGTCAACCCCAAGTTCCTGAAGA  
ACGCCGGCCGTGACTGCTCCCGCCGACGAGCCCCGTCTACGTGGGCCGGGTGGTGGAGTGGCATGGTCAA  
CTGCAACGATGACCAAGGTGTGCTGCTGGGACGCTGGGACAACAACCTACGGGGACGGCGTCAGCCCCATG  
20 TCCTGGATCGGACGCTGGACATCCTGCGCGCTGGAAGAACCAGGCTGCCAGCGCTCAAGTATGGCC  
AGTGCTGGGTCTTCGCGCCGCTGGCCTGCACAGTGTCTGAGGTGCCTGGGCATCCCTACCGCGCTCGTGAC  
CAACTACAACCTCGGCCCATGACCAGAACAGCAACCTTCTCATCGAGTACTTCCGCAATGAGTTTGGGGAG  
ATCCAGGGTGACAAGAGCGAGATGATCTGGAACCTTCCACTGCTGGGTGGAGTCTGATGACAGGCCGG  
ACCTGCAGCCGGGTACGAGGGCTGGCAGGCCCTGGACCCAACGCCCCAGGAGAAGAGCGAAGGGACGTA  
25 CTGCTGTGGCCAGTTCCAGTTCGTGCCATCAAGGAGGGCGACCTGAGCACCAAGTACGATGCGCCCTTT  
GTCTTTGCGGAGGTCAATGCCGACGTGGTAGACTGGATCCAGCAGGACGATGGGTCTGTGCACAAATCCA  
TCAACCGTTCCCTGATCGTTGGGCTGAAGATCAGCACTAAGAGCGTGGGCCGAGACGAGCGGGAGGATAT  
CACCCACACCTACAAATACCCAGAGGGGTCTCAGAGGAGAGGGAGGCCTTCACAAGGGCGAACCACCTG  
AAACAACCTGGCCGAGAAGGAGGAGACAGGGATGGCCATGCGGATCCGTGTGGGCCAGAGCATGAACATGG  
30 GCAGTACTTTGACGTCTTTGCCACATCACCAACAACACCGCTGAGGAGTACGTCTGCCGCTCCTGTGCT  
CTGTGCCCGCACCGTCAGCTACAATGGGATCTTGGGGCCGAGTGTGGCACCAAGTACCTGCTCAACCTC  
AACCTGGAGCCTTTCTCTGGTAAAGCCCTGTGTTTCTGGAGCATTTGTTGACCGCCAACCTGACAACATGC  
TAGGTAGTGACCTAAAAAAAAAAAAAAAAAAAA
- 35 >gi|13097681|gb|AAH03551.1|AAH03551 Similar to transglutaminase 2 (C  
polypeptide, protein-glutamine-gamma-glutamyltransferase) [Homo sapiens]  
MAEELVLERCDLELETNDRDHTADLCREKLVRVRRGQPFWLTLHFEGRNYESVDSLTFSSVVTGPAPSQE  
AGTKARFPLRDAVEEGDWTATVVDQDCTLSLQLTTPANAPIGLYRLSLEASTGYQSSFVLGHFILLFN  
40 AWC PADAVYLDSEERQEYVLTQQGFYQGS AKFIKNI PWNFGQFEDGILDICLILLDVNPKFLKNAGRD  
CSRSSPVYVGRVSGMVNCDNDQGVLLGRWDNNYGDGVSPMSWIGSVDILRRWKNHGCQRVKYQGCWVF  
AAVACTVLRCLGIPTRVVTNYS AHDQNSNLLIEYFRNEFGEIQDKSEMIWNFHCWVESWMTRPDLQPG  
YEGWQALDPTPQEKSEGTYCCGPVPVRAIKEGDLSTKYDAPFVFAEVNADVVDWIQQDDGSVHKSINRSL  
IVGLKISTKSVGRDEREDITHYKYPEGSSEEREAFTRANHLNKLAEKEETGMAMRIRVGQSMNMGSDFD  
VFAHITNNTAEYVCRLLLCARTVSYNGILGPECGTKYLLNLTLEPFSKALCSWSIC
- 45 >gi|339577|gb|M98478.1|HUMTGH1A Human transglutaminase mRNA, complete  
cds  
CAGGCGTGACGCCAGTTCTAAATCTTGAAACAGAACAAAACCTTCAAAGTACACCAAAATAGAACCTCCTT  
AAAGCATAAATCTCACGGAGGGTCTCGCCGCCAGTGGAAAGGAGCCACCGCCCCCGCCGACCATGGCCGA  
50 GGAGCTGGTCTTAGAGAGGTGTGATCTGGAGCTGGAGACCAATGGCCGAGACCACACACGGCCGACCTG  
TGCCGGGAGAAGCTGGTGGTGGCAGCGGGCCAGCCCTTCTGGCTGACCCTGCACCTTGTAGAGGCCGCAACT  
ACGAGGCCAGTGTAGACAGTCTCACCTTCAGTGTCTGACCGGCCAGCCCTAGCCAGGAGGCCGGGAC  
CAAGGCCCGTTTTTCCACTAAGAGATGCTGTGGAGGAGGGTGACTGGACAGCCACCGTGGTGGACCAGCAA  
GACTGCACCCCTCTCGCTGCAGCTCACCAACCCCGGCCAACGCCCCCATCGGCCTGTATCGCCTCAGCCTGG  
55 AGGCCTCCACTGGCTACCAGGGATCCAGCTTTGTGCTGGGCCACTTCATTTTGCTCTTCAACGCCTGGTG  
CCGACCGGATGTGTGTACCTGGACTCGGAAGAGGAGCGGCAGGAGTATGTCTCTACCCAGCAGGGCTTT  
ATCTACCAGGGCTCGGCCAAGTTCATCAAGAACATACCTTGAATTTTGGGCAGTTTGAAGATGGGATCC  
TAGACATCTGCCTGATCCTTCTAGATGTCAACCCCAAGTTCCTGAAGAACGCCGGCCGTGACTGCTCCCG  
CCGCAGCAGCCCCGTCTACGTGGGCCGGGTGTGGAGTGGCATGGTCAACTGCAACGATGACCAGGGTGTG  
60 CTGCTGGGACGCTGGGACAACAACCTACGGGGACGGCGTCAGCCCCATGTCTTGATCGGCAGCGTGGACA  
TCCTGCGGCGCTGGAAGAACCACGGCTGCCAGCGCTCAAGTATGGCCAGTGTGGGTCTTCGCCGCCGT

GGCCTGCACAGTGTCTGAGGTGCCTGGGCATCCCTACCCGCGTCTGTGACCAACTACAACCTCGGCCCATGAC  
 CAGAACAGCAACCTTCTCATCGAGTACTTCCGCAATGAGTTTGGGGAGATCCAGGGTGACAAGAGCGAGA  
 TGATCTGGAACCTTCCACTGCTGGGTGGAGTCTGTGGATGACCAGGCCGACCTGCAGCCGGGGTACGAGGG  
 CTGGCAGGCCCTGGACCAACGCCCCAGGAGAAGAGCGAAGGGACGTACTGCTGTGGCCAGTTCCAGTT  
 5 CGTGCCATCAAGGAGGGCGACCTGAGCACCAAGTACGATGCGCCCTTTGTCTTTGCGGAGGTCAATGCCG  
 ACGTGGTAGACTGGATCCAGCAGGACGATGGGTCTGTGCACAAATCCATCAACCGTTCCTGATCGTTGG  
 GCTGAAGATCAGCACTAAGAGCGTGGGCCGAGACGAGCGGGAGGATATCACCCACACCTACAAATACCCA  
 GAGGGGTCCTCAGAGGAGAGGGAGGCCCTTCACAAGGGCGAACACCTGAACAAACTGGCCGAGAAGGAGG  
 AGACAGGGATGGCCATGCGGATCCGTGTGGGCCAGAGCATGAACATGGGCAGTGACTTTGACGCTTTTGC  
 10 CCACATCACCAACAACACCGCTGAGGAGTACGCTGCGCCCTCCTGCTCTGTGCCCCGACCGTCAGCTAC  
 AATGGGATCTTGGGGCCCCGAGTGTGGCACCAGTACCTGCTCAACCTCAACCTGGAGCCTTTCTCTGGTA  
 AAGCCCTGTGTTCTCGGACATTTGTGTGACCGCCAACCTGACAACATGCTAGGTAGTGACCTAACCACTTA  
 GCATGTGTGATTTACCCACAGACACTTACATGGCGCTGACTCTGGGGCAGGCCCTGTCTTAAGCACTT  
 TATAATATCAACCCACTTA

15 >gi|339578|gb|AAA36739.1| transglutaminase  
 MAEELVLERCDLELETNGRDHHTADLCREKLVVRRGQPFWLTLHFEGRNYEASVDSLTFSSVVTGPAPSQE  
 AGTKARFPLRDAVEEGDWTATVVDQDCTLQLLTPANAPIGLYRLSLEASTGYQGSSFVLGHFILLFN  
 AWC PADAVYLDSEERQEYVLTQQGFIYQGS AKFIKNIPWNFGQFEDGILDICLILLDVNPKFLKNAGR  
 20 CSRSSPVYVGRVWSGMVNCNDDQGVLLGRWDNNYGDGVS PMSWIGSV DILRRWKNHGCQRVKYQGCWVF  
 AAVACTVLRCLGIPTRVVTNYS AHQNSNLLIEYFRNEFGEIQGDKSEMIWNFHCWVESWMT RPDLPQ  
 YEGWQALDPTPQEKSEGT YCCGPVPVRAIKEGDLSTKYDAPFVFAEVNADVVDWIQQDDGSVHKSINRSL  
 IVGLKISTKSVGRDEREDITHYKYPEGSSSEEREAFTRANHLNKLAEKEETGMAMRIRVGQSMNMGSDFD  
 VFAHITNNTAEYVCRLLLCARTVSYNGILGPECCKYLLNLNLEPFSKALCSWSIC

25 Cytosine deaminase

GH1-27-PCR-G3F1  
 CCAGCGGTGGCTCCAGTGTGCTGGTCTGCGGACGTGTGCCATGCGGAGCTGAATGCCATCATGAACAAAAAT  
 30 TCGACCGATGTGAAAGGCTGTAGTATGTATGTTGCCTTGTTCCCTTGTAATGAATGCGCTAAGCTCATCATC  
 CAGGCAGGTATAAAGAAGTGATTTTCTTGTGTTTGGATAAATACCATGATAGTGACGAGGCAACTGCTGCGAG  
 GCTCCTGTTTAATATGGCCGGGGTGACATTCCGGAAATTCATACCGAAGTGCAGCAAGATTGTCATTGACTT  
 TGATTCAATTAACAGCAGACCGAGTCAAAAGCTTCAGTGAGTTACATCTCATTCAATCTCCAGAAGATTGGG  
 ATTATCGTCTTCTAAGAGGTTGCTAATGCCTTTCATCTTGAAGTTACACATAACTTCTTACTAGCCAGTATG  
 35 GCAAAAGTAGGCATCTTAAGAATATAAAGCCTCCAATCTTCCTTACTGTCTCTCTTGTACATGGAATCTAC  
 ATGTGTTTGAATATTGCTTTAGGGATTAAAATAGGGGAGCCTGTGGTGGCCTGGTGCACAGGGGCTAGAA  
 CGAGAGTGCCTCCCCTTCTTGTGTCTCTGGCTGGCTGGGATGCTGTGGCTCTTCAGAGGAGCATCAGCCTGTC  
 TGTATCTGCTGCGATCCGGCAG

40 >gi|23503055|sp|P32321|DCTD\_HUMAN Deoxycytidylate deaminase (dCMP  
 deaminase)  
 MSEVSKKRDDYLEWPEYFMAVAFLSAQRSKDPNSQVGACIVNSENKIVGIGYNGMPNGCSDDVLPWRRRT  
 AENKLDTKYPYVCHAE LNAIMNKNSTDVKGCSMYVALFPCNECAKLIIQAGIKEVIFMSDKYHDSDEATA  
 ARLLFN MAGVTFRKFI PKCSKIVIDFDSINSRPSQKLQ

45 >gi|4503276|ref|NM\_001921.1| Homo sapiens dCMP deaminase (DCTD), mRNA  
 ATGAGTGAAGTTTCTGCAAGAAACGGGACGACTATTTGGAATGGCCAGAGTATTTTATGGCTGTGGCCT  
 TCTTATCAGCACAGAGAAGCAAAGATCCAAATTCAGGTCGGCGCCTGCATCGTGAATTCAGAAAACAA  
 GATTGTGCGGATTGGGTACAATGGGATGCCAAATGGGTGCAGTGATGACGTGTTGCTTGGAGAAGGACA  
 50 GCAGAGAATAAGCTGGACACCAAATACCCGTACGTGTGCCATGCGGAGCTGAATGCCATCATGAACAAAA  
 ATTCGACCGATGTGAAAGGCTGTAGTATGTATGTCGCCTTGTTCCCTTGTAATGAATGCGCTAAGCTCAT  
 CATCCAGGCAGGTATAAAGAAGTGATTTTACGCTCTGATAAATACCATGATAGTGACGAGGCAACTGCT  
 GCGAGGCTCCTGTTTAATATGGCCGGGGTGACATTCCGGAAATTCATACCGAAGTGCAGCAAGATTGTCA  
 TTGACTTTGATTCAATTAACAGCAGACCGAGTCAAAAGCTTCAGTGAGTTACATCTCATTCAATCTCCAG  
 55 AAGATTGGGATTATCGTCTTCTAAGAGGTTGCTAATGCCTTTCATCTTGAAGTTACACATAACTTCTTAC  
 TAGCCAGTATGGCAAAAGTAGGCATCTAAGAATATAAAGCCTCAAATCTTCTTACTGTCTCTCTTGTG  
 ACATGGAACTACATGTGTTTGAACATATTGCTTTAGGATTTAAAATAGGGGAGCCTGTGGTGGCCTGGTG  
 CACAGGGCTAGAACGAGAGTGCCTCCCCTTCTTGTGTCTCTGGCTGGCTGGGATGCTGGTGGCTCTTCAGA  
 GGAGCATCAGCTGTCTGTCTGCTGCGATCCGGCAGCCTCTCTTCACTGCTACATGTGCTGGAAGGAC  
 60 AAATAAATAATTGTGGTTGTGTTCTTAATGGGGACGAGCAGACACACTGATCTGAACATCTGGCCCAAGT  
 GAAGCATGGCATATAGTGCCCTTGAAGAAAATTAGGCCTCAAATGACAGTAGCATTGAAGTGTGCTG

CAGAGTTGAGGGAAACCCCCAGCCACCCTCCCGGAATCCGAGATAGGGTGGCACATCTGTCCTGACAGAC  
 GAGGAGTGTAACGAACCAGGAATATTTCTCCATTTCCTGCTCTCCCACTGCACACAGGGTGGTGGCACA  
 TTATCCCTCTGGGGGGTGGGGACGCTGTGTTTGGCTCAATTTGGGTTTGGTTGGTCACATGGAGCTCT  
 TCCATTTTCGTTTAGCTGAATAATGAGTTGTTCTAGAGGAGACAGCCTGTCTCTCCTTGTGCCCCCAA  
 5 GCCCATGCCCTGCCGTGGTGGCAGCTGGGGCTGTGGATGGGAGGGGTCCCCAACATGGATGTGTTGCCCC  
 TCCTCCGCATGCCAACGCAGTTCATGTACAAGGCCCTCTGCAACTGGAGAGAAAATTAATTCCTATCCC  
 GTGAGTGGATTGTGAGAAATTCCACCCACGTGGAGACAGCTTACTGCAGCACTGTTGGTGTTCGGAGCTC  
 TTCTGTGCCCTGGCTCCATGCTTTACCTACACAAGCATCACCTTCCTAATCACCGCGGGGCGGGGAGCG  
 TGTGGCTGTGCCCCCTTCTCTTTAATCTCATTTAATTTTTATTAAACATGCTCAGTACCTGTGTTGAGAAA  
 10 AGGCTTTCTTTATCCTAAAGATTATTACCTTTTAAAGTGCTCTTATATTTTCATGAGTTTTATTTTGT  
 CTCTGAGATTTGTATTCCACATTCTAGGGTATTCTGTAATTTGGCTCCTTACCAATATTATTAAATCT  
 TATTAAATCT

>gi|4503277|ref|NP\_001912.1| dCMP deaminase [Homo sapiens]  
 15 MSEVSKKRDDYLEWPEYFMAVAFLSAQRSKDPNSQVGACIVNSENKIVGIGYNGMPNGCSDDVLPWRRRT  
 AENKLDTKYPYVCHAEINAIMNKNSTDVKGCSMYVALFPCNECAKLI IQAGIKEVIFTSDKYHDSDEATA  
 ARLLFNMAGVTFRKFI PKCSKIVIDFDSINSRPSQKLQ

Peptidase M41 (Paraplegin)

GH1-40-PCR-G3F1

GTGGAAATGCCCTGTTACTCTGTGGGGATGACGGCAGTGGGCTGGCCATCCTGTGGTATGTTTTCCGTCTG  
 GCCGGGATGACTGGAAGGGAAGGTGGATTCACTGCTTTTAATCAGCTTAAAATGGCTCGTTTACCATTGTG  
 GATTGGAAGATGGGGAAAGGGAGTCAGCTTCAAAGACGTGGCAGGAATGCACGAAGCCAAACTGGAAGTCCG  
 25 CGAGTTTGTGGATTATCTGAAGAGCCCAGAACGCTTCCTCCAGCTTGGCGCCAGGTCCCCAAAGGGCGCACTG  
 CTGCTCGGCCCCCGGCTGTGGGAAGACGCTTCTGGCCAAGGCGGTGGCCACGGAGGCTCAGGTGCCCTTC  
 CTGGCGATGCCGGCCCAGAGTTTCGTGGAGGTTCATTGGAGGCTTCGGCGCTCCCGTGTGCGGAGCCTCTTTAA  
 GGAAGCCCCGAGCCCCGGGCCCCCTGCATCGTCTACATCGATAGATCGACGCGGTGGGCAAGAAGCGCTCCACC  
 ACCATGTCCGGTCTCTCCAACACNAGGAGGAGCAGACGCTCAACCAGCTTCTGGTNAGAAATGGATGGAAA  
 30 TGGGTACCACAGACTGTCATCGTCTGGCGTCCACGAAACCAGCTGACATTTTGGACGGTGCTCTTATA  
 GGCCAGGCCGAACCTGGGACCGGGACGTCTTCTTTGATCTC

>gi|4507172|ref|NM\_003119.1| Homo sapiens spastic paraplegia 7,  
 paraplegin (pure and complicated autosomal recessive) (SPG7), mRNA

TTTCAGGCCAACATGGCCGTGCTGCTGCTGCTGCTCCGTGCCCTCCGCCGGGGTCCAGGCCCGGGTCCCTC  
 35 GGCCGCTGTGGGGCCCAGGCCCGGCTGGAGTCCAGGGTTCCCCGCCAGGCCCGGGAGGGGGCGGCCGTA  
 CATGGCCAGCAGGCCCTCCGGGGACCTCGCCGAGGCTGGAGGCCGAGCTCTGCAGAGCTTACAATTGAGA  
 CTGCTAACCCCTACCTTTGAAGGGATCAACGGATTGTTGTTGAAACAACATTTAGTTCAGAACTCAGTCA  
 GACTCTGGCAACTTTTAGGTGGTACTTTCTATTTTAACACCTCAAGGTTGAAGCAGAAGAATAAGAGAA  
 40 GGATAAGTCGAAGGGGAAGGCGCCTGAAGAGGACGAAGAGGAGAGGAGACGCCGTGAGCGGGACGACCAG  
 ATGTACCGAGAGCGGCTGCGCACCTTGCTGGTTCATCGCGGTTGTTCATGAGCCTCCTGAATGCTCTCAGCA  
 CCAGCGGAGGCAGCATTTCTGGAACGACTTTGTCCACGAGATGCTGGCCAAGGGCGAGGTGCAGCGCGT  
 CCAGGTGGTGCCTGAGAGCGACGTGGTGAAGTCTACCTGCACCCTGGAGCCGTGGTGTGTTGGGCGGCCT  
 CGGCTAGCCTTGATGTACCGAATGCAGGTTGCAAAATATTGACAAGTTTGAAGAGAAGCTTCGAGCAGCTG  
 45 AAGATGAGCTGAATATCGAGGCCAAGGACAGGATCCCAAGTTTCTTACAAGCGAACAGGATTCTTTGGAAA  
 TGCCCTGTACTCTGTGGGGATGACGGCAGTGGGCTGGCCATCCTGTGGTATGTTTTCCGTCTGGCCGGG  
 ATGACTGGAAGGGAAGGTGGATTCACTGCTTTTAATCAGCTTAAAATGGCTCGTTTACCATTGTGGATG  
 GGAAGATGGGGAAAGGAGTCAGCTTCAAAGACGTGGCAGGAATGCACGAAGCCAAACTGGAAGTCCGCGA  
 GTTTGTGGATTATCTGAAGAGCCCAGAACGCTTCCCTCCAGCTTGGCGCCAAAGGTCCCAAAGGGCGCACTG  
 50 CTGCTCGGCCCCCGGCTGTGGGAAGACGCTGCTGGCCAAGGCGGTGGCCACGGAGGCTCAGGTGCCCT  
 TCCTGGCGATGGCCGGCCCAGAGTTTCGTGGAGGTTCATTGGAGGCTCGGCGCTGCCCGTGTGCGGAGCCT  
 CTTTAAGGAAGCCCCGAGCCCCGGGCCCCCTGCATCGTCTACATCGATGAGATCGACGCGGTGGGCAAGAA  
 CGCTCCACCACCATGTCCGGCTTCTCCAACACGAGGAGGAGCAGACGCTCAACCAGCTTCTGGTAGAAA  
 TGGATGGAATGGGTACCACAGACCATGTTCATCGTCTCGGCTCCACGAACCGAGCTGACATTTTGACGG  
 55 TGCTCTGATGAGGCCAGGCCGACTGGACCGGCAGCTCTTTCATTGATCTCCCCACGCTGCAGGAGAGGCGG  
 GAGATTTTTGAGCAGCACCTGAAGAGCCTGAAGCTGACCCAGTCCAGCACCTTTTACTCCCAGCGTCTGG  
 CAGAGCTGACACAGGATTTCAGTGGGGCTGACATCGCCAACATCTGCAATGAGGCTGCGCTGCACGCGGC  
 GCGGGAGGGACACACTTCCGTGCACACTCTCAACTTCGAGTACGCCGTGGAGCGCGTCTCTGCAGGGACT  
 GCCAAAAAGAGCAAGATCCTGTCCAAGGAAGAACAGAAAGTGGTTGCGTTTCATGAGTGGGCCCACGCT  
 60 TGGTGGGCTGGATGCTGGAGCACACGGAGGCGGTGATGAAGGTCTCCATAACCCCTCGGACAAACGCGC  
 CCTGGGCTTTGCTCAGATGCTCCCCAGAGACCAGCACCTCTTACCAAGGAGCAGCTGTTTGAGCGGATG

TGCATGGCCTTGGGAGGACGGGCGCTCGGAAGCACTGTCTTCAACGAGGTCACTTCTGGGGCACAGGACG  
 ACCTGAGGAAGGTCACCCGCATCGCCTACTCCATGGTGAAGCAGTTTGGGATGGCACCCTGGCATCGGGCC  
 CATCTCCTTCCCTGAGGCGCAGGAGGGCCTCATGGGCATCGGGCGGCGCCCTTCAGCCAAGGCCTGCAG  
 CAGATGATGGACCATGAAGCAAGACTGCTGGTGGCCAAGGCCTACAGACACACCGAGAAGGTGCTGCAGG  
 5 ACAACCTGGACAAGTTGCAGGCGCTGGCAAACGCCCTTCTGGAAAAGGAAGTGATAAACTATGAGGACAT  
 TGAGGCTCTCATTGGCCCCGCCGCCCATGGGCCGAAGAAAATGATCGCACCGCAGAGGTGGATCGACGCC  
 CAGAGGGAGAAACAGGACTTGGGCGAGGAGGAGACCGAAGAGACCCAGCAGCCTCCACTTGGAGGCGAAG  
 AGCCGACTTGGCCCAAGTAGTTGGGAGGTGTTGGCTGCACGTGCGGGTGGTCCGGGAAGTGAGGGCTCAC  
 TCAGCCACCCTGAGTTGCTTTTCAGCTGAGGTTTGCACTTCCTCTCGCGGCCCTCAGTAGTCCCTGCACA  
 10 GTGACTTCTGAGATCTGTTGATTGATGACCCTTTTCATGATTTTAAGTTTCTCTGCAGAACTACTGACG  
 GAGTCTGTGTTTTGTGAGTCGTTTCCCCTATGGGGAAGGTTATCAGTGCTTCCCGAGTGAGCATGGAACA  
 CTTTCGAGTTCCCAGGGTTATAGACAGTCGTTCCAGTGTGGCTGAGGCCACCCAGAGGCAGCAGAGCATT  
 CAGACTCCAAACAGACCCCTGTTCATGCCGACGCTTGACGACCGCCCCAGTTCTGTGGCTCCCTCGGA  
 ATGCTAAGGGGATCGGACATGAAAGGACCCCTGTGAGCCGATTGTCCTATCTCCAGCGGCCCTGTCATCCA  
 15 GCTCACTCATCAATGGGGCCAGTCAGGCCCAGGCACTGGGCTCCGGAGGACTCACCCTGCCCCCTGCTG  
 CCATGTGGACTGGTGCAAGTTGAGGACTTCTTGCTGGTCTAGTCACGCATGCAGTGTTGGGGATGCCTTG  
 GTTTTTACTGCTCTGAGAATTGTTGAGATACTTTACTAATAAACTGTGTAGTTGGAAAAAAAAAAAAAA  
 AAAAAA

20 >gi|4507173|ref|NP\_003110.1| paraplegin [Homo sapiens]  
 MAVLLLLLRLRRPGPGPRPLWGPAPSPGFPARPGRGRPYMASRPPGDLAEAGGRALQSLQLRLLTLP  
 TFEGINGLLLLKQHLVQNPVRLWQLLGGTFYFNTSRLKQKNKEKDKSKGKAPEEDEEERRRRERDDQMYRE  
 RLRTLLVIAVMSLLNALSTSGGSISWNDFVHEMLAKGEVQRVQVVPESDVVEVYLHPGAVVFGRPRAL  
 MYRMQVANIDKFEEKLRAAEDELNIEAKDRIPVSYKRTGFFGNALYVGMTAVGLAILWYVFRLAGMTGR  
 25 EGGFSAFNQLKMARFTIVDGKMGKGVSFKDVAGMHEAKLEVREFVDYLKSPERFLQLGAKVPKGALLGP  
 PGCGKTLAKAVATEAQVPFLAMAGPEFVEVIGGLGAARVRSLFKEARARAPCIVYIDEIDAVGKKRSTT  
 MSGFSNTEEEQTLNQLLVEMDGMGTDDHVIVLASTNRADILDGALMRPGRDLDRHVFIDLPTLQERREIFE  
 QHLKSLKLTQSSTFYSQLAEELTPGFSGADIANICNEAALHAAREGHTSVHTLNFEYAVERVLAGTAKKS  
 KILSKEEQKSVAFHESGHALVGWMLLEHTEAVMKVSIPTRNAALGFAQMLPRDQHLFTKEQLFERMCML  
 30 GGRASEALSFNEVTSGAQDDLKRVTRIAYSMVKQFGMAPGIGPISFPEAQEGLMGIRRPFSQGLQQMMD  
 HEARLLVAKAYRHTEKVLQDNLDKLQALANALLEKEVINYEDIEALIGPPPHGPKKMIAPQRWIDAQREK  
 QDLGEEETEETQQPPLGGEEPTWPK

35 CD13 Amino peptidase  
 GH1-72-PCR-G3F1  
 AGGCCAGGCCTAGGGCGGGGTTGGCATGAGCGGGCAGCGCGCTGGGAGGTGCTCAGGCAGCCTGGGTCATCA  
 GGAAGTAGACTGGCTCACAGGCAGAGAGAACGTGGGCTGGAGACTTTGTCCTTGAGGGGAGGACACTGGTG  
 40 CTCGGGCTCCAGGAATGGAGGCCCTGCACCAGCCGCTGGGATGGACACATGTGGGCACCTTGCAATGGGGGCC  
 GGGTGACTTCAAGGGCTGGGGACTATTTGCTGTTTTCTGTGAACCACTGGAGCACCACCTCCTTGTCTCCT  
 TCACCCACTTATGTTGCTTTTCGTCTTCTCCAGGGGCTTGCTCCAGGGCCCGGTGCCTTAGCCGAAGCCTGT  
 TTCCTCGTTTCCT

45 >gi|113743|sp|P15144|AMPN\_HUMAN Amino peptidase N (Microsomal  
 aminopeptidase) (GP150) (Myeloid plasma membrane glycoprotein CD13)  
 MAKGFYISKSLGILGILLGVAAVCTIIALSVMVYSQEKKNANSSPVASTTPSASATNPASATTLQSKA  
 WNRRLPNTLKPDSYQVTLRPYLTPNDRGLYVFKGSSTVRFTCKEATDVIIHSHKLNLYTLSSQHRVLR  
 GVGGSQPPDIDKTELVEPTLYLVVHLKGSVVKDSQYEMDSEFEGELADDLAGFYRSEYMEGNVRKVATT  
 50 QMQAADARKSFPCFDEPAMKAEFNITLIHPKDLTALSNMLPKGPSTPLPEDPNWNVTEFHTTPKMSTYLL  
 AFIVSEFDYVEKQASNGVLIRIWARPSAIAAGHG DYALNVTGPILNFFAGHYDTPYPLPKSDQIGLPDFN  
 AGAMENWGLVTYRENSLLFDPLSSSSSNKERVVTVIAHELAHQWFGNLVTIEWWNLWLNEGFASYEYL  
 GADYAEPTWNLKDLMLVNDVYRMAVDALASSHPLSTPASEINTPAQISELFDALISYSKGASVLRMLSSF  
 LSEDVFKQGLASYLHTFAYQNTIYLNLDHQLQEA VNNRSIQLPTTERDIMNRWTLQMGFPVITVDTSTGT  
 55 LSQEHFLDPDSNVTRPSEFN YVWIVPITSIRDGRQQDYWLMDVRAQNDLFSTSGNEWVLLNLNVTGYY  
 RVNYDEENWRKIQTQLQRDHSAPVINRAQIINDAFNLASAHKVPVTLALNNTLFLIEERQYMPWEAALS  
 SLSYFKLMFDRSEVYGPKNYLKKQVTPLFIHFRNNTNNWREIPENLMDQYSEVNAISTACSNVPECEE  
 MVSGLFKQWMENPNNNPIHPNLRSTVYCNAIAQGGEEWDFAWEQFRNATLVNEADKLRAALACSKELWI  
 LNRYLSYTLNPD LIRKQDATSTIISITNNVIGQGLVWDFVQSNWKKPFNDYGGGSFSFSNLIQAVTRRFS  
 60 TEYELQQLEQFKKDNEETGFGSGTRALEQALEKTKANIKWVKENKEVVLQWFTENSK

>gi|4502094|ref|NM\_001150.1| Homo sapiens alanyl (membrane)  
aminopeptidase (aminopeptidase N, aminopeptidase M, microsomal  
aminopeptidase, CD13, p150) (ANPEP), mRNA

5 TAATTTTGGCCAGTCTGCCTGTTGTGGGGCTCCTCCCCTTTGGGGATATAAGCCCGCCTGGGGCTGCT  
CCGTTCTCTGCCTGGCCTGAGGCTCCCTGAGCCGCCTCCCCACCATCACCATGGCCAAGGGCTTCTATAT  
TTCCAAGTCCCTGGGCATCCTGGGGATCCTCCTGGGCGTGGCAGCCGTGTGCACAATCATCGCACTGTCA  
GTGGTGTACTCCAGGAGAAGAACAAGAACGCCAACAGCTCCCCCGTGGCCTCCACCACCCCGTCCGCCT  
CAGCCACCACCAACCCCGCCTCGGCCACCACCTTGGACCAAAGTAAAGCGTGGAATCGTTACCGCCTCCC  
CAACACGCTGAAACCCGATTCTACCAGGTGACGCTGAGACCGTACCTCACCCCAATGACAGGGGCGCTG  
10 TACGTTTTTAAGGGCTCCAGCACCGTCCGTTTCACCTGCAAGGAGGCCACTGACGTCATCATCCACA  
GCAAGAAGCTCAACTACACCTCAGCCAGGGGCAAGGGTGGTCTGCGTGGTGTGGGAGGCTCCAGCC  
CCCCGACATTGACAAGACTGAGCTGGTGGAGCCACCGAGTACCTGGTGGTGCACCTCAAGGGCTCCCTG  
GTGAAGGACAGCCAGTATGAGATGGACAGCGAGTTTCGAGGGGGAGTTGGCAGATGACCTGGCGGGCTTCT  
ACCGCAGCGAGTACATGGAGGGCAATGTGAGAAAGTGGTGGCCACTACACAGATGCAGGCTGCAGATGC  
15 CCGGAAGTCTTCCCATGCTTCGATGAGCCGGCCATGAAGGCCGAGTTCAACATCACGCTTATCCACCCC  
AAGGACCTGACAGCCCTGTCCAACATGCTTCCCAAAGTCCCAGCACCCCACTTCCAGAAGACCCCAACT  
GGAATGTCACTGAGTTCCACACCACGCCCAAGATGTCCACGTACTTGCTGGCCTTCATTGTGCTGAGTT  
CGACTACGTGGAGAAGCAGGCATCCAATGGTGTCTTGATCCGGATCTGGGCCCGGGCCAGTGCCATTGCG  
GCGGGCCACGGCGATTATGCCCTGAACGTGACGGGCCCCATCCTTAACCTTCTTGTCTGGTCAATTATGACA  
20 CACCTTACCCTACCCAAATCAGACCAAGTGGCCTGCCAGACTTCAACGCCGGCGCCATGGAGAACTG  
GGGACTGGTGGACTACCGGGAGAATCCCTGCTGTTTCGACCCCTGTCTCCTCCAGCAGCAACAAGGAG  
CGGGTGGTCACTGTGATTGCTCATGAGCTGGCCACCAGTGGTTTCGGGAACCTGGTGACCATAGAGTGGT  
GGAATGACCTGTGGCTGAACGAGGGCTTCGCCCTCTACGTGGAGTACCTGGGTGCTGACTATGCGGAGCC  
CACCTGGAACCTTGAAGACCTCATGGTGTGATGATGTGTACCGCGTGTGTCAGTGGATGCACTGGCC  
25 TCCTCCACCCGCTGTCCACACCCGCCTCGGAGATCAACACGCCGGCCAGATCAGTGAGCTGTTTGACG  
CCATCTCCTACAGCAAGGGCGCCTCAGTCTCAGGATGCTCTCCAGCTTCTGTCCGAGGACGTATTCAA  
GCAGGGCTTGGCGTCTTACCTCCACACCTTTGCCCTACCAGAACACCATCTACCTGAACCTGTGGGACCAC  
CTGCAGGAGGCTGTGAACAACCGGTCCATCCAACCTCCCCACCACCGTGCAGGACATCATGAACCGCTGGA  
CCCTGCAGATGGGCTTCCCGGTATCATCGGTGGATACCAAGCAGGGGACCCCTTTCCAGGACACTTCTCT  
30 CTTTGACCCCGATTCCAATGTTACCCGCCCTCAGAACTCAACTACGTGTGGATTGTGCCCATCATCTCC  
ATCAGAGATGGCAGACAGCAGCAGGACTACTGGCTGATAGATGTAAGAGCCAGAACGATCTCTTCAGCA  
CATCAGGCAATGAGTGGGTCTGCTGAACCTCAATGTGACGGGCTATTACCGGGTGAACACGACGAAGA  
GAACTGGAGGAAGATTGAGACTCAGCTGCAGAGAGACCACTCGGCCATCCCTGTCAATCGGGCACAG  
ATCATTAAATGACGCCTTCAACCTGGCCAGTGCCCATAAAGTCCCTGTCACTCTGGCGCTGAACAACACCC  
35 TCTTCTTGATTGAAGAGAGACAGTACATGCCCTGGGAGGCCGCCCTGAGCAGCCTGAGCTACTTCAAGCT  
CATGTTTGACCGCTCCGAGGTCTATGGCCCCATGAAGAACTACCTGAAGAAGCAGGTACACCCCTCTTC  
ATTCACTTCAGAAATAATACCAACAACCTGGAGGGAGATCCCAGAAAACCTGATGGACCAGTACAGCGAGG  
TTAATGCCATCAGCACCGCCTGCTCCAACGGAGTTCCAGAGTGTGAGGAGATGGTCTCTGGCCTTTTCAA  
GCAGTGGATGGAGAACCCCAATAATAACCCGATCCACCCCAACCTGCGGTCCACCGTCTACTGCAACGCT  
40 ATCGCCAGGGCGGGGAGGAGGAGTGGGACTTCGCCCTGGGAGCAGTTCCGAAATGCCACACTGGTCAATG  
AGGCTGACAAGCTCCGGGCAGCCCTGGCCTGCAGCAAAGAGTTGTGGATCCTGAACAGGTACCTGAGCTA  
CACCTGAACCCGACTTAATCCGGAAGCAGGACGCCACCTCTACCATCATCAGCATTACCAACAACGTC  
ATTGGGCAAGGTCTGGTCTGGGACTTTGTCCAGAGCAACTGGAAGAAGCTTTTTAACGATTATGGTGGTG  
GCTCGTTCTCCTTCTCCAACCTCATCCAGGCAGTGACACGACGATTCTCCACCGAGTATGAGCTGCAGCA  
45 GCTGGAGCAGTTCAAGAAGGACAACGAGGAAAACAGGCTTCGGCTCAGGCACCCGGGCCCTGGAGCAAGCC  
CTGGAGAAGACGAAAGCCAACATCAAGTGGGTGAAGGAGAACAAGGAGGTGGTGTCTCAGTGGTTACAG  
AAAACAGCAAATAGTCCCCAGCCCTTGAAGTACCCGGCCCCGATGCAAGGTGCCCCACATGTGTCCATCC  
CAGCGCTGGTGCAGGGCCTCCATTCTGGAGCCCCGAGGACCAAGTGTCTCTCCCTCAAGGACAAAGTCT  
CCAGCCACGTTCTCTCTGCTGTGAGCCAGTCTAGTTCTGTATGACCCAGGCTGCCTGAGCACCTCCCA  
50 GCCCCTGCCCCCTCATGCCAACCCCGCCCTAGGCCTGGCATGGCACCTGTGCGCCAGTGCCCTGGGGCTGA  
TCTCAGGGAAGCCAGCTCCAGGGCCAGATGAGCAGAAGCTCTCGATGGACAATGAACGGCCTTGCTGGG  
GGCCGCCCTGTACCCTCTTTACCTTTCCCTAAAGACCCTAAATCTGAGGAATCAACAGGGCAGCAGATC  
TGATATATTTTTTTCTAAGAGAAAATGTAAATAAAGGATTTCTAGATGAAAAAAAAAAAAAAAAAAAA

55 >gi|4502095|ref|NP\_001141.1| membrane alanine aminopeptidase precursor;  
microsomal aminopeptidase; Alanyl (membrane) aminopeptidase  
(aminopeptidase N, aminopeptidase M, [Homo sapiens])  
MAKGFYISKSLGILGILLGVAAVCTIIALSVVYSQEKKNANSSPVASTTPSASATTNPASATTLQDSKA  
WNRRLPNTLKPDSYQVTLRPYLTPNDRGLYVFKGSSTVRFTCKEATDVIIHSHKKNLYTSLQGHRVRLR  
60 GVGGSQPPDIDKTELVEPTEYLVVHLKGS LVKDSQYEMDSEFEGELADDLAGFYRSEYMEGNVRKV VATT  
QMQAADARKSFPCFDEPAMKAEFNITLIHPKDLTALSNMLPKGPSTPLPEDPNWNVTEFHTTPKMSTYLL

AFIVSEFDYVEKQASNGVLIRIWARPSAIAAGHGDIYALNVTGPILNFFAGHYDTPYPLPKSDQIGLPDFN  
 AGAMENWGLVTYRENSLLFDPLSSSSSNKERVVTVIAHELAHQWFGNLVTIEWWNDLWLNIEGFASYVEYL  
 GADYAEPTWNKDLMLVNDVYRVMVDALASSHPLSTPASEINTPAQISELFDASISYKASVLRMLSSF  
 LSEDFVKQGLASYLHTFAYQNTIYLNLDHLEAVNNRSIQLPTTVRDIMNRWTLQMGFPVITVDTSTGT  
 5 LSQEHFLLDPSNVTRPSEFNIVVIVPITSIRDGRQQQDYWLIDVRAQNDLFSTSGNEWVLLNLNVTGY  
 RVNYDEENWRKIQTQLQRDHSAPVINRAQIINDAFNLASAHKVPVTLALNNTLFLIEERQYMPWEAALS  
 SLSYFKLMFDRSEVYGPMKNYLKKQVTPLFIFHRNNTNNWREIPENLMDQYSEVNAISTACSNVPECEE  
 MVSGLFKQWMENPNNNPIHPNLRSTVYCNAIAQGGEEEDFAWEQFRNATLVNEADKLRAALACSKELWI  
 10 LNRYLSYTLNPD LIRKQDATSTIISITNNVIGQGLVWDFVQSNWKKLFNDYGGGSFSSFNLIQAVTRRFS  
 TEYELQQLEQFKDNEETGFGSGTRALEQALEKTKANIKWVKENKEVLQWFTENSK

PRK-1

15 GH1-54-PCR-G3F1  
 TCCTTTCCCGCCACGCACTACAGCACCCCTGTTGCAAGCCCGCGCCGCTCACAGGGACCCCTGAGGTACGAGTG  
 GTGGGCTGCAGAGACCTCCCAGAGACCATCCCGTGGAACCCTACCCCTCAATGGGGGGACCTGGGACCCCCA  
 GACAGCGCCCCCTTCCCTGAGCCGCCCAGCCCGGGGCCGAGTAACCCAGCACAGTGGTTAGATAGATAAAG  
 CGGCCGCTCGACTAGTCTGAGGTCTGATACTCACTGACGTGATACGT  
 20 >gi|4506072|ref|NM\_002741.1| Homo sapiens protein kinase C-like 1  
 (PRKCL1), mRNA  
 TGAGTAAATCGATACATCATACGCGCGCTCCTCTGGCCGCCCCCTCCCTCCGACGATCGGGGACCCCTGGCG  
 GCGGCAGGAGGACATGGCCAGCGACGCCGTGCAGAGTGAGCCTCGCAGCTGGTCCCTGCTAGAGCAGCT  
 25 GGGCCTGGCCGGGGCAGACCTGGCGGCCCCCGGGGTACAGCAGCAGCTGGAGCTGGAGCGGGAGCGGCTG  
 CGGCGGGAAATCCGCAAGGAGCTGAAGCTGAAGGAGGGTGCTGAGAACCTGCGGCGGGCCACCACTGACC  
 TGGGCCGACGCTGGGCCCCGTAGAGCTGCTGCTGCGGGGCTCCTCGCGCCGCTCGACCTGCTGCACCA  
 GCAGCTGCAGGAGCTGCACGCCCCACGTGGTGCTTCCCGACCCGGCGGCCACCCACGATGGCCCCCAGTCC  
 CCTGGTGGGGTGGCCCCACCTGCTCGGCCACCAACCTGAGCCGCGTGGCGGGCCTGGAGAAGCAGTTGG  
 30 CCATTGAGCTGAAGGTGAAGCAGGGGGCGGAGAACATGATCCAGACCTACAGCAATGGCAGCACCAAGGA  
 CCGGAAGCTGCTGCTGACAGCCAGCAGATGTTGCAGGACAGTAAGACCAAGATTGACATCATCCGCATG  
 CAACTCCGCCGGGCGCTGCAGGCCGACCAGCTGGAGAACCAGGCAGCCCCGGATGACACCCAAGGGAGTC  
 CTGACCTGGGGGCTGTGGAGCTGCGCATCGAAGAGCTGCGGCACCACTTCCGAGTGGAGCACGCGGTGGC  
 CGAGGGTGCCAAGAACGTACTGCGCCTGCTCAGCGCTGCCAAGGCCCGGACCGCAAGGCAGTCAGCGAG  
 35 GCCCAGGAGAAATTGACAGAATCCAACCAGAAGCTGGGGCTGCTGCGGGAGGCTCTGGAGCGGAGACTTG  
 GGGAGCTGCCCCGCCGACCACCCCAAGGGGCGGCTGCTGCGAGAAGAGCTCGCTGCGGCCTCCTCCGCTGC  
 CTTACGACCCCGCTGGCCGGGCCCCCTTCCCGCCACGCACCTACAGCACCTGTGCAAGCCCCGCGCGCTC  
 ACAGGGACCTTGGAGGTACGAGTGGTGGGCTGCAGAGACCTCCAGAGACCATCCCGTGGAACCCATCCC  
 CCTCAATGGGGGACCTGGGACCCAGACCCGCGCCCCCTTCCCTGAGCCGCCAGCCCCGGGGCCTTTTA  
 40 CAGCCGAAGCGGAAGCCTCAGTGGCCGGAGCAGCCTCAAAGCAGAAGCCGAGAACACCAGTGAAGTCAGC  
 ACTGTGCTTAAGCTGGATAACACAGTGGTGGGGCAGACGTCTTGAAGCCATGTGGCCCCAATGCCTGGG  
 ACCAGAGCTTCACTCTGGAGCTGGAAAGGGCACGGGAACTGGAGTTGGCTGTGTTCTGGCGGGACACGCG  
 GGGCCTGTGTGCCCTCAAATTCCTGAAGTTGGAGGATTTCTTGGACAATGAGAGGCATGAGGTGCAGCTG  
 GACATGGAACCCAGGGCTGCCTGGTGGCTGAGGTACCTTCCGCAACCCTGTCAATTGAGAGGATTCTCTC  
 45 GGCTCCGACGGCAGAAGAAAATTTCTCCAAGCAGCAAGGGAAGGCGTTCCAGCGTGCTAGGCAGATGAA  
 CATCGATGTCGCCACGTGGGTGCGGCTGCTCCGGAGGCTCATCCCCAATGCCACGGGCACAGGCACCTTT  
 AGCCCTGGGGCTTCTCCAGGATCCGAGGCCCGGACCCAGGGTGACATATCGGTGGAGAAGCTGAACCTCG  
 GCACTGACTCGGACAGCTCACCTCAGAAGAGCTCGCGGGATCCTCCTTCCAGCCCATCGAGCCTGAGCTC  
 CCCCATCCAGGAATCCACTGCTCCCGAGCTGCCTTCGGAGACCCAGGAGACCCAGGCCCGCCCTGTGC  
 50 AGCCCTCTGAGGAAGTCACCTCTGACCCTCGAAGATTTCAAGTTCTGGCGGTGCTGGGCGGGGTCATT  
 TTGGGAAGGTGCTCCTCTCCGAATTCGGGCCAGTGGGGAGCTGTTCCGCATCAAGGCTCTGAAGAAAGG  
 GGACATTGTGGCCCCGAGACGAGGTGGAGAGCCTGATGTGTGAGAAGCGGATATTGGCGGCAGTGACCACT  
 GCGGGACACCCCTTCTGGTGAACCTCTTCGGCTGTTTCCAGACACCGGAGCACGTGTGCTTCGTGATGG  
 AGTACTCGGCCGGTGGGGACCTGATGCTGCACATCCACAGCGACGTGTTCTCTGAGCCCCGTGCCATCTT  
 55 TTATTCGCCCTGCGTGGTGCTGGGCCTACAGTTTCTTACGAACACAAGATCGTCTACAGGGACCTGAAG  
 TTGGACAATTTGCTCCTGACACCGAGGGCTACGTCAGATCGCAGACTTTGGCCTCTAGGAAGGAGGA  
 TGGGCTATGGGGACCGGACCATCTGTGTGGGACCCCGGAGTTCTTGGCCCCCTGAGGTGCTGACCGGA  
 CACGTGCTACACGCGAGCTGTGGACTGGTGGGGACTGGGTGTGCTGCTCTACGAGATGCTGGTTGGCGAG  
 TCCCCATTCCAGGGGATGATGAGGAGGAGGTCTTCGACAGCATCGTCAACGACGAGGTTTCGTACCCCC  
 60 GCTTCTGTGCGGCCGAAGCCATCGGCATCATGAGAAGGCTGCTTCGGAGGAACCCAGAGCGGAGGCTGGG  
 ATCTAGCGAGAGAGATGCAGAAGATGTGAAGAAACAGCCCTTCTTCAGGACTCTGGGCTGGGAAGCCCTG

TTGGCCCCGGCGCCTGCCACCGCCCTTTGTGCCACGCTGTCCGGCCGCACCGACGTCAGCAACTTCGACG  
 AGGAGTTCACCGGGGAGGCCCCCACACTGAGCCCCGCCCGCGACGCGCGGCCCTCACAGCCGCGGAGCA  
 GGCAGCCTTCCTGGACTTCGACTTCGTGGCCGGGGGCTGCTAGCCCCCTCCCCTGCCCCCTGCCCCCTGCC  
 CTGCCCGAGAGCTCTTAGTTTTTAAAAAGGCCCTTTGGGATTTGCCGGAIAAAAAAAAAAAAAA

5

>gi|4506073|ref|NP\_002732.1| protein kinase C-like 1; serine-threonine  
 kinase N [Homo sapiens]

MASDAVQSEPRSWSLLEQLGLAGADLAAPGVQQQLELERERLRREIRKELKLKEGAENLRATTDLGRSL  
 GPVELLLRGSSRRLLDLHQQLQELHAHVLPDPAATHDGPQSPGAGGPTCSATNLSRVAGLEKQLAIELK  
 10 VKQGAENMIQTYNSGSTKDRKLLLTAAQMLQDSKTKIDIIRMQLRRALQADQLENQAAPDDTQGSPLGA  
 VELRIEELRHFRVEHAVAEGAKNVLRLLSAAKAPDRKAVSEAQEKLTESNQKLGLLREALERRLGELPA  
 DHPKGRLLREELAAASSAAFSTRLAGPFPATHYSTLCKPAPLTGTLEVRVVGCRDLPETIPWNPTPSMGG  
 PGTPDSRPPFLSRPARGLYSRGSLSGRSSLKAEAEENTSEVSTVLKLDNTVVVGQTSWKPCGPNAWDQSFT  
 LELERARELELAVFWRDQRLCALFKLEDFLDNERHEVQLDMEPQGCCLVAEVTFRNPVIERIPRLRRQ  
 15 KKIFSKQQGKAFQRRARQMNIDVATWVRLRLRRLIPNATGTGTFSPGASPGSEARTTGDISVEKLNLTGDS  
 SSPQKSSRDPPSSPSSLSPIQESTAPELPSETQETPGPALCSPLRKSPLTLEDFKFLAVLGRGHFGKVL  
 LSEFRPSGELFAIKALKKGDIVARDEVESLMCEKRILAAVTSAGHPFLVNLFGCFQTPHEVCFVMEYSAG  
 GDLMLHIHSDVFSEPRAFYFYSACVVLGLQFLHEHKIVYRDLKLDNLLDTEGYVKIADFGLCKEGMGYGD  
 RTSTFCGTPEFLAPEVLTDTSYTRAVDWGLGVLLYEMLVGESPFPGDDEEEVFDSIVNDEVRYRPRFLSA  
 20 EAIGIMRRLRLRNPERRLGSSERDAEDVKKQPFRTLGEALLARRLPPFPVPTLSGRTDVSNFDEEFTG  
 EAPTLSPPRDARPLTAAEQAAFLDFDFVAGGC

Zip kinase

25

GH1-68-PCR-G3F1

GCTGGGTTTCATTTTCGAGTATTCGCGGGCCTGCTCTCAGCTAGGTTTAGCCCGTTCTGTGACCCCTCCACGTGC  
 ACTCGTGGTCACTGTGGCACCGTGAGGGTTGGGACCCACCGAGGCGCAAGGCGGCCGAATGCGCCTGTTTCA  
 CCCGAGAGGTTTTCGCGGTAGTTGCCGGACATTCGCGGGGTGCTGCCTGTTGTCTGCCATTATGCCCAGGAG  
 GAGGTGAGTGGGGAGGGTGGGATGGACGGCGGACGAGCAGTCCCCACGCTGCTTGGTGGCGCGCGCTT  
 30 GGTGGGGTCTTCCACTGTGTGCCCTTCTCGCCGAGGCGGCTCCCCCGCGTGTGGGGTGCCTTCCGCGAC  
 TCCTCCGCACGCGAGAAACCAGCACAGTGGTTAGAGTAGATAAAGCGGGCGAGTCGACTAGATCTGAGGTCT  
 GATACTCACTGACTGTTTCGTAA

>gi|4557510|ref|NM\_001348.1| Homo sapiens death-associated protein  
 kinase 3 (DAPK3), mRNA

GTGCGCATTAGGGGACTCCTGAGGTCCTATCTCCAGGCTGCGGTGACTGCACTTTCCCTGGAGTGGAAGC  
 TGCTGGAAGGCGGACCGGCCGCCATGTCCACGTTCAAGCAGGAGGACGTGGAGGACCATATGAGATGGG  
 GGAGGAGCTGGGCAGCGGCCAGTTTTCGATCGTGCAGGAGTCCCGCAGAGGCGCAAGGCGCAAGGAGTAC  
 GCAGCCCAAGTTTCATCAAGAAGCGCCGCTGTCTATCCAGCCGCGTGGGGTGAGCCGGGAGGAGATCGAGC  
 40 GGGAGGTGAACATCCTGCGGGAGATCCGGCACCCCAACATCATCACCTGCACGACATCTTCGAGAACA  
 GACGGACGTGGTCTCTATCTGAGCTGGTCTCTGCGGGGAGCTCTTTGACTTCCTGGCGGAGAAAGAG  
 TCGCTGACGGAGGACGAGGCCACCCAGTTTCTCAAGCAGATCCTGGACGGCGTTCACTACCTGCACTCTA  
 AGCGCATCGCACACTTTGACCTGAAGCCGGAACATCATGCTGCTGGACAAGAACGTGCCCCAACCCACG  
 AATCAAGCTCATCGACTTCGGCATCGCGCACAAGATCGAGGCGGGGAACGAGTTCAAGAACATCTTCGGC  
 45 ACCCCGGAGTTTGTGGCCCCAGAGATTGTGAACTATGAGCCGCTGGGCCTGGAGGCGGACATGTGGAGCA  
 TCGGTGTCATCACCTATATCCTCCTGAGCGGTGCATCCCCGTTCTTGGGCGAGACCAAGCAGGAGACGCT  
 CACCAACATCTCAGCGGTGAACACGAGTTCGACGAGGAGTACTTCAGCAACACACGAGCTGGCCAAG  
 GACTTCATTCGCGCGCTGCTCGTCAAAGATCCCAAGCGGAGAATGACCATTGCCAGAGCCTGGAACATT  
 CCTGGATTAAAGGCGATCCGGCGGCGGAACGTGCGTGGTGAGGACAGCGGCCGCAAGCCCCGAGCGGCGGCG  
 50 CCTGAAGACCACGCGTCTGAAGGAGTACACCATCAAGTCGCACTCCAGCTTGCCGCCCAACAACAGCTAC  
 GCCGACTTCGAGCGCTTCTCAAGGTGCTGGAGGAGGCGGCGGCCGAGGAGGGCCTGCGCGAGCTGC  
 AGCGCAGCCGGCGGCTCTGCCACGAGGACGTGGAGGCGTGGCCGCCATCTACGAGGAGAAGGAGGCTTG  
 GTACCGCGAGGAGAGCGACAGCCTGGGCCAGGACCTGCGGAGGCTACGGCAGGAGCTGCTCAAGACCGAG  
 GCGCTCAAGCGGCAGGCGCAGGAGGAGGCCAAGGGCGCGCTGCTGGGGACCAGCGGCCTCAAGCGCCGCT  
 55 TCAGCCGCCCTGGAGAACCCTACGAGGCGCTGGCCAAGCAAGTAGCCTCCGAGATGCGCTTCGTGACGGA  
 CCTCGTGCAGCCCTGGAGCAGGAGAAGCTGCGAGGCGTGGAGTGGGGCTGCGCTAGGCGCAGCTGGGGT  
 GGGCCAGGCCCGGAGCAGCGGAGCTCGGCTGCGGTGGGGGCGCTTCCTGTGGACGCTGCGCCTGCCA  
 TCGCCCGGGTGCTGTCTTGGCCAGCGCCACCAGGCTGGAGGCGGAGTGGGAGGAGCTGGAGCCAGGCC  
 CGTAAGTTCGAGGAGGAGGGTGGGTGTGGGACGGGGCTGCTTCTCTACACAGCCTCTACGCTGGCCTTCA  
 60 CCTTACCCCTGCATCGTGGTGACCTGGGACCTCCAGGACGCTGGCCTGTGGCACCGTGAGGGTTG  
 GGACCCACCGAGGCGCAGAGGCGGCCCAATGCAGCCCTGGTTCAGGCCCGAGGAGGGTTTTCGGGTAG

TTGCACGGACAATTTCGGCGGGGTGCTGCCTGTTGCTGCCATTAGCCCAGGAGGAGGTCGTGGGACGGGGA  
GGGTGGGATGGACGGCGGACAGGCAGTCCCCACGCTGCTGGGTGGCGCCGGGCTTGGTGGGGTCTTCCAC  
TGTGTGCCCTTCTCGCCGAGGCCGGTCCCCCGGGTGTGGGGTGGCCTGCTGCGGACTCCTCCGCGAGCCC  
CATCGTCGCGCCTGTGGACGCCTAGGCAAGAGCGGCCCTCTGCAGCCAAGAGAAATAAAATACTGGCTTC  
CAGAT

5

>gi|4557511|ref|NP\_001339.1| death-associated protein kinase 3 [Homo sapiens]  
MSTFRQEDVEDHYEMGEELGSGQFAIVRKCRQKGTGKEYAAKFIKKRRLSSRRRGVSREEIEREVNIRE  
IRHPNIITLHDIFENKTDVVLILELVSGGELFDLAEKESLTEDEATQFLKQILDGVHYLHRSKRIAHFDL  
KPENIMLLDKNVPNPRIKLIDFGIAHKIEAGNEFKNIFGTPEFVAPEIVNYEPLGLEADMWSIGVITYIL  
LSGASPFLETGKETLTNISAVNYDFDEEYFSNTSELAKDFIRLLVKDPKRRMTIAQSLEHSWIKAIRR  
RNVRGEDSGRKPERRLKTTRLKEYTIKSHSLPPNNSYADFERFSKVLEEAEEGLRELQSRRLCH  
EDVEALAAIYEEKEAWYREESDSLQDLRRLRQELLKTEALKRQAQEEAKGALLGTSGLKRRFSRLNRY  
EALAKQVASEMRFVQDLVRALEQEKLGVECGLR

10

15

>gi|2911155|dbj|AB007144.1| Homo sapiens mRNA for ZIP-kinase, complete cds

20

25

30

35

40

45

50

>gi|2911156|dbj|BAA24955.1| ZIP-kinase [Homo sapiens]  
MSTFRQEDVEDHYEMGEELGSGQFAIVRKCRQKGTGKEYAAKFIKKRRLSSRRRGVSREEIEREVNIRE  
IRHPNIITLHDIFENKTDVVLILELVSGGELFDLAEKESLTEDEATQFLKQILDGVHYLHRSKRIAHFDL  
KPENIMLLDKNVPNPRIKLIDFGIAHKIEAGNEFKNIFGTPEFVAPEIVNYEPLGLEADMWSIGVITYIL  
LSGASPFLETGKETLTNISAVNYDFDEEYFSNTSELAKDFIRLLVKDPKRRMTIAQSLEHSWIKAIRR  
RNVRGEDSGRKPERRLKTTRLKEYTIKSHSLPPNNSYADFERFSKVLEEAEEGLRELQSRRLCH  
EDVEALAAIYEEKEAWYREESDSLQDLRRLRQELLKTEALKRQAQEEAKGALLGTSGLKRRFSRLNRY  
EALAKQVASEMRFVQDLVRALEQEKLGVECGLR

55

60

>gi|5162883|dbj|AB022341.1| Homo sapiens mRNA for ZIP kinase, complete cds

GCACTTTCCCTGGAGTGGAAAGCTGCTGGAAGGCGGACCGGCCCATGTCCACGTTTCAGGCAGGAGGACG  
 TGGAGGACCATTATGAGATGGGGGAGGAGCTGGGCAGCGGCCAGTTTGCATCGTGCAGGAAAGTGCCGGCA  
 GAAGGGCACGGGCAAGGAGTACGCAGCCAAGTTTCATCAAGAAGCGCCGCTGTCTATCCAGCCGGCGTGGG  
 GTGAGCCGGGAGGAGATCGAGCGGGAGGTGAACATCCTGCGGGAGATCCGGCACCCCAACATCATCACCC  
 5 TGCACGACATCTTCGAGAACAAGACGGACGTGGTCTCATCTGGAGCTGGTCTCTGGCGGGGAGCTCTT  
 TGACTTCCTGGCGGAGAAGGAGTGCCTGACGGAGGACGAGGCCACCCAGTTCTCAAGCAGATCCTGGAC  
 GGCCTTCACTACCTGCACTCTAAGCGCATCGCACACTTTTGACCTGAAGCCGGAAAACATCATGCTGCTGG  
 ACAAGAAGCTGCCCAACCCACGAATCAAGCTCATCGACTTCGGCATCGCGCACAAAGATCGAGGCGGGGAA  
 CGAGTTCAAGAACATCTTCGGCACCCCGGAGTTTGTGGCCCCAGAGATTGTGAACATATGAGCCGCTGGGC  
 10 CTGGAGGCGGACATGTGGAGCATCGGTGTCTATCACCTATATCCTCCTGAGCGGTGCATCCCCGTTCCTGG  
 GCGAGACCAAGCAGGAGACGCTCACCAACATCTCAGCCGTGAACACGACTTCGACGAGGAGTACTTCAG  
 CAACACCAGCGAGCTGGCCAAGGACTTCATTTCGCCGGCTGCTCGTCAAAGATCCCAAGCGGAGAATGACC  
 ATTGCCCAGAGCCTGGAACATTCTGGATTAAGGCGATCCGGCGGCGGAACGTGCGTGGTGAGGACAGCG  
 GCCGCAAGCCCGAGCGGCGGCGCTGAAGACCACGCGTCTGAAGGAGTACACCATCAAGTCGCACTCCAG  
 15 CTTGCCGCCCAACAACAGCTACGCCGACTTCGAGCGCTTCTCCAAGGTGCTGGAGGAGGCGGCGGCC  
 GAGGAGGGCCTGCGCGAGCTGCAGCGCAGCCGGCGGCTCTGCCACGAGGACGTGGAGGCGCTGGCCGCCA  
 TCTACGAGGAGAAGGAGGCTGGTACCGCGAGGAGAGCGACAGCCTGGGCCAGGACCTGCGGAGGCTACG  
 GCAGGAGCTGCTCAAGACCGAGGCGCTCAAGCGGCAGGCGCAGGAGGAGGCCAAGGGCGCGCTGCTGGGG  
 ACCAGCGGCCTCAAGCGCCGCTTCAGCCGCTTGAGAACCGCTACGAGGCGCTGGCCCAAGCAAGTAGCCT  
 20 CCGAGATGCGCTTCGTGCAGGACCTCGTGCGCGCCCTGGAGCAGGAGAAGCTGCAGGCGCTGGAGTGCGG  
 GCTGCGCTAGGCGCAGTGGGGTGGGCCAGGCCCGGACAGCCGGAGCTCGGCCTGCGGTGGGGGCGCTT  
 CTTGTGGACGCTGCGCCTCCCATCGCCCGGCTGCTGTCTTGGCCAGCGCCACCAGGCTGGAGGCGGAG  
 TGGGAGGAGCTGGAGCCAGGCCCGTAAGTTCGACGGCAGGGGTGGGTGTGGGACGGGGCTGCTTCTCTAC  
 ACATCCTCCACGCTGGCCTTCACCTTCACCCCTGCATCGTGGTGACCCTGGGACCCTCCAGGCAGCGTG  
 25 GCCTGTGGCACCGTGAGGGTTGGGACCCACCGAGGCGCAGAGGCGGCCCAATGCAGCCCTGGTTACGGC  
 CCGGAGGAGGGTTTGGCGGTAGTTGCACGGACAATTGGCGGGGTGCTGCCTGTTGCTGCCATTAGCCCA  
 GGAGGAGGTGCTGGGACGGGGAGGGTGGGATGGACGGCGGACAGGCAGTCCCCACGCTGCTGGGTGGCGC  
 CGGGCTTGGTGGGGTCTTCCACTGTGTGCCCTTCTCGCCGAGGCCGGTCCCCGGGTGTGGGGTGCCTG  
 CTGCGGACTCCTCCGCGAGCCCCATCGTCGCGCTGTGGACGCCTAGGCAAGAGCGGCCCTCTGCAGCCA  
 30 AGAGAAATAAAATACTGGCTTCCAG

>gi|5162884|dbj|BAA81746.1| ZIP kinase [Homo sapiens]  
 MSTFRQEDVEDHYEMGEELGSGQFAIVRKCRQKGTGKEYAAKFIKKRRLSSSRRGVSREEIEREVNIRE  
 35 IRHPNIIITLHDIFENKTDVVLILELVSGGELFDFLAEKESLDEATQFLKQILDGVHYLHSHKRIAHFDL  
 KPENIMLLDKNPNPRIKLIDFGIAHKIEAGNEFKNIFGTPEFVAPEIVNYEPLGLEADMWSIGVITYIL  
 LSGASPFLETGKETLTNISAVNYDFDEEYFSNTSELAKDFIRLLVKDPKRRMTIAQSLEHSHWIKAIRR  
 RNVRGEDSGRKPERRRLKTTTLKEYTIKSHSLPPNNSYADFERFSKVL EAAAAEEGLRELQRSRLCH  
 EDVEALAAIYEEKEAWYREESDSLQDLRLRQLLELKEALKRQAQEEAKGALLGTSGLKRRFRSRLNRY  
 40 EALAKQVASEMRVQDLVRALEQEKLQGVCEGLR

Gas6

GH1-50-PCR-G3F1  
 GCGCAGGAATCTGGTCATCAAGGTCAACAGGGATGCTGTCATGAAAATCGCGGTGGCCGGGGACTTGTTC  
 45 ACCGGAGCGAGGACTGTATCATCTGAACCTTACCGTGGGAGGTATTCCCTTCCATGAGAAGGACTACGTGCA  
 GCTATAAAACCTCGTCTGGATGCTGCACTGAAGAGCGCGCAGAAACCAACAC

>gi|4557616|ref|NM\_000820.1| Homo sapiens growth arrest-specific 6  
 (GAS6), mRNA  
 50 CCGCAGCCGCCCGCCGCCGCCGCCGCGATGTGACCTTCAGGGCCGCCAGGACGGGATGACCGGAGCCT  
 CCGCCCCGCGGCGCCCGCTCGCCTCGGCCTCCCGGGCGCTCTGACCGCGCTCCCCGGCCCCCATGGCC  
 CCTTCGCTCTCGCCCGGGCCCGCCCGCTGCGCCGCGCGCCGAGCTGCTGCTGCTGCTGCTGGCCGCGG  
 AGTGCGCGCTTGCCGCGCTGTTGCCGGCGCGCGAGGCCACGCAGTTCTGCGGCCAGGCAGCGCCGCGC  
 CTTTCAGGTCTTCGAGGAGGCCAAGCAGGGCCACCTGGAGAGGGAGTGCGTGGAGGAGCTGTGCAGCCGC  
 55 GAGGAGGCGCGGAGGTGTTTCGAGAACGACCCGAGACGGATTATTTTACCCAAGATACTTAGACTGCA  
 TCAACAAGTATGGGTCTCCGTACACCAAAAACCTCAGGCTTCGCCACCTGCGTGCAAAACCTGCCTGACCA  
 GTGCACGCCCAACCCCTGCGATAGGAAGGGGACCCAAAGCCTGCCAGGACCTCATGGGCAACTTCTTCTGC  
 CTGTGTAAAGCTGGCTGGGGGGGCCGGCTCTGCGACAAAGATGTCAACGAATGCAGCCAGGAGAACGGGG  
 GCTGCCCTCCAGATCTGCCACAACAAGCCGGGTAGCTTCCACTGTTCTGCCACAGCGGCTTCGAGCTCTC  
 60 CTCTGATGGCAGGACCTGCCAAGACATAGACGAGTGCGCAGACTCGGAGGCCTGCGGGGAGGCGCGCTGC  
 AAGAACCTGCCCGGCTCCTACTCCTGCCTCTGTGACGAGGGCTTTGCGTACAGCTCCAGGAGAAGGCTT

GCCGAGATGTGGACGAGTGTCTGCAGGGCCGCTGTGAGCAGGTCTGCGTGAACCTCCCCAGGGAGCTACAC  
 CTGCCACTGTGACGGGCGTGGGGGCTCAAGCTGTCCAGGACATGGACACCTGTGAGGACATCTTGCCG  
 TCGTGGCCTTACGCGTGGCCAAGAGTGTGAAGTCTTGTACCTGGGCCGGATGTTTCACTGGGACCCCCG  
 5 TGATCCGACTGCGCTTCAAGAGGCTGCAGCCACCAGGCTGGTAGCTGAGTTTGAATCTCCGACCTTTGA  
 CCCCAGGGCATCTCTCTTTGCGGAGGCCACCAGGACAGCACCTGGATCGTGTCTGGCCCTGAGAGCC  
 GGCCGGCTGGAGCTGCAGCTGCGCTACAACGGTGTGCGCCGTGTACCAGCAGCGGCCCGGTCAACAAC  
 ATGGCATGTGGCAGACAATCTCTGTTGAGGAGCTGGCGCGGAATCTGGTCATCAAGGTCAACAGGGATGC  
 TGTATGAAAATCGCGGTGGCCGGGACTTGTTCACCGGAGCGAGGACTGTATCATCTGAACCTGACC  
 10 GTGGGAGGTATTCCTTCCATGAGAAGGACCTCGTGCAGCCTATAAACCTCGTCTGGATGGCTGCATGA  
 GGAGCTGGAACCTGGCTGAACGGAGAAGACACCACCATCCAGGAAACGGTGAAAGTGAACACGAGGATGCA  
 GTGCTTCTCGGTGACGGAGAGAGGCTCTTTCTACCCGGGAGCGGCTTCGCCTTCTACAGCCTGGACTAC  
 ATGCGGACCCCTCTGGACGTGCGGACTGAATCAACCTGGGAAGTAGAAGTCGTGGCTCACATCCGCCAG  
 CCGCAGACACAGGCGTGTGTTGCGCTCTGGGCCCCGACCTCCGTGCGTGCCTCTCTCTGTGGCACT  
 GGTAGACTATCACTCCACGAAGAACTCAAGAAGCAGCTGGTGGTCTGGCCGTGGAGCATAACGGCCTTG  
 15 GCCCTAATGGAGATCAAGGTCTGCGACGGCCAAGAGCACGTGGTCACCGTCTCGCTGAGGGACGGTGAGG  
 CCACCCTGGAGGTGGACGGCACCAGGGGCCAGAGCGAGGTGAGCGCCGCGCAGCTGCAGGAGAGGCTGGC  
 CGTGCTCGAGAGGCACCTGCGGAGCCCCGTGCTCACCTTTGCTGGCGGCCTGCCAGATGTGCCGTGACT  
 TCAGCGCCAGTCACCGCGTTCTACCGCGGTGCATGACACTGGAGGTCAACCGGAGGCTGCTGGACCTGG  
 ACGAGGCGGCGTACAAGCACAGCGACATCACGGCCACTCCTGCCCCCGTGGAGCCCCCGCGAGCCTA  
 20 GGCCCCACGGGACGCGGAGGCTTCTCAGTCTCTGTCAGACAGCGGAGGAGCCTGGGGGCTCCTC  
 ACCACGTGGGGCATGCTGAGAGCTGGGCTTTCCTCTGTGACCATCCCGGCCTGTAACATATCTGTAAAT  
 AGTGAGATGGACTTGGGGCTCTGACCGCGCACTCAGCCGTGGGCGCGGGGAGGCGGGGAGGCGGCGCA  
 GCGCAGAGCGGGCTCGAAGAAAATAATTCTCTATTATTTTATTACCAAGCGCTTCTTTCTGACTCTAAA  
 ATATGGAAT  
 25  
 >gi|4557617|ref|NP\_000811.1| growth arrest-specific 6; AXL stimulatory  
 factor [Homo sapiens]  
 MAPSLSPGPAALRRAPQLLLLLLLAAECALAAALLPAREATQFLRPRQRRAFQVFEEAKQGHLERECVEELC  
 30 SREEAREVFENDPETDYFYPRYLDCKINKYGSPTKNSGFATCVQNLDPQCTPNPCDRKGTQACQDLMGNF  
 FCLCKAGWGGRLCDKDVNECSQENGGCLQICHNKPGSFHCSCHSGFELSSDGRTCQDIDECADSEACGEA  
 RCKNLPGSYSCLDEGFAYSSQEKACRDVDECLQGRCEQVCVNSPGSYTCHCDGRGGLKLSQDMDTCEDI  
 LPCVPFVAKSVKSLYLGRMFSGTPVIRLRFKRLQPTRLVAEFDFRTFDPEGILLFAGGHQDSTWIVLAL  
 RAGRLELQLRYNGVGRVTSSGPVINHGMMQWQTSVEELARNLVIKVNRAVMKIAVAGDLFQPERGLYHLN  
 LTVGGIIPFHEKDLVQPINPRLDGCMRSWNWLNGETTTIQTETVKVNTRMQCFVTERGSFYPGSGFAFYSL  
 35 DYMRTPLDVGTSTWEVEVVAHIRPAADTGVLFWAPDLRAVPLSVALVDYHSTKKLKKQLVVLAVEHT  
 ALALMEIKVCDGQEHVVTVSLRDGEATLEVDGTRGQSEVSAAQLQERLAVLERHLRSPVLTFAAGGLPDPV  
 VTSAPVTAFYRGCMTELVNRLLDLDEAAYKHSIDTAHSCPPVEPAAA  
 40  
 SRm160  
 GH1-67-PCR-G3F1  
 GGTTCCTCTCCTCGTCTTCTCCTCCCCCTTCCCCTTCTAAGCCTGGCCCTCAGGCCTTGCCCAAACCTGCAAG  
 CCCCAGAAGCCACCCCTTGCAGCGGAGGTCCGCAGCCCCGGAAGCCAATAACTCCCTCAGGGGACTCTC  
 45 GGTCCCTCAGCTACTCGCCTGTGGAGCGTCGCCGTCCCTCGCCCCAGCCCTCACCACGGGACCAGCAGAGCA  
 GCAGCAGTGAGCGGGGTTCGCGGAGAGGCCAGCGTGGGGACAGCCGTCCCCAGCCACAAGCGCAGAGGAG  
 ACACCTAGCCCTCGGCCATGAGACACCGCTCCTCCAGGTCTCCATAAATTGTCTTTGGGGGATTNCAACACA  
 CCCATGCTCTTGAGCCACAAGGAGTGTCTCTTCTCCCCAGCAGAACCCTGGAAGGTCTTGTCTGCGTCT  
 CCTTTTAACCTTNGCAGCCTTTGATTGGAGGGCGTCCCCTTTTCTCCCCCTTTTTTTAG  
 50  
 >gi|19923465|ref|NM\_016333.2| Homo sapiens serine/arginine repetitive  
 matrix 2 (SRRM2), mRNA  
 GCGGCCAGGCGGGGTGCGAGTGGCGCAGTCGGAGCCCGTTGCGGCCCTGAGGAAGCGAGGAGGCGTCG  
 GCGTCGGCTGAGGCGGGCGGACCGGCGAGGCGAGGCGGCGGCCAGGCCCGAGGGACTCGGGAGCTCGA  
 55 GCAGCGGCGGCGGAAGACCTCTCCCCCTCGGAGGCGGCGGGCGGAGGCGGCGGGAGCGGTGGTGCCCCC  
 CCCGGGCACGGGGCCATGTACAACGGGATCGGGCTGCCGACGCCCCGGGGCAGCGGACCAACGGCTACG  
 TCCAGCGCAACCTGTCCCTGGTGGGGGCGCGGGGTGAGCGGCTGACTACAAGGAGAGGAGGAAT  
 GCGGCGCCTGGAGCTGCCCTGGTGAAGCGGCTAATCCTGACATCCTGGACCACGAGGAGCAAGCGCGC  
 GTCGAGCTGCGATGCCTCGAGCTGGAGGAGATGATGGAAGAGCAGGGGTACGAGGAACAGCAAATTCAGG  
 AAAAAGTGGCGACCTTTGACTCATGTTGCTGGAGAAGGATGTGAACCTGGGGGCAAGGAGGAGACCCC  
 60 AGGGCAGAGGCCAGCGGTACGGGAGACTCACCAGTTGGCAGAATTAAATGAGAAGAAGATGAAAGACTC  
 CGTGCTGCCTTTGGCATCAGTGATTCTTACGTAGATGGCAGCTCTTTTGATCCTCAGCGTCTGCCCCAG

AAGCTAAACAACCAGCTCCTGAGCCTCCCAAACCTTACAGCCTTGTTTCGGGAGTCTAGCAGTTCTCGCTC  
 ACCAACCCCAAAGCAGAAGAAGAAGAAAAAGAAAGATAGAGGACGCAGGTCAGAGAGCAGCTCTCCT  
 CGACGGGAGAGAAAGAAAAGCTCAAAGAAGAAGAAGCACAGGTCAGAATCTGAGTCCAAGAAACGTAAGC  
 ATAGGTCTCCCACTCAAAGAGCAAACGTAAATCTAAGGACAAAAGCGAAAGCGGTCTCGAAGTACAAC  
 5 ACCAGCCCCCAAGAGCCGCCGGGCCACCCTTCAACTTCTGCTGACTCTGCTTCTCTCCGATACTTCC  
 CGCAGTCGGTCTCGAAGTGCTGCAGCTAAAACCTCATACAACTGCCTTGGCTGGGGCGAAGTCTTCCCTG  
 CTTTCAGGGCGACGCGGGGAGGGAGATGCGCCTTTTCAGTGAACCAGGTACTACCAGCACACAACGGCCTAG  
 TAGCCCGGAGACTGCTACGAAACAGCCTAGCAGCCCTTATGAAGACAAAGATAAAGACAAGAAGGAGAAA  
 TCTGCAACTCGACCTAGCCCCCTCTCCGAAAGGAGCAGCAGAGGCCAGAACCACCTGCTCCCACTCCGC  
 10 TCCTTGCTGAGCGACATGGCGGCTCCCCACAACCCCTTGCAACCACCCCTTAAGCCAGGAGCCAGTGAA  
 CCCCCATCTGAGGCCTCTCCAACCTCGGGACCGTTACCACCTAAGTCTCCCGAGAAACTTCCCCAGTCT  
 TCTTCTCAGAGAGCAGCCCACCATCCCCCTCAACCTACCAAAGTTTCTCGGCATGCCAGCTCTTCCCCAG  
 AAAGTCCTAAACCTGCTCCAGCTCCAGGGTCCCACCGAGAGATTTCCTTCTTCTCCACATCTAAGAATCG  
 CTCACATGGCCGAGCAAACGGGATAAATCACATTCTCATACCCCTCCCGTAGGATGGGGAGGTCCCGT  
 15 AGCCCTGCCACCGCTAAGAGAGGGCGATCTCGGTCTCGAACCCTTACCAAGAGAGGTCAATTCTCGATCCC  
 GATCTCCCCAGTGGCGTAGGTCCAGGTCTGCACAGAGGTGGGGAAGATCTAGAAGCCCCCAGCGACGTGG  
 CCGCTCTAGGTCTCCTCAGCGACCAGGCTGGTCTAGGAGCAGAAATACCCAGAGAAGAGGCAGGTCTAGG  
 TCAGCAAGGCGAGGGAGGTCCCACTTAGATCCCCAGCCACTAGGGGTAGATCTCGTTCTAGAACCACAG  
 CCGCCGGGGCAGGTCCCGCTCTAGAACACCTGCCAGCGAGATCAGATCCAGAACTCCCACAGGCG  
 20 TAGGTCTCGGTCTAGAACCACCGCCGAGGGGCGAGGTCTCGGTCTAGAACCACCTGCTAGGCGCAGATCT  
 AGGACCCGATCACCAGTACGACGCAGGTCTCGTAGTAGATCACCAGCCAGGAGAAGTGGCAGGTCTACGCT  
 CTAGAACCCTCAGCTAGACGTGGCCGCTCACGCTCCAGAACCCTCAGCCAGACGTGGCCGCTCACGCTCTAG  
 AACCCCTCAGCTAGACGCAGTGGTCTCGCTCACGCTCCAGAACCACCGCCAGGAGAGGGAGGTCTCGGTCTAGG  
 ACACCAAGACGAGGAAGATCCCGCAGTAGAAGCTTAGTTAGACGTGGAAGATCTCACTCTAGAACCCTC  
 25 AAAGAAGAGGCAGATCTGGCTCATCTTCAGAGCGGAAAAACAAATCCAGAACATCTCAAAGAAGAAGCAG  
 GTCCAATTCAAGCCAGAAATGAAGAAATCTCGCATTTCTTCAAGGCGGAGCAGGTCTCTCTCTTACCA  
 CGGTCCAAAGCAAATCTCGCTTGTCTTTGAGGCGCAGCCTTTCAGGGTCTTCCCCATGCCCTAAGCAAA  
 AGTCACAGACACACCCAGGCGCAGTCGCTCTGGATCTTCCCAACCTAAAGCTAAATCTAGAACGCCACC  
 CAGACGCAGTCGCTCCAGTTCTTCTCCGCCACCTAAACAGAAATCTAAGACACCATCAAGACAAAGTCAT  
 30 TCCAGTTCATCTCCTCATCTAAAGTGAAATCTGGAACACCACCGAGGCAAGGGTCCATAACAAGTCCCC  
 AGGCCAATGAGCAATCTGTAACGCCACAGAGACGGAGCTGTTTTGAATCATCACCTGACCTGAGTTGAA  
 ATCTAGGACCCCTTCTAGACATAGCTGCTCAGGGTCTCTCCTCTAGAGTGAAATCTAGCACACCTCCC  
 AGACAGAGCCCATCTAGGTTCATCATCTCCACAACCCAAAGTGAAGGCAATAATATACCAAGACAAAGAA  
 GCCATTCTGGCTCCTCTTCTCCAAGTCTTAGTAGGGTGACGTCGAGAACAACCTCCACGGCGAAGCAGATC  
 35 AGTATCTCCCTGCTCCAATGTGGAATCCAGATTGTTGCCAAGATACAGTCATTCTGGGTCTCTCTCACC  
 GATACCAAAGTGAAACCTGAAACACCGCCAAGACAAAGTCATCAGGGTCTATTTACCATAACCCAAAG  
 TAAAGGCCCAAACCTCCACCGGGGCCAAGTCTTCTGGATCAAAGTCACCATGTCCCCAAGAGAAGTCTAA  
 AGACTCACTAGTTCAAGTTGCCCTGGATCCCTCTCTCTCTGTGCAGGAGTAAATCTAGCACACCACCA  
 GCGGAGAGCTATTTTGGTGTCTCATCTCTGCAACTGAAAGGACAATCTCAAACCTTACCAGACCACAGAT  
 40 CTGATACTTCAAGTCCAGAAGTGAGACAGAGTCATTGAGAATCACCATCTCTGCAGAGCAAATCTCAAAC  
 ATCACCTAAGGGAGGTGCGTCCAGGTCTTCATCTCCAGTCACTGAGCTGGCATCCAGATCTCCAATAAGA  
 CAAGATAGAGGTGAGTTCTCAGCGAGTCCTATGTTGAAATCTGGAATGTCTCTGAGCAGAGCAGGTTC  
 AGTCTGACTCTTCTTCATATCTACAGTGGACTCGAATCTCTCTTGGGGCAGAGTAGATTGGAGACTGC  
 TGAATCAAAGAGAAAATGGCCTTACCCCTCAGGAGGATGCTACTGCATCACCTCCTAGACAGAAAGAC  
 45 AAATTTAGTCCCTTTCCAGTACAGGATAGGCCTGAGTCTTCACTGGTATTCAAAGACACACTTAGAACCC  
 CGCCAAGAGAAAGAAGTGGTGTCTGGGTCTATCTCCAGAAACAAAAGAGCAAAATAGTGCATTGCCTACGTC  
 AAGCCAAGATGAAGAGTTAATGGAGGTGGTAGAGTCTGAAGAACCAGGCGCAAAATCCTGTCTCAT  
 TTGTCTTCAGAACTTAAAGAAATGTCCACAAGTAACCTTTGAATCATCTCCTGAAGTAGAAGAAAGCCCTG  
 CTGTGTCTTTGACTCTTGATCAGAGCCAGTCAACAGGCTTCTTTGGAAGCAGTAGAAGTCCCTTCAATGGC  
 50 CTCATCTTGGGGTGGGCCACATTTTTCTCCAGAACATAAAGAACTGTCTAACTCCCCACTCAGGGAGAAC  
 AGCTTTGGATCACCTTTAGAATTTAGAACTCAGGCCCACTTGGTACAGAAATGAATACTGGATTTTCTT  
 CTGAGGTTAAAGAAGATTTGAATGGACCGTTTCTTAATCAGCTGGAACAGATCCATCTCTAGACATGAA  
 AGAACATCGACAAGATCCTCTGGACACAGCAGTTCTGAGTTATCCCCAGATGCAGTGGAAAAGGCAGGG  
 ATGCTTTCAAATCAGAGCATCTCTTACCTGTGCTTGATGCTGTACCCAGAACACCCCTCGAGAGAAAGAA  
 55 GTAGTTCTGCATCTTCTCCTGAAATGAAAGATGGTTTACCAGAACTCCATCAAGGAGAGCAGGTCTGG  
 GTCTTCTCCAGGACTTAGAGATGGGTCTGGGACTCCCTCGAGGCACAGCCTGTCTGGGTCTCTCTGGGA  
 ATGAAAGATATACCTAGAACCCTATTTAGAGGGAGGAAGCGAATGTGATTCTTCCCCAGAACCAGAAAGCTT  
 TGCCTCAGACTCCTAGGCCGAGGAGTCGTTCTCCATCATCCCCAGAGCTCAACAACAAGTGTCTTACCCC  
 CCAGAGAGAAAGAAGCGGGTCAGAATCATCAGTTGATCAGAAAACCTGTGGCTCGGACTCCCCTGGGGCAG  
 60 AGAAGTCGTTCCGGATCCTCTCAAGAACTTGATGTGAAACCCAGTGCATCCCTCAGGAAAGAAGTGAGT  
 CAGACTCTTCTCCAGATTCTAAAGCCAAGACACGAACCCCACTTCGGCAGAGGAGTGGGTCTGGATCATC

TCCAGAGGTTGACAGCAAATCTCGACTATCCCCCTCGGCGCAGTAGGTCTGGTTCTCCCCCTGAAGTGAAA  
 GATAAGCCAAGAGCAGCACCCAGGGCACAGAGTGGTTCTGATTCTCTCTGAACCTAAAGCTCCAGCCC  
 CTCGGGGCCCTTCCCAGACGAAGCAGATCAGGTTTCATCAAGCAAAGGCAGAGGCCCTTCTCTGAAGGAAG  
 CAGCAGTACCGAGTCTCTCTGAACATCCGCCCAAATCCAGAAGTCTCGCAGAGGTTCCAGGTCATCA  
 5 CCAGAGCCCAAGACCAAGTCTCGTACACCACCTCGACGTCGAGCTCTCGATCATCTCCGAGCTAACAA  
 GGAAGGCCAGACTGTCCCGTAGAAGCCGCTCTGCCTCATCTCACCAGAAACTCGCTCTAGAAGTCCCCC  
 AAGGCACCGGAGAAGTCCCTCAGTGTCTTCCCCGGAGCCAGCCGAAAAATCGAGGTCTTCACGCCGACGG  
 CGCTCAGCTTCATCTCCACGCACTAAGACAACCTCAAGGAGAGGGCCGCTCTCTTTCGCCAAAGCCTCGTG  
 GACTCCAGAGGTCCCGTTCCCGCTCAAGGAGAGAGAAAACAAGAACAACCCGACGTCGAGATAGGTCTGG  
 10 ATCTTCTCAGTCAACCTCTCGGCGAAGACAGCGGAGCCGGTCAAGGTCTCGCGGTTACTCGGCGGCGGAGG  
 GGAGGCTCTGGTTATCACTCAAGGTCACTGCGCGGAGGAAAGTTCCCGGACCTCCTCTCGACGCCGAA  
 GAGGCCGCTCTCGGACACCCCCAACCACTCGGAAGCGTTCTCGCTCACGCACATCACCAGCCCCGTGGAA  
 ACGCTCTAGATCTCGAGCCTCTCCAGCCACTCACCGGCGATCCAGGTCCAGAACCCCCCTGATAAGCCGA  
 CGTAGGTCCAGATCTCGAACTTCACCAGTCAGCCGGAGACGGTCAAGGTCCAGGACTTCAGTGACTCGAC  
 15 GAAGATCCCGGTCAAGAGCATCCCCAGTGAGCAGAAGGCGATCCAGATCCAGAACGCCACCAGTAACCCG  
 CCGTCGTTCAAGGTCTAGAACGCCAACACACGCCGCCGCTCCCGTTCTAGAAGTCCACCAGTGACTCGC  
 AGAAGGTCCAGATCCAGGACTCCACCAGTAACCAGGAGGCGATCTCGAAGCAGAACTTCGCCTATCACTC  
 GCAGAAGATCAAGATCCAGAACATCTCCGGTCACCCGAAGGAGATCTCGATCTCGCACATCTCCAGTAAC  
 TCGAAGAAGGTCCCGCTCTCGAACCTCACCAGTGACACGCCGCCGCTCTAGGTCCCGGACACCTCCAGCT  
 20 ATTTCGGCGCGCTCTAGATCTCGAACGCCACTGTTACCACGCAAACGTTCTCGAAGTCGCTCACCACCTTG  
 CTATCCGCCGCGCTCCAGATCCCGTACTCCACGAACAGCTCGGGGTAAACGGTCTTAAACAAGATCTCC  
 TCCAGCCATCCGAGGCGTTCTGCATCTGGAAGTAGTTCTGATCGTTACGATCTGCTACTCTCCAGCA  
 ACAAGAAATCATTCTGGTTACGGACACCTCCAGTAGCACTCAACAGTTCCAGAATGAGCTGCTTCAGTC  
 GTCCTAGCATGTCCCCAACACCTCTTGATCGCTGCAGATCACCTGGAATGCTTGAACCCCTTGGCAGCTC  
 25 TAGAACACCCATGTCTGTCTGTCAGCAAGCCGGCGGCTCCATGATGGATGGTCCAGGTCCCCGAATACCT  
 GACCACCAGAGAACATCTGTGCCAGAAAAATCATGCTCAGTCCAGGATTGCACTTGCCCTGACAGCTATCA  
 GTCTTGGCACCGCTCGGCCTCCTCCGTCCATGTCTGCTGCTGGCCTTGCTGCAAGAATGTCCAGGTTCC  
 AGCCCCGGTGCTCTCATGAGTCTCAGAACCGCACCAGCAGCCAACCTTGCCAGCAGGATTCCTGCGACCC  
 TCTCGGCGCAGCATGAACCTAGCCAGCGCCAGGACACTGCCATTCCAACAGCAGTGAACCTGGCTGACT  
 30 TCTGAACGCGCATGTCAGCAGCGGCCATGAACCTTGCCAGCCCCAGAACAGCGGTGGCACCTTCCGGTGT  
 GAACCTGGCTGACCTTCGCACTCCACAGCCCCAGCTGTGAACCTAGCAGGGGCCAGAACCCCCAGCTGCC  
 TTGGCAGCTCTGAGTCTCACAGGCTCTGGCACACCACCAACTGCTGCAAACTATCCCTCCAGCTCCAGAA  
 CACCACAGGCTCCAGCCTCTGCAAACTGGTGGGTCTCCGCTGTCACATGCCACAGCTCCTGTGAATAT  
 TGCCGGCTCCAGAACCGCCGAGCCTTGCCCCCGCGAGCCTCACCAGTGCTAGGATGGCTCCAGCATTG  
 35 TCTGGTGCAACCTCACCAGCCCCAGGGTGCCCCCTTCTGCTACGAGCGTGTCTAGTGGCAGAACCTCAC  
 CACCGCTCCTTGACCGAGCTAGGTCCAGAACACCACCGTCTGCCCAAGCCAATCTAGGATGACCTCTGA  
 ACGGGCTCCCTCCCCTTCTCTAGAATGGGCCAGGCTCCTTCACAGTCTCTTCTCCCTCCAGCACAGGAT  
 CAGCCGAGGTCTCCTGTGCTTCTGCTTTTTCAGACCAATCCCGTTGTTTGATTGCCAGACCACCCCTG  
 TAGCAGGGTCTCAGTCCCTTCTCTGCGGCGAGTGGAACGACCAGTCTCTGTGCTGGTGATCACAATGG  
 40 CATGCTCTCTGTCCCTGCCCCCTGGGGTGCCCCACTCTGATGTGGGGGAGCCACCTGCCTCTACTGGGGCC  
 CAGCAGCCTTCTGCATTAGCCGCCCTGCAGCCAGCAAAGGAGCGGCGAGTTCTCTCTCGTCGTCGTCGT  
 CCTCTAGCTCCTCCTCCTCTTCATCATCGTCGTCGTCGTCCTCCTCCTCTCTGGCTCCAGTTCTAGTGA  
 CTCAGAGGGCTCTAGCCTTCTGTGCAACCTGAGGTGGCACTGAAGAGGGTCCCCAGCCCCACCCAGCC  
 CCAAAGGAGGCTGTTTCGAGAGGGACGTCTCCGGAGCCAACCCAGCCAAACGGAAGAGGCGCTCTAGCA  
 45 GTTCCAGTTCCAGCTCCTCCTCTTCATCTTCTCCTCCTCCTCCTCCTCCTCCTCCTCCTCCTCCTCCTC  
 CTCTTCTTCTTCTTCTCCTCCTCATCTTCTCCTCCTCCTCGTCGTCCTTCTCCTCCTTCCCCTGCTAAGCCTGGC  
 CCTCAGGCCTTGCCCAACCTGCAAGCCCCAAGAAGCCACCCCTGGCGAGCGGAGGTCCCGCAGCCCCC  
 GGAAGCCAATAGACTCCCTCAGGGACTCTCGTCCCTCAGCTACTCGCCTGTGGAGCGCTCGCCGTCCCTC  
 GCCCCAGCCCTCACCACGGGACCAGCAGAGCAGCAGTGAAGCGGGTTCCCGGAGAGGCCAGCGTGGG  
 50 GACAGCCGCTCCCCAGCCACAAGCGCAGGAGGGAGACACCTAGCCCTCGGCCCATGAGACACCGCTCCT  
 CCAGGTCTCCATAAATTGTCTTTGGGGGATTCCACCACACCCAATGCTCTGGAGCCACAAGGAGTGTCCC  
 TTCTTCCCCAGCAGAGCCGTGGGAGGGTCTTGTCTGCTCTCCTTTGAACCTTGGCAGCCCTTGGATGGA  
 GGGCTCCCTTTCCCTCCCCTTTTTTTTTTCTTTGTTCTGTGAAATGTTAATCTCCGTGAGTTCTTCTCTG  
 GTTCATGTGTTCTGGGGGGTTTGGGGTGGGAGGGAATGCAGATGGGAGTTGGGGGAGGGGAGGATACAGT  
 55 TCAGGATACCCAGCCTGGAGTCAGGGCCAGGGAGGCTGGCCCCACTTGTATCCAGAAGTCCCAGGGG  
 TAGTTGTGATGGTGGTGGGACTGGAGGTTGTATAAGGTGTTCTTGGAAGGAAGGGGAGGAGTTGGAAT  
 TGATTGGTCCCTAGTGTCCCCCATGAGGTTGTGAACCCCTCCCCCAACTTTTCATGTTTCTTAAAGGCA  
 TTTTGGTTTTTTTAAATCTGTACAGCAAGAGCAACTTTTTCTGTCAAATAAAAATGAGAAATGCAGG

60 >gi|19923466|ref|NP\_057417.2| splicing coactivator subunit SRm300; RNA  
 binding protein; AT-rich element binding factor [Homo sapiens]

MYNGIGLPTPRGSGTNGYVQRNLSLVRGRRGERPDYKGEELRRLLEAALVKRPNPDILDHERKRRVELRC  
 LELEEMMEEQGYEEQIQEKVATFRLMLEKDVNPGGKEETPGQRPVAVTETHQLAELNEKKNERLRAAFG  
 ISDSYVDGSSFDPPQRRAREAKQPAPEPPKPYSLVRESSSSRSPTPKQKKKKKKDRGRSSESSSPRRERK  
 KSSKKKKHRSESESKKRKHSRSTPKSKRKSDDKKRKRSTTPAPKSRRHRSTADSASSSDTSRSTRSR  
 5 SAAAKTHTTALAGRSPSPASGRRGEGDAPFSEPGTTSTQRPSSPETATKQPSSPYEDKDKDKKEKSATRP  
 SPSERSSTGPEPPAPTPLLAERHGGSPQPLATTPLSQEPVNPPEASPTRDRSPPKSPEKLPQSSSES  
 SPPSPQPTKVSRRHASSSPESPKPAPAPGSHREISSSPTSKNRSHGRAKRDKSHSHTPSRRMGRSRSPATA  
 KRGRSRSTPTKRGRSRSRSPQWRRSRSAQRWGRSRSPQRRGRSRSPQRPGWSRSRNTQRRGRSRARRG  
 RSHSRSPATRGRSRSRSTPARRGRSRSRSTPARRRSRSTPTRRRSRSRSTPARRGRSRSRSTPARRRSRTRSP  
 10 VRRRSRSTRSPARRSGRSRSTPARRGRSRSRSTPARRGRSRSTPARRSGRSRSTPARRGRSRSTPRRG  
 RSRSRSLVRRGRSHSRTPQRRGRSGSSSERKNKSRTSQRRSRSNSSPEMKKSRISSRRSRSLSSPRSKAK  
 SRLSLRRSLSGSSPCPKQKSQTTPRRRSRSGSSQPKAKSRTPRRRSRSSSSPPPKQKSQTSPRQSHSSSSP  
 HPKVKSQTTPRQGSITSPQANEQSVTPQRRSCFESSPDPELKSRTPSRHSCSGSSPPRVKSSTPPRQSPS  
 RSSSPQPKVKAIISPRQRSHSGSSSPSRVTSRTTPRRRSRVSPCS NVESRLLPYSHSGSSSPDTKVK  
 15 PETPPRQSHSGSISPYPKVKAQTPPGPSLSGSKSPCQEKSKDSLQSCPGSLSLCAGVKSSTPPGESYF  
 GVSSLQLKGQSQTSPDHRSDTSSPEVRQSHSESPSLQSKSQTSPKGGRSRSSSPVTELASRSPIRQDRGE  
 FSASPMLKSGMSPEQSRFQSDSSSYPTVDSNLLGQSRLETAESKEKMALPPQEDATASPPRQDKFSPF  
 PVQDRPESSSLVFKDTRLTPPRERSGAGSSPETKEQNSALPTSSQDEELMEVVEKSEEPAGQILSHLSEL  
 KEMSTSNFESSPEVEERPAVSLTLDQSQSQASLEAVEVPSMASSWGGPHFSPPEHKELSN SPLRENSFGSP  
 20 LEFRNSGGLPTGEMNTGFSSEVKEDLNGPFLNQLTDPISLDMKEQSTRSSGHSSSELSPDAVEKAGMSSNQ  
 SISSPVLDAYVPTPSRERSSSASSPEMKDGLPRTPSRRSRSGSSPGLRDGSGTSPRHSLSGSSPGMKDIP  
 RTPFRGRSECDSSPEPKALPQTTPRPSRSPSSPELNNKCLTPQRERSGSESSVDQKTVAARTPLGQRSRG  
 SSQELDVKPSASPQERSESDSSPDSKAKTRTPLRQRSRGSSPEVDSKRLSPRRSRSGSSPEVKDKPRA  
 APRAQSGSDSSPEPKAPAPRALPRRSRGSSSKGRGPSPEGSSSTESSPEHPPKSRTARRGSRSSPEPKT  
 25 KSRTPPRRRSRSPPELTKARLSRRSRSSASSPETRSRTPPRHRRSPSVSSPEPAEKSRSSRRRRSASS  
 PRTKTTSRGRSPSPKPRGLQRSRSRREKTRTTRRRDRSGSSQSTSRRRQRSRSRSTVTRRRRGSGY  
 HSRSPARQESSRTSSRRRRGRSRTPTTSRKRSRSTSPAPWKRSRSPATHRRSRSTPLISRRRSRS  
 RTSVPVRRRSRSTSVTRRRRSRSPVSRRRSRSTPPVTRRRRSRSTPTTRRRRSRSTPPVTRRRRS  
 RTPPVTRRRRSRSTSPITRRRSRSTSPVTRRRRSRSTSPVTRRRRSRSTSPVTRRRRSRSTPPAIRRS  
 30 RSRTPLLRKRSRSPLAIRRRSRSTPRTARGKSLTRSPPAIRRRSAGSSSDRSRATPPATRNHS  
 GSRTPPVALNSSRMSCFRSPMSPTPLDRCSRPGMLEPLGSSRTPM SVLQQAGGSMMDGPGPRI PDHQRT  
 SVPENHAQSRIALALTAISLGTARPPPSMSAAGLAARMSQVPAPVPLMSLRTAPAAANLASRIPAASAAAM  
 NLASARTPAIPTAVNLADSRTPAAAAAMNLASPTAVAPSAVNLA DPRTPTAPAVNLAGARTPAALAALS  
 LTGSGTPPTAANYPSSSRTPQAPASANLVGPRSAHATAPVNIAGSR TAAALAPASLTSARMAPALSGANL  
 35 TSPRVPLSAYERVSGRTSPPLLD RARSRTPPSAPSQSRMTSERAPSPSSRMGQAPSQSLLP AQDQPRSP  
 VPSAFSDQSRCLIAQTTPVAGSQSLSSGAVATTTSSAGDHNGMLSV PAPGVPHSDVGEPPASTGAQQPSA  
 LAALQPAKERRSSSSSSSSSSSSSSSSSSSSSSSSSSSSSSSSSGSSSSDSEGSSLPVQPEVALKRVSP TPAPKEAV  
 REGRPPEPTPAKRKRSSSPAPKPGPQALP  
 KPASPCKPPGERRSRSPRKPIDSLRDSRSLSYSPVERRRSPSPQPSPRDQSSSSSERGSRRGQRGDSRSP  
 40 SHKRRRETSPSRPMRHRSSRSP

>gi|6649241|gb|AF201422.1|AF201422 Homo sapiens splicing coactivator  
 subunit SRm300 (SRM300) mRNA, complete cds  
 GCAGCGCGGCCAGGCGGGGTGCGAGTGGCGCAGTCGGAGCCCGTTGCGGCCCTGAGGAAGCGAGGAGG  
 45 CGTCGGCGTCGGCTGAGGCGGGCGGACCGGCGAGGCGAGGCGGGCGGCCCCAGGCCCGGGGACTCGGGAGC  
 TCGAGCAGCGGCGGGCGGCAAGACCTCTCCCCCTCGGAGGCGGGCGGGCGGAGGCGGGGAGCGGTGGTG  
 CCCCCCGGGCACGGGGCCATGTACAACGGGATCGGGCTGCCGACGCCCGGGGACGCGGCACCAACGGC  
 TACGTCCAGCGCAACCTGTCCCTGGTGCGGGGCCCGGGGTGAGCGGCCTGACTACAAGGGAGAGGAGG  
 AACTGCGGCGCCTGGAGGCTGCCCTGGTGAAAGCGGCCTAATCCTGACATCCTGGACCACGAGCGCAAGCG  
 50 GCGCGTCGAGCTGCGATGCCTCGAGCTGGAGGAGATGATGGAAGAGCAAGGGTACGAGGAACAGCAAATT  
 CAGGAAAAAGTGGCGACCTTTGACTCATGTTGCTGGAGAAGGATGTGAACCCTGGGGGCAAGGAGGAGA  
 CCCAGGGCAGAGGCCAGCGGTACGGGAGACTACCAAGTTGGCAGAATTAAATGAGAAGAAGAATGAAAG  
 ACTCCGTGCTGCCTTTGGCATCAGTGATTCTTACGTAGATGGCAGCTCTTTTGATCCTCAGCGTCGTGCC  
 CGAGAAGCTAAACAACCAGCTCCTGAGCCTCCCAAACCTTACAGCCTTGTTTCGGGAGTCTAACAATTCTC  
 55 GCTCACAAACCCCAAAGCAGAAGAAGAAAAAGAAAGATAGAGGACGCAGGTACAGAGACGAGCTCT  
 CCTCGACGGGAGAGAAAGAAAGCTCAAAGAAGAAGAACACAGGTACAGATCTGAGTCCAAGAAACGTA  
 AGCATAGGTCTCCCACTCAAAGAGCAAACGTAAATCTAAGGACAAAAGCGAAAGCGGTCTCGAAGTAC  
 AACACCAGCCCCAAGAGCCGCGGGCCACCCTTCAACTTCTGCTGACTCTGCTTCTCTCTCCGATACT  
 TCCCGCAGTCGGTCTCGAAGTGCTGCAGCTAAACTCATACAACCTGCCTTGGCTGGGCGAAGTCTTCCC  
 60 CTGCTTCAGGGCGACGCGGGGAGGAGATGCGCCTTTCAGTGAACCAAGTACTACCAAGCACACAACGGG  
 CTAGTAGCCCGGAGACTGCTACGAAACAGCCTAGCAGCCCTTATGAAGACAAAGATAAAGACAAGAAGGA

GAAATCTGCAACTCGACCTAGCCCCCTCTCCGGAAGGAGCAGCACAGGCCAGAACACCTGCTCCCACT  
 CCGCTCCTTGCTGAGCGACATGGCGGCTCCCCACAACCCCTTGCAACCACCCCTTAAGCCAGGAGCCAG  
 TGAACCCCCCATCTGAGGCCTCTCCAACCTCGGGACCGTTACCACCTAAGTCTCCCGAGAACTTCCCCA  
 GTCTTCTTCTCAGAGAGCAGCCACCATCCCCCTCAACCTACCAAAGTTTCTCGGCATGCCAGCTCTTCC  
 5 CCAGAAAGTCTTAAACCTGCTCCAGCTCCAGGTCCCACCGAGAGATTTCTTCTTCTCCACATCTAAGA  
 ATCGCTCACATGGCCGAGCAAAACGGGATAAATCACATTCTCATACCCCTCCCGTAGGATGGGGAGGTC  
 CCGTAGCCCTGCCACCGCTAAGAGAGGGCGATCTCGGTCTCGAACCCCTACCAAGAGAGGTCATTCTCGA  
 TCCCGATCTCCCCAGTGGCGTAGGTCCAGGTCTGCACAGAGGTGGGAAGATCTAGAAGCCCCCAGCGAC  
 GTGGCCGCTCTAGGTCTCCTCAGCGACCAGGTGGTCTAGGAGCAGAAATACCCAGAGAAGAGGCAGGTCT  
 10 TAGGTACGAAGGCGAGGGAGGTCCCCTCTAGATCCCCAGCCACTAGGGGTAGATCTCGTTCTAGAACA  
 CCAGCCCGCCGGGCGAGGTCCCGCTCTAGAACACCTGCCAGGCGGAGATCAGATCCAGAACTCCCACCA  
 GCGTAGGTCTCGGTCTAGAACACCAGCCCGAGGGGCGAGGTCTCGGTCTAGAACACCTGCTAGGCGCAG  
 ATCTAGGACCCGATCACCAGTACGACGCAGGTCTCGTAGTAGATCACCAGCCAGGAGAAGTGGCAGGTCA  
 CGCTCTAGAACCCAGCTAGACGTGGCCGCTCACGCTCCAGAACCCAGCCAGACGTGGCCGCTCACGCT  
 15 CTAGAACCCAGCTAGACGCAGTGGTCTGCTCACGCTCCAGAACACCAGCCAGGAGAGGGAGGTCTCGGTCT  
 TAGGACACCAAGACGAGGAAGATCCCGCAGTAGAAGCTTAGTTAGACGTGGAAGATCTCACTCTAGAACA  
 CCTCAAAGAAGAGGCAGATCTGGCTCATCTTCAGAGCGGAAAAACAAATCCAGAACATCTCAAAGAAGAA  
 GCAGGTCCAATTCAAGCCCAGAAATGAAGAAATCTCGCATTTCTTCAAGGCGGAGCAGGTCTCTCTCTTC  
 ACCACGGTCCAAGCAAAATCTCGCTGTCTTTGAGGCGCAGCCTTTTCAAGGTCTTCCCCATGCCCTAAG  
 20 CAAAAGTACAGACACCACCCAGGCGCAGTCTGCTTCTGATCCTCCCAACCTAAAGCTAAATCTAGAACGC  
 CACCCAGACGCAGTCTGCTCCAGTTCTTCTCCGCCACCTAAACAGAAATCTAAGACACCATCAAGACAAAG  
 TCATTCCAGTTTCTCTCATCTCTAAAGTGAATCTGGAACACCACCGAGGCAAGGGTCCATAACAAGT  
 CCCCAGGCCAATGAGCAATCTGTAACGCCACAGAGACGGAGCTGTTTTGAATCATCACCTGACCTGAGT  
 TGAAATCTAGGACCCCTTCTAGACATAGCTGCTCAGGTCTCTCTCTCTAGAGTGAAATCTAGCACACC  
 25 TCCCAGACAGAGCCCATCTAGGTCTCATCTCTCCACAACCCAAAGTGAAGGCAATAATATCACCAGACAA  
 AGAAGCCATTCTGGCTCCTCTTCTCCAAGTCCTAGTAGGGTGACGTCGAGAACAACCTCCACGGCGAAGCA  
 GATCAGTATCTCCCTGCTCCAATGTGGAATCCAGATTGTTTGCCAAGATACAGTCATTCTGGGTCTCTCTC  
 ACCAGATACCAAAGTGAACCTGAAACACCGCCAAGACAAAGTCACTCAGGGTCTATTTACCATAACCC  
 AAGTAAAGGCCCAAACCTCCACGGGGCCAAGTCTTTCTGGATCAAAGTCAAGTCCCCCAAGAGCAAGT  
 30 CTAAGACTCACTAGTTCAAAGTTGCCCTGGATCCCTCTCTCTCTGTGCAGGAGTAAATCTAGACACC  
 ACCAGGCGAGAGCTATTTTGGTGTCTCATCTCTGCAACTGAAAGGACAATCTCAAACCTTACCAGACCAC  
 AGATCTGATACTTCAAGTCCAGAAGTGAGACAGAGTCATTTCAGAATCACCATCTCTGCAGAGCAAATCTC  
 AAACATCACCTAAGGGAGGTCCGTCCAGGTCTTCATCTCCAGTCACTGAGCTGGCATCCAGATCTCCAAT  
 AAGACAAGATAGAGGTGAGTTCTCAGCGAGTCTATGTTGAAATCTGGAATGTCTCTCTGAGCAGAGCAGG  
 35 TTCCAGTCTGACTCTTCTTCATATCCTACAGTGGACTCGAATTTCTCTCTTGGGGCAGAGTAGATTGGAGA  
 CTGCTGAATCAAAGAGAAAATGGCCTTACCCCTCAGGAGGATGCTACTGCATCACCTCCTAGACAGAA  
 AGACAAATTTAGTCCCTTTCCAGTACAGGATAGGCCTGAGTCTTCACTGGTATTCAAAGACACACTTAGA  
 ACCCGCCAAGGGAAGAAGTGGTCTGGGTCTATCTCCAGAAACAAAGAGCAAAAGTAGTCATTTGCCTA  
 CGTCAAGCCAAGATGAAGAGTTAATGGAGGTGGTAGAGAAGTCTGAAGAACCCGAGGCCAAATCCTGTCT  
 40 TCATTTGTCTTCAGAACTTAAAGAAATGTCCACAAGTAACCTTTGAATCATCTCTCTGAAGTAGAAGAAAGG  
 CCTGCTGTGTCTTTGACTCTTGATCAGAGCCAGTCACAGGCTTCTTTGGAAGCAGTAGAAGTCCCTTCAA  
 TGGCCTCATCTTGGGGTGGGCCACATTTTCTCCAGAACATAAAGAAGTGTCTAACTCCCCACTCAGGGA  
 GAACAGCTTTGGATCACCTTTAGAATTTAGAACTCAGGCCACTTGGTACAGAAATGAATACTGGATTT  
 TCTTCTGAGGTTAAAGAAGATTTGAATGGACCGTTTCTTAATCAGCTGGAACAGATCCATCTCTAGACA  
 45 TGAAAGAACAATCGACAAGATCCTCTGGACACAGCAGTTCTGAGTTATCCCCAGATGCAGTGGAAAAGGC  
 AGGGATGTCTTCAAATCAGAGCATCTCTTACCTGTGCTTGATGCTGTACCCAGAACACCCCTCGAGAGAA  
 AGAAGTAGTTCTGCATCTTCTCCTGAAATGAAAGATGGTTTACCCAGAACTCCATCAAGGAGAAGCAGGT  
 CTGGGTCTTCTCCAGGACTTAGAGATGGGTCTGGGACTCCCTCGAGGCACAGCCTGTCTGGGTCTCTCC  
 TGGAAATGAAAGATATACCTAGAACGCCATCTAGAGGGAGAAGCGAATGTGATTCTTCCCCAGAACCGAAA  
 50 GCTTTGCCCTCAGACTCCTAGGCCGAGGAGTCGTTCTCCATCATCCCCAGAGCTCAACAACAAGTGTCTTA  
 CCCCCAGAGAGAAAGAAGCGGGTCAGAATCATCAGTTGATCAGAAAAGTGTGGCTCGGACTCCCCTGGG  
 GCAGAGAAGTCGTTCCGGATCCTCTCAAGAACTTGATGTGAAACCCAGTGCATCCCCCTCAGGAAAGAAGT  
 GAGTCAGACTCTTCTCCAGATTCTAAAGCCAAGACACGAACCCCACTTCGGCAGAGGAGTCGGTCTGGAT  
 CATCTCCAGAGGTTGACAGCAAATCTCGACTATCCCCCTCGGCGCAGTAGGTCTGGTCTCTCCCTGAAGT  
 55 GAAAGATAAGCCAAGAGCAGCACCCAGGGCACAGAGTGGTTCTGATTCTCTCTGAACTAAAGCTCCA  
 GCCCTCGGGCCCTTCCCAGACGAAGCAGATCAGGTTTCATCAAGCAAGGAGGAGCCCTTCTCCTGTAAG  
 GAAGCAGCAGTACCGAGTCTCTCCTGAAACATCGGCCCAATCCAGAAGTCTCGCAGAGGTTCCAGGTC  
 ATCACCAGAGCCCAAGACCAAGTCTCGTACACCACCTCGACGTCGCAGCTCTCGATCATCTCCGGAGCTA  
 ACAAGGAAGGCCAGACTGTCCCGTAGAAGCCGCTCTGCCTCATCTCACCAGAACTCGCTCTAGAAGTCTC  
 60 CCCCAGGACCCGAGAGAGTCCCTCAGTGTCTTCCCCGAGCCAGCCGAAAAATCGAGGTCTTACGCGCG  
 ACGGCGCTCAGCTTCATCTCCACGCACTAAGACAACCTCAAGGAGAGGCGCTCTCTCTCGCCAAAGCCT

CGTGGACTCCAGAGGTCCCGTTCCCGCTCAAGGAGAGAGAAAACAAGAACAACCCGACGTCGAGATAGGT  
 CTGGATCCTCTCAGTCAACCTCTCGGCGAAGACAGCGGAGCCGGTCAAGGTCGCGGGTTACTCGGCGGCG  
 GAGGGGAGGCTCTGGTTATCACTCAAGGTCACCTGCCCGGAGGAAAGTTCCCGGACCTCCTCTCGACGC  
 CGAAGAGGCCGCTCTCGGACACCCCCAACCCAGTCGGAAGCGTTCTCGCTACGCACATCACCAGCCCCGT  
 5 GGAACGCTCTAGATCTCGAGCCTCTCCAGCCACTCACCGGCGATCCAGGTCCAGAACCCCCCTGATAAG  
 CCGACGTAGGTCCAGATCTCGAACTTCACCAGTCAGCCGGAGACGGTCAAGGTCCAGGACTTCAGTGACT  
 CGACGAAGATCCCGGTCAAGAGCATCCCCAGTGAGCAGAAGGCGATCCAGATCCAGAACGCCACCAGTAA  
 CCCGCCGTCGTTCAAGGTCTAGAACGCCAACACGCCGCCGCTCCCGTTCTAGAACTCCACCAGTGAC  
 TCGCAGAAGGTCCAGATCCAGGACTCCACCAGTAACCAGGAGGCGATCTCGAAGCAGAACTTCGCCTATC  
 10 ACTCGCAGAAGATCAAGATCCAGAATCTCCGGTCACCCGAAGGAGATCTCGATCTCGCACATCTCCAG  
 TAACTCGAAGAAGGTCCCGCTCTCGAACCTCACCAGTGACACGCCGCCGCTCTAGGTCGCCGACACCTCC  
 AGCACAGGATCAGCCGAGGTCTCCTGTGCTTTCTGCTTTTTCAGACCAATCCCGTTGTTTGTATTGCCAG  
 ACCACCCCTGTAGCAGGGTCTCAGTCCCTTTCTCTGCGGAGTGGCAACGACCACGTCCTCTGCTGGTG  
 ATCACAATGGCATGCTCTCTGTCCCTGCCCTGGGGTGGCCACTCTGATGTGGGGGAGCCACCTGCCTC  
 15 TACTGGGGCCAGCAGCCTTCTGCATTAGCCGCCCTGCAGCCAGCAAAGGAGCGGCGGAGTTCTCTCTCG  
 TCGTCGTCGTCCTCTAGCTCCTCCTCTTCTCATCATCGTCGTCGTCCTCCTCCTCCTCTGCTCCA  
 GTTCTAGTGACTCAGAGGGCTCTAGCTTCTGTGCAACCTGAGTGGCACTGAAGAGGTCCCCAGCCCCAC  
 CCCAGCCCCAAAGGAGGCTGTTTCGAGAGGGACGTCCTCCGGAGCCAACCCAGCCAAACGGAAGAGGCGC  
 TCTAGCAGTTCCAGTTCAGCTCCTCCTCTTCTCCTCTCCTCCTCCTCCTCCTCCTCCTCCTCCTCCTCCT  
 20 CCTCTTCTCCTTCTTCTTCTTCTTCTCCTCCTCCTCCTCCTCCTCCTCCTCCTCCTCCTCCTCCTCCTCCT  
 AGCCTGGCCCTTGGCCCTTGGCCAAACCTGCAAGCCCCAAGAAGCCACCCCTGGCGAGCGGAGTCCCG  
 CAGCCCCCGGAAGCCAATAGACTCCCTCAGGGACTCTCGGTCCCTCAGCTACTCGCCTGTGGAGCGTCGC  
 CGTCCCTCGCCCCAGCCCTCACCACGGGACCAGCAGAGCAGCAGCAGTGAGCGGGGTTCGCGAGAGGCC  
 AGCGTGGGGACAGCCGCTCCCCAGCCACAAGCGCAGGAGGGAGACACCTAGCCCTCGGCCCATGAGACA  
 25 CCGCTCCTCCAGGTCTCCATAAATTGTCTTTGGGGGATTCCACCACACCCAATGCTCTGGAGCCACAAGG  
 AGTGTCCCTTCTTCCCCAGCAGAGCCGTGGGAGGGTCTTGTCTGCTCTCCTTTGAACCTTGGCAGCCCT  
 TGGATGGAGGGCTCCCTTTCCCTCCCTTTTTTTTTTCTTTGTTCCTGTGAAATGTTAATCTCCGTGAGT  
 TCTTCTGTGTTTATGTTTTCTGGGGGGTGTGGGGTGGGAAGGAATCAAATGGGAATTGGGGGAAGGGAA  
 GATACAGTTTCAGGATACCCAGCCTGGAATTAGGGCAGGGAAGCATGGCCCCACTTGATACCGAAGTT  
 30 ACCAGGGGTGATTGTGATGGTGGTGGGACTGGAGTTGTATAAGGTGTTCTTGAAAGGAGGTGGAAT  
 TAGTTGGTCCCTACTGTCCCCCATGAGGTTGTGAACCCCTCCCCCAACTTTTTCATGTTTCTTAAAGGCA  
 TTTTGGTTTTTTTAAATCTGTACAGCAAGAGCAACTTTTTCTGTCAAATAAAAATGAGAAATGCAGGAAA  
 AAAAAAAAAAAAAAAAAA

35 >gi|6649242|gb|AAF21439.1|AF201422\_1 splicing coactivator subunit SRm300  
 [Homo sapiens]  
 MYNGIGLPTPRSGTNGYVQRNLSLVRGRRGERPDYKGEELRRLEAALVKRPNPDILDHERKRRVELRC  
 LELEEMMEEQGYEEQIQEKVATFRLMLLEKDVNPGGKEETPGQRPVAVTETHQLAELNEKKNERLRAAFG  
 40 ISDSYVDGSSFDPPRRAREAKQPAPEPPKPYSLVRESNNSRSQPQSRRRRKRRKIEDAGQRAALLDGRER  
 KAQRRRSTGQNLSPRNVSIGLPLQORANVNLRTKSESGLVQHQPRAAGPTVQLLLTLLPPLILPAVGLE  
 VLQLKLIQLPWLGEVLPLLQGDAGREMRLSVNQVLPSTQRASSPETATKQPSSPYEDKDKDKKEKSATRP  
 SPSERSSTGPEPPAPTPLLAERHGGSPQPLATTPLSQEPVNPPEASPTRDRSPPKSPEKLPQSSSES  
 SPPSPQPTKVSRRHASSPESPKPAPAPGSHREISSSPTSKNRSHGRAKRDKSHSTPSRRMGRSRSPATA  
 45 KRGRSRRTPTKRGRHSRSPQWRRSRSAQRWGRSRSPQRRGRSRSPQRPWGRSRNTQRRGRSRARRG  
 RSHSRSPATRGRSRRTPARRGRSRSPARRRSRRTPTRRRSRRTPARRGRSRSPARRRSRTRSP  
 VRRRSRSPARRSGRSRRTPARRGRSRSPARRGRSRSPARRSGRSRSPARRGRSRSPARRGRSRTPRRG  
 RSRSLVRRGRSHSRTPQRRGRSGSSSERKNKRTSQRRSRNSSPPEMKKSRISSRRSRSLSSPRSKAK  
 SRLSLRRSLSGSSPCPKQKSQTPPRRSRSGSSQPKAKSRTPRRSRSSSSPPPKQKSQTSPRQSHSSSSP  
 50 HPKVKSQTTPRQGSITSPQANEQSVTPQRRSCFESSPDPELKSRTPSRHSCSGSSPPRVKSSTPPRQSPS  
 RSSSPQPKVKAIISPRQRSHSGSSSPSRVTSRTTPRRSRSVSPCSNVESRLLPYSHSGSSSPDTKVK  
 PETPPRQSHSGSISPYPKVKAQTPPGPSLSGSKSPCPQEKSKDSLQSCPGSLSLCAGVKSSSTPPGESYF  
 GVSSLQLKGQSQTSPDHRSDTSSPEVRQSHSESPSLQSKSQTSPKGGRSRSSSPVTELASRSPIRQDRGE  
 FSASPMLKSGMSPEQSRFQSDSSSYPTVDSNLLGQSRLETAESKEKMALPPQEDATASPPRQDKFSPF  
 55 PVQDRPESSLVFKDTRLTPPRERSGAGSSPETKEQNSALPTSSQDEELMEVVEKSEEPAGQILSHLSSEL  
 KEMSTSNFESSPEVEERPAVSLTLDQSQSQASLEAVEVPSMASSWGGPHFSPHKEKLSNSPLRENSFGSP  
 LEFRNSGPLGTEMNTGFSSEVKEDLNGPFLNQLTDPISLDMKEQSTRSSGHSSSELSPDAVEKAGMSSNQ  
 SSSPVLDAVPRTPSRERSASSPEMKDGLPRTPSRRSRSGSSPGLRDGSGTSPSRHSLSGSSPGMKDIP  
 60 RTPSRGRSECDSSPEPKALPQTPRPRSRSPSSPELNNKCLTPQRERSGSESSVDQKTAVARTPLGQSRSG  
 SSQELDVKPSASPQERSESDSSPDSKAKTRTPLRQRRSRSGSSPEVDSKRLSPRRSRSGSSPEVKDKPRA  
 APRAQSGSDSSPEPKAPAPRALPRRSRSGSSSKGRGPSPEGSSSTESSPEHPPKSRTARRGRSSPEPKT  
 KSRTPPRRRSSRSPELTRKARLSRRSRSSSPETRSTPPRHRRSPSVSSPEPAEKSRSSRRRSASS

GH1-90-PCR-G3F1

15 GCTCCCTAAAGAACAAAGCTCAGGCGCGGGGACCTGCCGTTTGTCTGTGCCCCGCCGAATGGCCCGGAAAGGCG  
CCGGGGATGGCTCCGACGAAGAGGTAGATGGCAAAGCGGATGGGGCTGAGGCCAAACCTGCCGAATAAGCCT  
CTTCTCCTGCAGCCTGAGATGGATGGACAGACAGACACCACAGCCTCCCC

```
>gi|22507396|ref|NM_002473.2| Homo sapiens myosin, heavy polypeptide 9,
non-muscle (MYH9), mRNA
```

ATGGCACAGCAAGCTGCCGATAAGTATCTCTATGTGGATAAAAACTTCATCAACAATCCGCTGGCCAGG  
20 CCGACTGGGCTGCCAAGAAGCTGGTATGGGTGCCTTCCGACAAGAGTGGCTTTGAGCCAGCCAGCCTCAA  
GGAGGAGGTGGGCGAAGAGGCCATCGTGGAGCTGGTGGAGAATGGGAAGAAGGTGAAGGTGAACAAGGAT  
GACATCCAGAAGATGAACCCGCCAAGTTCTCCAAGGTGGAGGACATGGCAGAGCTCACGTGCCTCAACG  
AAGCCTCGGTGCTGCACAACCTCAAGGAGCGTTACTACTCAGGGCTCATCTACACCTATTTCAGGCCTGTT  
25 CTGTGTGGTTCATCAATCCTTACAAGAACCTGCCCATCTACTCTGAAGAGATTGTGGAATGTACAAGGGC  
AAGAAGAGGACGAGATGCCCTTACATCTATGCCATCAGACACACCGCTACAGAGATATGATGCAAG  
ACCGAGAAGATCAATCCATCTTGTGCACTGGTGAATCTGGAGCTGGCAAGACGAGAACACCAAGAAGG  
CATCCAGTATCTGGCGTACGTGGCGTCTCGCACAAAGAGCAAGAAGGACCAGGGCGAGCTGGAGCGGCAG  
CTGCTGCAGGCCAACCCCATCCTGGAGGCCTTCGGGAACGCCAAGACCGTGAAGAATGACAACCTCTCCC  
GCTTCGGCAAATTCATTGCGATCAACTTTGATGTCAATGGCTACATTGTTGGAGCCAACATTGAGACTTA  
30 TCTTTTGGAGAAATCTCGTGCTATCCGCCAAGCCAAGGAAGAACGGACCTTCCACATCTTCTATTATCTC  
CTGTCTGGGGCTGGAGAGCACCTGAAGACCGATCTCCTGTTGGAGCCGTACAACAAATACCGCTTCTCTGT  
CCAATGGACACGTCAACCATCCCCGGGCAGCAGGACAAGGACATGTTCCAGGAGACCATGGAGGCCATGAG  
GATTATGGGCATCCCAAGAAGAGGAGCAAAATGGGCCTGCTGCGGGTCATCTCAGGGGTCTTTCAGCTCGGC  
AATCATCGTCTTCAAGAAGGAGCGGAACATGACCAAGGCGTCCATCGCCGACAACACAGCTGCCCAAAGG  
35 TGTCCTCATCTCTTGGGTATCAATGTGACCTGATTTCCACGAGGAATCCCTCACCCCGCGCATCAAGGTGGG  
ACGGGATTACGTCCAGAAGGCGCAGACTAAAAGAGCAGGCTGACTTTGCCATCGAGGCCTTGGCCAAGGCG  
ACCTATGAGCGGATGTTCCGCTGGCTGGTGCTGCGCATCAACAAGGCTCTGGACAAGACCAAGAGGCAGG  
GCGCCTCCTTCATCGGGATCCTGGACATTGCCGGCTTCGAGATCTTTGATCTGAACTCGTTTGGAGCAGCT  
GTGCATCAATTACCAATGAGAAGCTGCAGCAGCTCTTCAACCACACCATGTTTCATCCTGGAGCAGGAG  
40 GAGTACCAGCGCAGGGGCATCGAGTGGAACCTTCATCGACTTTGGCCCTCGACCTGCAGCCCTGCATCGACC  
TCATTGAGAAGCCAGCAGGCCCCCGGGCATTCTGGCCCTGCTGGACGAGGAGTGCTGGTTCCCCAAAGC  
CACCGAAGAGACTTCGTGGAGAAGGTGATGCAGGAGCAGGCGACCCCAAGTTCCAGAAGGCCAAG  
CAGCTGAAGGACAAGCTGATTTCTGCATTATCCACTATGCGCGCAAGTGGATTACAAAGCTGACGAGT  
GGCTGATGAAGAACATGGATCCCCCTGAATGACAACATCGCCACACTGCTCCACCAGTCTCTGACAAGTT  
45 TGTCTCGGAGCTGTGGAAGGATGTGGACC GCATCATCGGCCTGGACCAGGTGGCCGGCATGTGCGGAGACC  
GCACTGCCCCGGGGCCTTCAAGACGCGGAAGGGCATGTTCCGCACTGTGGGGCAGCTTTACAAGGAGCAGC  
TGGCCAAGCTGATGGCTACGCTGAGGAACACGAACCCCAACTTTGTCCGCTGCATCATCCCCAACACGA  
GAAGAAGGCCGGCAAGCTGGACCCGCATCTCGTGCTGGACCAGCTGCGCTGCAACGGTGTTCTCGAGGGC  
ATCCGTATCTGCCGCCAGGGCTTCCCCAACAGGGTGGTCTTCCAGGAGTTTCCGCAGAGATATGAGATCC  
50 TGACTCCAACTCCATTCCCAAGGGTTTCATGGACGGGAAGCAGGCGTGCGTGCATGATAAAGCCCT  
GGAGCTCGACAGCAATCTGTACCGCATTTGCCACAGACAAATCTTCTCCGTGCCGTGTGCTGGCCAC  
CTGGAGGAGGAGCGAGACTGAAGATACCGACGTCATCAGATAGGTTTCCAGGCCTGCTGCAGGGGCTACC  
TGGCCAGGAAAGCATTTGCCAAGCGGCAGCAGCAGCTTACCGCCATGAAGGTCCTCCAGCGGAACCTGCGC  
TGCCTACCTGAAGCTGCGGAACCTGGCAGTGGTGGCGGCTCTTCACCAAGGTCAAGCCGCTGCTGCAGGTG  
55 AGCCGGCAGGAGGAGGAGATGATGGCCAAGGAGGAGGAGCTGGTGAAGGTGAGAGAGAAGCAGCTGGCTG  
CGGAGAACAGGCTCACGGAGATGGAGACGCTGCAGTCTCAGCTCATGGCAGAGAAATTGCAGCTGCAGGA  
GCAGCTCCAGGCAGAAACCGAGCTGTGTGCCGAGGCTGAGGAGCTCCGGGCCCGCTGACCGCCAAGAAG  
CAGGAATTAGAAGAGATCTGCCATGACCTAGAGGCCAGGGTGGAGGAGGAGGAGCGCTGCCAGCACCT  
TGCAGGCGGAGAAGAAGAAGATGCAGACAGAACCATCAGGAGCTTGAGGAGCAGCTGGAGGAGGAGGAGAG  
60 CGCCCGCGAGAAGCTGCAGCTGGAGAAGGTGACCACCGAGGCGAAGCTGAAAAAGCTGGAGGAGGAGCAG  
ATCATCTCTGGAGGAGCAGAACTGCAAGCTGGCCAAGGAAAAAGAACTGCTGGAAGACAGAATAGCTGAGT

TCACCACCAACCTCACAGAAGAGGAGGAGAAAATCTAAGAGCCTCGCCAAGCTCAAGAACAAGCATGAGGC  
 AATGATCACTGACTTGGAAGAGCGCCTCCGCAGGGAGGAGAAGCAGCGACAGGAGCTGGAGAAGACCCGC  
 CGGAAGCTGGAGGGAGACTCCACAGACCTCAGCGACCAGATCGCCGAGCTCCAGGCCAGATCGCGGAGC  
 TCAAGATGCAGCTGGCCAAGAAAGAGGAGGAGCTCCAGGCCGCCCTGGCCAGAGTGGAAGAGGAAGCTGC  
 5 CCAGAAGAATCATGGCCCTCAAGAAGATCCGGGAGCTGGAATCTCAGATCTCTGAACTCCAGGAAGACCTG  
 GAGTCTGAGCGTGCTTCCAGGAATAAAGCTGAGAAGCAGAAACGGGACCTTGGGGAAGAGCTAGAGGCTC  
 TGAAAACAGAGTTGGAGGACACGCTGGATTCCACAGCTGCCAGCAGGAGCTCAGGTCAAAACGTGAGCA  
 GGAGGTGAACATCCTGAAGAAGACCCTGGAGGAGGAGGCCAAGACCCACGAGGCCAGATCCAGGAGATG  
 AGGCAGAAGCACTCACAGGCCGTGGAGGAGCTGGCGGAGCAGCTGGAGCAGACGAAGCGGGTGAAAGCAA  
 10 ACCTCGAGAAGGCAAAGCAGACTCTGGAGAACGAGCGGGGGGAGCTGGCCAACGAGGTGAAGGTGCTGCT  
 GCAGGGCAAAGGGGACTCGGAGCACAAGCGCAAGAAAGTGGAGGCGCAGCTGCAGGAGCTGCAGGTCAAG  
 TTCAACGAGGGAGAGCGCGTGCGCACAGAGCTGGCCGACAAGGTACCAAGCTGCAGGTGGAGCTGGACA  
 ACGTGACCGGGCTTCTCAGCCAGTCCGACAGCAAGTCCAGCAAGCTACCAAGGACTTCTCCGCGCTGGA  
 GTCCAGCTGCAGGACACTCAGGAGCTGCTGCAGGAGGAGAACCAGGAGCAAGCTGAGCCTGAGCACCAAG  
 15 CTCAAGCAGGTGGAGGACGAGAAGAATTCCTTCCGGGAGCAGCTGGAGGAGGAGGAGGAGGCCAAGCACA  
 ACCTGGAGAAGCAGATCGCCACCCTCCATGCCAGGTGGCCGACATGAAAAAGAAGATGGAGGACAGTGT  
 GGGGTGCCTGGAACTGCTGAGGAGGTGAAGAGGAAGCTCCAGAAGGACCTGGAGGGCCTGAGCCAGCGG  
 CACGAGGAGAAGGTGGCCGCCTACGACAAGCTGGAGAAGACCAAGACGCGGCTGCAGCAGGAGCTGGACG  
 ACCTGCTGGTGGACCTGGACCACCAGCGCCAGAGCGGTGCAACCTGGAGAAGAAGCAGAAGAAGTTTGA  
 20 CCAGCTCCTGGCGGAGGAGAAGACCATCTCTGCCAAGTATGCAGAGGAGCGCGACCGGCTGAGCGGAG  
 CCCCAGAGAAGGAGACCAAGGCTCTGTCTGCTGGCCCGGCGCCTGGAGGAAGCCATGGAGCAGAAGCGG  
 AGCTGGAGCGGCTCAACAAGCAGTTCCGCACGGAGATGGAGGACCTTATGAGCTCCAAGGATGATGTGGG  
 CAAGAGTGTCCACGAGCTGGAGAAGTCCAAGCGGGCCCTAGAGCAGCAGGTGGAGGAGATGAAGACGCAG  
 CTGGAAGAGCTGGAGGACGAGCTGCAGGCCACCGAAGATGCCAAGCTGCGGTTGGAGGTCAACCTGCAGG  
 25 CCATGAAGGCCAGTTTCGAGCGGGACCTGCAGGGCCGGGACGAGCAGAGCGAGGAGAAGAAGAAGCAGCT  
 GGTCAAGCAGGTGCGGGAGATGGAGGCAGAGCTGGAGGACGAGAGGAAGCAGCGCTCGATGGCAGTGGCC  
 GCCCGGAAGAAGCTGGAGATGGACCTGAAGGACCTGGAGGCGCACATCGACTCGGCCAACAAGAACCAGG  
 ACGAAGCCATCAAACAGCTGCGGAAGCTGCAGGCCCAGATGAAGGACTGCATGCGCGAGCTGGATGACAC  
 CCGCGCCTCTCGTGAGGAGATCCTGGCCAGGCCAAAGAGAACGAGAAGAAGCTGAAGAGCATGGAGGCC  
 30 GAGATGATCCAGTTGAGGAGGAGTGGCAGCCGCGAGCGTGCCTAAGCGCCAGGCGCAGGAGGAGCGG  
 ATGAGCTGGCTGACGAGATCGCCAACAGCAGCGGCAAGGAGCCCTGGCGTTAGAGGAGAAGCGGCGTCT  
 GGAGGCCCGCATCGCCAGCTGGAGGAGGAGCTGGAGGAGGAGCAGGGCAACACGGAGCTGATCAACGAC  
 CGGCTGAAGAAGGCCAACCTGCAGATCGACCAGATCAACACCGACCTGAACCTGGAGCGCAGCCACGCCC  
 AGAAGAACGAGAATGCTCGGCAGCAGCTGGAACGCCAGAACAAGGAGCTTAAGGTCAAGCTGCAGGAGAT  
 35 GGAGGGCAGTGTCAAGTCCAAGTACAAGGCCTCCATCACCGCCCTCGAGGCCAAGATTGCACAGCTGGAG  
 GAGCAGCTGGACAACGAGACCAAGGAGCGCCAGGCAGCCTGCAAACAGGTGCGTTCGGACCGAGAAGAAGC  
 TGAAGGATGTGCTGCTGCAGGTGGATGACGAGCGGAGGAACGCCGAGCAGTACAAGGACCAGGCCGACAA  
 GGCATCTACCCGCTGAAGCAGCTCAAGCGGCAGCTGGAGGAGGCCGAAGAGGAGGCCAGCGGCCCAAC  
 GCCTCCCGCCGGAAGACTGCAGCGGAGCTGGAGGAGTGCCTGAGACGCGCCGATGCCATGAACCGCGAAG  
 40 TCAGCTCCCTAAAGAACAAGCTCAGGCGCGGGGACCTGCCGTTTGTGCTGCCCCGCCGAATGGCCCCGAA  
 AGGCGCCGGGGATGGCTCCGACGAAGAGGTAGATGGCAAAGCGGATGGGGCTGAGGCCAAACCTGCCGAA  
 TAAGCCTCTTCTCCTGCAGCCTGAGATGGATGGACAGACAGACACCACAGCCTCCCTTCCAGACCCCG  
 CAGCACGCCTCTCCCCACCTTCTTGGGACTGCTGTGAACATGCCTCCTCCTGCCCTCCGCCCCGTCCCC  
 CATCCCGTTTCCCTCCAGGTGTTGTTGAGGGCATTGCGTTCCTCTGCTGCATCCCTTCCAGCTCCCTC  
 45 CCTGCTCAGAATCTGATACCAAAGAGACAGGGCCCCGGGCCAGGCAGAGAGCGACCAGCAGGCTCCTCA  
 GCCCTCTCTTGCCAAAAGACACAAGATGTTGAGGCGAGCAGGGCAGGCCCGGGGAGGGGCCAGAGTTT  
 TCTATGAATCTATTTTTCTTCAGACTGAGGCCTTTTGGTAGTCGGAGCCCCCGCAGTCGTACGCTCCCT  
 GACGTCTGCCACCAGCGCCCCCACTCCTCCTCTTCTTGTGTTTGAATCACACGTGGTGACCTCAC  
 ACACCTCTGCCCTTGGGCCTCCCACTCCCATGGCTCTGGGCGGTCCAGAAGGAGCAGGCCCTGGGCCTC  
 50 CACCTCTGTGCAGGGCACAGAAGGCTGGGGTGGGGGGAGGAGTGGATTCTCCCCACCTGTCCCAGGCA  
 GCGCCACTGTCCGCTGTCTCCCTCCTGATTCTAAAATGTCTCAAGTGCAATGCCCCCTCCCTCCTTTAC  
 CGAGGACAGCCTGCCTCTGCCACAGCAAGGCTGTGCGGGTCAAGCTGGAAAGGCCAGCAGCCTTCCAGTG  
 GCTTCTCCCAACACTCTTGGGGACCAATATATTTAATGGTTAAGGGACTTGTCCCAAGTCTGACAGCCA  
 GAGCGTTAGAGGGGCCAGCGGCCCTCCAGGCAGATCTTGTGTCTACTCTAGGACTGGGCCCCGAGGGTGGT  
 55 TTACCTGCACCGTTGACTCAGTATAGTTTAAAAATCTGCCACCTGCACAGGTATTTTTGAAAGCAAAATA  
 AGGTTTTCTTTTTTCCCTTTCTTGTATAAATGATAAAATCCGAGTCTTTCTCAGTCCCTTGTTTAG  
 AAGAGAGTAGCTCGTCTCCTCAGTCTGAGGCTGCGCAATTTACTTGTATTCTTAATCTGTTTGTAT  
 ATGCTGCATTGAGACTTACGGCAAGAAGGCATTTTTTTTTTTTTTAAAGGAAACAAACTCTCAAATCATGAA  
 GTGATATAAAAGCTGCATATGCCTACAAAGCTCTGAATTCAGGTCCCAGTTGCTGTACAAAGGAGTGAG  
 60 TGAAACTCCCACCTACCCCTTTTTTATATAATAAAAGTGCCTTAGCATGTGTTGCAGCTGTCAACACT  
 ACAGTAAGCTGGTTTACAGATGTTTTCCACTGAGCATCACAATAAAGAGAACCATGTGCTACGA

>gi|12667788|ref|NP\_002464.1| myosin, heavy polypeptide 9, non-muscle  
[Homo sapiens]

5 MAQQAADKYLYVDKNFINNPLAQADWAAKLVWVPSDKSGFEPASLKEEVGEEAIVELVENGKKVKVKNKD  
DIQKMNPPKFSKVEDMAELTCLNEASVLHNLKERYYSGLIYTYSGLFVVINPYKNLPYSEEIVEMYKG  
KKRHEMPPHIYAITDTAYRSMMQDREDQSILCTGESGAGKTENTKKVIQYLAYVASSHKSCKDQGELEERQ  
LLQANPILEAFGNAKTVKNDNSSRFGKFIRINFDVNGYIVGANIETYLLEKSRAIRQAKEERTFHFIFYL  
10 LSGAGEHLKTDLLLEPYNKYRFLSNGHVTIPGQQDKDMFQETMEAMRIMGIPEEEQMGLLRVISGVLQLG  
NIVFKKERNTDQASMPDNTAAQKVSHLLGINVTDFTRGILTPIKVGDRDYVQKAQTKEQADFAIEALAKA  
TYERMFRWLVLRLINKALDKTKRQASFIGILDIAGFEIFDLNSFEQLCINYTNEKLQQLFNHTMFILEQE  
EYQREGIEWNFIDFGLDLQPCIDLIEKPAGPPGILALLDEECWFPKATDKSFVEKVMQEQGTHPKFQKPK  
QLKDKADFCIIHYAGKVDYKADEWLMKNMDDLNDNIATLLHQSSDKFVSELWKDVDRIIIGLDQVAGMSET  
ALPGAFKTRKGMFRVTGQLYKEQLAKLMATLRNTNPNFVRCIIPNHEKKAGKLDPHLVLDQLRCNGVLEG  
15 IRICRQGFPNRVVFQEFRQRYEILTPNSIPKGFMDGKQACVLMIKALELDSNLYRIGQSKVFFRAGVLAH  
LEEERDLKITDVIIGFQACCRGYLARKAFAKRQQQLTAMKVLQRNCAAYLKLNRWQWWRLFTKVKPLLQV  
SRQEEEMMAKEEEELVKVREKQLAAENRLTEMETLQSQLMAEKLQLQEQLQAETELCAEAEELRARLTAKK  
QELEEICHDLARVEEEEEERCQHLQAEKKMQQNIQELEEQLEEEEESARQKLQLEKVTTEAKLKKLEEEQ  
IILEDQNCCLAKEKKLLEDRIAEFTTNLTETEEESKSLAKLKNKHEAMITDLEERLRREEKQRQELEKTR  
RKLEGDSTDLSQIAELQAIQIAELKMQLAKEEELQAALARVEEEAAQKNMALKKIRELESQISELQEDL  
20 ESERASRNKAQKQKRDLEGELEALKTELEDLTLDSTAAQQLRSKREQEVNILKKTLEEEAKTHEAQIQEM  
RQKHSQAVEELAEQLEQTKRVKANLEKAKQTLENERGELANEVKKVLLQKGKGDSEHKRKKVEAQLEQVQK  
FNEGERVRELADKVTKLQVELDNVTGLLSQSDSKSSKLTDFSALESQLODQELLQEEENRQKLSLSTK  
LKQVEDEKNSFREQLEEEEEAKHNLEKQIATLHAQVADMKKMEDSVGCLETAEEVKKRKLQKDLGLSQR  
HEEKVAAYDKLEKTKTRLQQLDDLLVDLDHQRQSACNLEKKQKKFDQLLAEKTIISAKYAEERDRAEAE  
25 AREKETKALSLARALEEAMEQKAELERLNNKQFRTEMEDLMSSKDDVGKSVHELEKSKRALEQQVEEMKTQ  
LEELEDELQATEDAKLRLEVNLQAMKAQFERDLQGRDEQSEEKKKQLVRQVREMAELEDERKQQRSMVA  
ARKKLEMDLKDLEAHIDSANKNRDEAIKQLRLKLAQMKDCMRELDLTRASREEILAQAKENEKKLKSMEA  
EMIQLQEELAAAEAKRQAQQRDELADEIANSSGKGALALEEKKRLEARIAQLEEELEEEQGNTELIND  
RLKKANLQIDQINTDLNLSHAQKNENARQQLERQNKELVKVLQEMEGTVKSKYKASITALEAKIAQLE  
30 EQLDNETKERQAAACKQVRRTEKKKLKDVLLQVDDERRNAEQYKDQADKASTRLKQLRKQLEEAEEEAQRAN  
ASRRKLQRELEDATETADAMNREVSSLKNKLRRGDLFPVVPRRMARKGAGDGSDEEVDGKADGAEAKPAE

Calmodulin 2

35 GH1-84-PCR-G3F1  
GCTGTCTGTAAATACCTGGTGCTAACATCCCATGCCGCTCCCTCCTCACGATGCACCCACCGCCCTGAGGGC  
CCGTCCCTAGGAATGGATGTGGGGATGGTCGCTTTGTAATGTGCTGGTTCTCTTTTTTTTTCTTTCCCTCTT  
TGGCCCTTAAGACTTTTCATTTTGTTCAGAACCATGCTGGGCTAGCTAAAGGGTGGGGAGAGGGAAGATGGGC  
40 CCCACCACGCTCTCAAGAGAACGCACCTGCAATAAAACAGTCTTGTGCGCCAGCTGCCCAGGGGACGGCAG  
>gi|13477324|gb|BC005137.1|BC005137 Homo sapiens, calmodulin 2  
(phosphorylase kinase, delta), clone MGC:1447 IMAGE:3504793, mRNA,  
complete cds  
GGCACGAGGGCGCGGGCGGAGCTGGAAGTCTGTCAGCTGCTGCCGCCGCCGGAGGAACCTTGATCCCCGTG  
45 CTCCGGACACCCCGGGCCTCGCCATGGCTGACCAGCTGACTGAGGAGCAGATTGCAGAGTTCAAGGAGGC  
CTTCTCCCTCTTTGACAAGGATGGAGATGGCACTATCACCACCAAGGAGTTGGGGACAGTGATGAGATCC  
CTGGGACAGAACCCCACTGAAGCAGAGCTGCAGGATATGATCAATGAGGTGGATGCAGATGGGAACGGGA  
CCATTGACTTCCCGGAGTTCTTGACCATGATGGCCAGAAAGATGAAGGACACAGACAGATGAGGAGGAGAT  
CCGAGAGGCGTTCCGTGTCTTTGACAAGGATGGGAATGGCTACATCAGCGCCGAGAGCTGCGTCACGTA  
50 ATGACGAACCTGGGGGAGAAGCTGACCGATGAGGAGGTGGATGAGATGATCAGGGAGGCTGACATCGATG  
GAGATGGCCAGGTCAATTATGAAGAGTTTGTACAGATGATGACTGCAAAGTGAAGGCCCCCGGGCAGCT  
GGCGATGCCCGTTCTCTTGATCTCTCTTCTCGCGCGCGCACTCTCTTCAACACTCCCTTGCGTACC  
CCGTTCTAGCAAACACCAATTGATTGACTGAGAATCTGATAAAGCAACAAAGATTTGTCCCAAGCTGC  
ATGATTGCTCTTTCTCCTTCTTCCCTGAGTCTCTCTCCATGCCCCCTCATCTCTTCTTTTGCCCTCGCCT  
55 CTTCATCCATGTCTTCCAAGGCCTGATGCATTATAGTTGAAGCCCTCCCAGATCCCTTGAGGGAGC  
CTCTGCCCTCCTCCAGCCCGATGGCTCTCTCCATTTTGGTTTGTCTCTCTGTTGTCATCTTATTT  
TGGTGCTGGGGTGGCTGCCAGCCTGTCCCGGACCTGCTGGGAGGGACAAGAGGCCCTCCCCAGGCA  
GAAGAGCATGCCCTTTGCCGTTGCATGCAACCAGCCCTGTGATTCCACGTGCAGATCCCAGCAGCCTGTT  
GGGGCAGGGGTGCCAAGAGAGGCATTCCAGAAGGACTGAGGGGGCGTTGAGGAATTGTGGCGTTGACTGG  
60 ATGTGGCCAGGAGGGGTGCGAGGGGGCCAACTCACAGAAGGGGACTGACAGTGGGCAACACTCACATCC  
CACTGGCTGCTGTTCTGAAACCATCTGATTGGCTTTCTGAGTTTGGCTGGGTGGGGACTGCTCATTGG

- CCACTCTGCAGATTGGACTTGGCCGCGTTCCTGAAGCGCTCTCGAGCTGTTCTGTAAATACCTGGTGCTA  
ACATCCCATGCCGCTCCCTCCTCACGATGCACCCACCGCCCTGAGGGCCCGTCTAGGAATGGATGTGGG  
GATGGTCGCTTTGTAAATGTGCTGGTTCTCTTTTTTTTTTTCTTTCCCTCTATGGCCCTTAAGACTTTCATT  
TGTTCAGAACCATGCTGGGCTAGCTAAAGGGTGGGGAGAGGAAGATGGGCCCCACCACGCTCTCAAGA  
5 GAACGCACCTGCAATAAAACAGTCTTGTCGGCCAGCTGCCAGGGGACGGCAGCTACAGCAGCCTCTGCG  
TCCTGGTCCGCCAGCACCTCCCGCTTCTCCGTGGTGACTTGGCGCCGCTTCCCTCACATCTGTGCTCCGTG  
CCCTCTTCCCTGCCTCTTCCCTCGCCACCTGCCTGCCCCATACTCCCCCAGCGGAGAGCATGATCCGT  
GCCCTTGCTTCTGACTTTTCGCTCTGGGACAAGTAAGTCAATGTGGGCAGTTTCACTCGTCTGGGTTTTTT  
CCCCTTTTCTGTTTCAATTCATCTGGCTCCCCCACCACCTCCCCACCCACCCCTGCTTCCC  
10 CTCAGTGCCAGGTCGATCAAGTGGCTTTTCTGGGACCTGCCAGCTTTGAGAATCTCTTCTCATCCAC  
CCTCTGGCACCAGCCTCTGAGGGAAGGAGGGATGGGGCATAGTGGGAGACCCAGCCAAGAGCTGAGGGT  
AAGGTCAAGGTAGGCGTGAGGCTGTGGACATTTTCGGAATGTTTTGGTTTTTTTTTAAACCGGGCAA  
TATTGTGTTCAAGCTGTGAAGAAAAATATATATCAATGTTTTCCAATAAAATACAGTGACTACC'T  
GAAAAAAAAAAAAAAAAAAAAA
- 15 >gi|13477325|gb|AAH05137.1|AAH05137 calmodulin 2 (phosphorylase kinase,  
delta) [Homo sapiens]  
MADQLTEEQIAEFKEAFSLFDKDGDTITTKELGTVMRSLGQNPTAELODMINEVDADGNGTIDFPEFL  
TMMARKMKD TDSEEEIREAFRVFDKDGNGYISAAELRHVMTNLGEKLTDEEVDEMIREADIDGDGQVNYE  
20 EFVQMMTAK
- Novel Symporter
- GH1-178-PCR-G3F1  
25 CTGGGTTCCTTGCAGACTTGGCTGGAGATCACGATGATGCCCTCACTGTCCTCAGTGAAACTCAAACTCCA  
TCACAGAGCCATCTCCAATGCTCAAGTAGCGGCCCTTCCCTGCCAGGCCCGGCCGGGCGACCCGAGTGGGCG  
ATCGCGGAGCAGGTGCGGGCCAGAGGCCGCTCCCTTCCGGAGGCTCTCACCTGCCACAGCCACCGCTGCAC  
CGCAGGAACCCAGCACAGTGGTTAGATTGATAAGCGGCCGCTCGACTAGTCTGAGGTCTGATACTCACTGAC  
TGTCGTAT
- 30 Novel Semaphorin
- GH1-204-PCR-G3F1  
35 AAAAAATCTACTTCTAAGCTTGTTCTTATTGTTGGCAGAATTCAGGTCCTTGTGGCTGTAGGACCGAGGCC  
CAGCTTCCTGCTGATGTCCGCTGGAGACTGCTGTGCTCAGCTCCCAGAGGCCACCCCATTCCTGGACACGTGGC  
CCCTCCATCTCAAACCTGCAGTGGGTGTTAAACCCCTTCTCATGCTTCTCATCTCTACTTCAGGAATACAGAT  
AGTGTCTGGTGGCTTGACGTGATTTTAATGAATTTGGACTCCATGTGGATTGTTGTCGTCTCCCTATTCCGAG  
CTGCGGGCAGGGAGAGGGCCCTCGCGCCGCCCTCAGCAGCCGGCGGGCGGAGGTAGACGAGCGGGGACGG  
AAGGACAGACCGACGTGCGCAGCTGGAATCATGTGAGGGCAACCGGGGAAGGTGGAGCAGATGAGCACACA  
40 CAGGAGCCGTCTCCTCACCGCCGCCCTCTCAGCATGGAACAGAGGCGCCCTGGCCCCGGGCCCTGGAGGTG  
GACAGCCGCTCTGTGGTCCTGCT
- Novel Zn finger helicase
- GH1-31-PCR-G3F1  
45 GAAGGAGAAGATGGTATAAACTGGTCCATCAGTGACAAAGACATTGAGGCCAGATAGCTAATAACCGAACA  
CCTGGAAGATGGACCCAGCGGTACTATTACGACAAAAACATTTCTGTGAAATTGTGACAAACGTGGTCA  
TTTATCAAAAACTGCCCTTACCACGAAAAGTTCGTGCTGCTTCTGTGCTCCAGGAGAGGACATCTCCT  
GTATTCTGTTCAGCCCCCTTTGCGAATACTGTCTGTGCCTAATGTTTGACCACTCATGTCTTTTCAGAC  
50 ATTCCTGGATAAACAGTGTGACCGATGTATATTGCTAGGCACTATACAGATTCTTGCCAGAAATCTTGAGG  
CAGTTATCACCTTACGACAACTTGACACCCCAAAGCCGAAAACCTTTCCGCAAAAACCCGCACAGTGGTT  
TGATTGATTAAGGCGGCGCTCGACTAGTCTGAGGTCTGATACTCACTGAC
- Novel Sugar transporter
- GH1-175-PCR-G3F1  
55 ACCGAGCCAGAGAATGTACCAATGGCACAGTGGGCGGCACAGCAGAGCCGGGGCACGAGGAGGTGAGCTGG  
ATGAACGGCTGGCTCAGCTGCCAGGCCAGGACGAGATGCTAAATTTGGCCTTCACTGTGGGCTCCTTTCTG  
CTCAGTGCCATCACCTGCCCCTGGGTATCGTCATGGACAAGTATGGCCCGAGGAAGCTCAGGCTGCTGGGC  
60 AGCGCTGCTTCGCGGTTTTCTGCTTGCTGATTGCGT

## human Plexin-A2

	gctgccggga	ggagcggcat	ccgcgccaga	ctggagcggg	agggcggcgg	agggcagttg
	ctgggaat	ttcagccgag	agggcgagcg	atccggagag	agaccccgag	agcttgggag
5	cggtagggcg	tgcgagcgcc	gcagccagcg	gagcaaacct	cgaaatagat	ctggaaagcc
	aggctcccgg	aggaaatggg	actgtgaacg	aaccggagag	caagaaggga	aggaagcgcc
	gggattgctg	atgtcagagg	agcccggaaa	gtcgcgctgg	aaaaatctga	agacagccgg
	ggctctgctt	cttcctcagg	agagacaccg	ccggccgccc	ccacacgccc	cctcggcgcc
	tccgggtgcc	ccctgagagc	cggcgacagc	gcccagccgg	gctgctgcgg	ggcgacggag
10	gactgagggg	cgcgccgagc	ggagaccgag	gagcgacttc	aggaatacag	ataagtgtct
	ggtggcttga	cgtggat	aatgaat	gactccatgt	ggatttggtc	gtctccctga
	ttccgagctg	cgggcaggga	gaggggcctc	gcgcgcctct	cagcagccgg	cggcgccgga
	ggtagaccga	gcggggacgg	aaggacagac	cgacgtcgcc	gagctggaat	catgtgaggg
	ccaaccgggg	aaggtggagc	agatgagcac	acacaggagc	cgtctcctca	ccgcgcctcc
15	tctcagcatg	gaacagaggc	ggccctggcc	ccgggcccctg	gaggtggaca	gccgctctgt
	ggtcctgctc	tcagtgggtc	gggtgctgct	ggccccccca	gcagccggca	tgccctcagtt
	cagcaccttc	cactctgaga	atcgtgactg	gaccttcaac	cacttgaccg	tccaccaagg
	gacggggggc	gtctatgtgg	gggccatcaa	ccgggtctat	aagctgacag	gcaacctgac
	catccagggtg	gtccataaga	cagggccaga	agaggacaac	aagtcttggt	acccgcccct
20	catcgtgcag	ccctgcagcg	aagtgtcac	cctcaccaac	aatgtcaaca	agctgctcat
	cattgactac	tctgagaacc	gctgtctggc	ctgtgggagc	ctctaccagg	gggtctgcaa
	gctgtctcgg	ctggatgacc	tcttcatcct	ggtggagcca	tcccacaaga	aggagcacta
	cctgtccagt	gtcaacaaga	cgggcacccat	gtacgggggtg	attgtgcgct	ctgaggggtga
	ggatggcaag	ctcttcatcg	gcacggctgt	ggatgggaag	caggattact	tcccgaccct
25	gtccagccgg	aagctgcccc	gagaccctga	gtcctcagcc	atgctcgact	atgagctaca
	cagcgat	gtctcctctc	tcatcaagat	cccttcagac	accctggccc	tgggtctccca
	ctttgacatc	ttctacatct	acggctttgc	tagtgggggc	tttgtctact	ttctcactgt
	ccagcccag	acccctgagg	gtgtggccat	caactccgct	ggagacctct	tctacacctc
	acgcctcgtg	cggctctgca	aggatgacc	caagtccac	tcatacgtgt	ccctgccctt
30	cggctgcacc	cgggcggggg	tggatcaccg	cctcctgcag	gctgcttacc	ggcccaagcc
	tggggactca	ctggcccagg	ccttcaatat	caccagccag	gacgatgtac	tctttgccat
	cttctccaaa	gggcagaagc	agtatcacca	cccgcgccgat	gactctgccc	tgtgtgcctt
	ccctatccgg	gccatcaact	tgcagatcaa	ggagcgccctg	cagtctctgt	accagggcga
	gggcaacctg	gagctcaact	ggctgctggg	gaaggacgtc	cagtgcacca	aggcgccctgt
35	ccccatcgat	gataacttct	gtggactgga	catcaaccag	cccctgggag	gctcaactcc
	agtggagggc	ctgaccctgt	acaccaccag	cagggaccgc	atgacctctg	tggcctccta
	cgtttacaac	ggctacagcg	tggtttttgt	ggggactaag	agtggcaagc	tgaaaaagat
	tcggggccgac	ggtccccccc	atggtggggg	ccagtacgag	atggtctctg	tgtcaagga
	cggaaagccc	atcctccggg	acatggccct	ctccattgat	cagcgctacc	tgtaacgtcat
40	gtctgagaga	caggtcacca	gggtccccgt	ggagtcatgt	gagcagtata	cgacttgtgg
	ggagtgcctg	agctctgggg	accctcactg	tggctgggtgt	gccctgcaca	acatgtgtct
	ccgcagggac	aaatgccaac	aggcctggga	acctaactga	tttgctgcca	gcatcagcca
	gtgtgtgagc	cttgacgtgc	atcccagcag	catctcagta	tctgagcaca	gccgggtgct
	tagcctggta	gtgagtgatg	ctcctgatct	atctgcgggt	atcgccctgtg	cctttgggaa
45	cctgacagag	gtggaggggg	aggtgtccgg	gagccaggtc	atctgcatct	cacctggggc
	caaggatgtc	cctgtcatcc	cgctggatca	agactggttt	gggctggagc	tacagctgag
	gtccaaggag	acaggggaaga	tatttgtcag	caccgagttc	aagttttaca	actgcagtgc
	ccaccaactg	tgctgtctct	gtgtcaacag	cgccttccgc	tgccattgggt	gcaagtaccg
	caacctctgc	actcatgacc	ccaccacctg	ctccttccag	gagggccgga	tcaatat
50	agaggactgt	ccccagctgg	tgccacacaga	ggagatcttg	attccagtcg	gggaggtaaa
	gccaatcacc	cttaaggcgc	gaaatctgcc	ccagccgcag	tccggccagc	gaggctatga
	gtgtgtcctc	aacatacaag	gagccatcca	ccgggtcccc	gctctgcgct	tcaacagctc
	cagcgttcag	tgtcagaaca	gctcgtacca	gtatgatggc	atggacatca	gcaatctggc
	cgtggatttc	gctgtgggtgt	ggaacggcaa	tttcatcatt	gacaaccctc	aggacctgaa
55	agtccatctc	tacaagtgtg	cagcccagcg	ggagagctgc	ggcctctgcc	tcaaggccga
	ccggaagt	gagtgtggct	ggtgcagcgg	cgagcgcagg	tgacacctcc	accagactg
	taccagccct	tccagccct	ggctcgactg	gtccagccac	aatgtcaagt	gctccaacct
	tcaaatcacc	gagat	cgggtgctgg	accgccggaa	ggagggacgc	gagtgacat
	ccatggcgtg	aacctgggtc	tggacttctc	cgagatcgcc	caccatgtgc	aggtggctgg
60	ggtgccctgc	acgcccctcc	caggggaata	catcatcgct	gagcagattg	tctgtgagat
	gggccatgcc	ctcgtgggaa	ccacctccgg	gccagtacgc	ctgtgtattg	gcgagtgtaa

	gccagagttc	atgacgaagt	cccatcagca	gtacaccttc	gtgaaccctt	ctgtgctgtc
	actcaaccca	atccgaggtc	ccgagtcagg	aggcactatg	gtgaccatta	ccggccatta
	ccttggggct	gggagcagcg	tggcagtcta	cctgggcaac	cagacctgcg	agttctacgg
5	gaggtcaatg	agtgaatcg	tgtgtgtctc	acccccatca	tccaatggcc	ttggcccggt
	ccctgtttct	gtgagtgtcg	accgagccca	tgtggatagc	aacctgcagt	ttgagtacat
	agatgaccct	cggttccagc	gcacgcagcc	agagtggagc	attgccagtg	gccacacacc
	cctgaccatc	acaggcttca	acctggatgt	cattcaggag	ccaaggatcc	gagtcaaatt
	caatggcaaa	gaatctgtca	atgtgtgtaa	agttgtgaac	acaaccaccc	tcacctgcct
10	ggcacccctc	ctgaccacgg	actaccgccc	tggcctggac	actgtggaac	gcccagatga
	gtttggattt	gtctttaaca	atgtccaatc	cttgctaatt	tacaacgaca	ccaagtatat
	ctactacccc	aacctgacct	ttgaactgct	tagccctact	ggagtcttgg	atcaaaagcc
	aggatcgccc	atcattctga	agggcaaaaa	cctctgccct	cctgcctctg	gagggggcaa
	actcaactac	actgtgtctc	tgggagagac	cccttgtgct	gtcaccgat	ctgagacca
	gcttctctgc	gagcctccca	acctcaccgg	gcagcacaag	gtcatggttc	acgtgggcgg
15	gatgggtgtc	tgccttggct	cggtgagtg	catctcagac	agcttgtctg	ccctgccagc
	catcgtcagc	atcgccggcg	gcggcagcct	cctcctcatc	atcgtcatca	tgcctcctcat
	tgcctacaag	cgcaagtctc	gagaaaatga	cctcactctc	aagcggctgc	aaatgcagat
	ggacaatctg	gagtcocgtg	tggccttgg	gtgcaaggaa	gcttttgcct	agctccagac
	ggatatcaat	gagttgacca	gtgacctgga	ccgctcagga	atcccttacc	tggagctatc
20	tacctacgct	atgcgagtc	tgttcccggg	catcgaggac	caccccgctc	tcggggatcg
	ggaggtacaa	ggaaacgggc	agcagcacgt	ggagaaggcc	ctgaagctct	ttggccagct
	catcaacaac	aaggtgttcc	tgtgtacctt	catccgcacc	ctggagctgc	agcgcagttt
	ctccatgcgc	gaccggggca	acgtggcttc	gctcatcatg	accggcctgc	agggccgcct
	ggaatatgcc	actgatgtcc	tcaagcagct	gctctctgac	ctcatcgata	agaacctgga
25	gaacaagaac	cacccaagc	tgtactccg	gaggacagag	tctgtggctg	aaaagatgct
	gaccaattgg	ttcgccctcc	tcctgcacaa	gttcctaaag	gagtgcgcag	gggagccact
	cttcatgcta	tactgtgcca	tcaagcagca	gatggagaag	ggccccattg	atgccatcac
	gggagaggcc	cgctactccc	tgagcgagga	caagctcatc	cggcagcaga	tcgagtacaa
	gaccttgatc	ctgaactgcg	tcaaccttga	caacgagaac	agtccagaga	tcccagtga
30	ggtgtttaa	tgtgacacca	tcacacaggt	caaggagaag	attcttgatg	ccgtgtataa
	gaatgtgccc	tattcccagc	ggccgagggc	agtggacatg	gacttggagt	ggcgccaagg
	ccggatcgcc	cggttcgtgc	tgcaagatga	ggacatcacc	accaagattg	agggtgactg
	gaagcggctc	aacacactga	tgcattatca	ggtgtcagac	aggtcggtgg	tggctctggt
	ccccaaacag	acctcctcct	acaacatccc	tgcctctgcc	agcatctccc	ggacgtccat
35	cagcagatac	ggtgactcct	ccttcaggta	tacgggcagc	cccagacagc	tgcgggtccc
	ggccccgatg	atcaccccag	acctggaaa	tggggccaag	gtgtggcatc	tgggtgaagaa
	ccatgaccac	ggtgaccaga	aggaggggtg	ccggggcagc	aagatgggtg	ccgagatcta
	cctgaccggg	ctactggcca	ccaagggcac	cctgcagaag	tttgtggagc	acttggttga
	gaccttggtc	agcactgtgc	accggggcag	cgctctcccc	ctggccatca	agtacatggt
40	tgatttccta	gatgagcagg	cagacaggca	cagcatccat	gacacagatg	tgcggcacac
	ctggaaaagc	aactgcctcc	ctctgcgctt	ctgggtgaac	gtgattaaga	acccccagtt
	cgtgtttgac	atccacaagg	gcagcatcac	ggacgcctgc	ctctctgtgg	tggcccagac
	cttcatggac	tcttgttcaa	cgtcagagca	ccggctgggc	aaggactccc	cctccaacaa
	gctgctctat	gccaaaggaca	tccccagcta	caagagctgg	gtggagagat	actacgcaga
45	catcgccaag	ctcccagcca	tcagtgacca	ggacatgaat	gcctacctcg	ccgagcagtc
	ccgcttcac	gccgtggagt	tcaacatgct	gagtgcctc	aatgagatct	actcctatgt
	cagcaagtat	agtgaggagc	tcacggggc	cctagagcag	gatgagcagg	cacggcggca
	gcggctggct	tataagggtg	agcagctcat	taatgccatg	tccattgaga	gctgagagga
	ggagcctcgc	attcctggga	agagggacct	gtccaagctg	tcacactggg	agttctcagat
50	ggaaggacaa	gtgatgggga	tcaggcccca	gagcttgctg	tcccctgaga	ccccatcctg
	gggagagggg	aggactcctc	tccctacgcc	agccaagttt	cgtcatagcc	agttccagct
	gggagagaca	gtgggcgtcg	tccatcctca	gtgagaacac	cagagaaccc	ggggccggga
	gaaggtggtt	cttcaagccg	agaggcacga	gctggggaca	gttctgcctc	tgtgactgct
	gctttgcatg	aaaactcatt	tgatgtatat	tggggaaata	atgagaactt	tatttaattt
55	ttttaagaaa	aagggaaaaa	aacagaaata	aaacaaaaag	ccgccctgtt	aatcccgtcc
	aacttttgtt	taattctgat	ttctgtctcc	cttccatctt	ttctcccatt	cctccttctt
	tatataatgc	ctatttccaa	atgccagaga	aagcagagat	gctgagagac	attggagaga
	aatgactgt	ctccttttcc	ttgaaattaa	aaaaaaaaaa	aaaaagagaa	agaggagaag
	aagaatgatg	agcacaagta	tgcaccaaac	acttcgcaaa	aacagaggcc	agtaaaacct
60	ggaattatcc	cggcagccag	aggagtatgg	aacttcaga	actttgcaca	aattgcaaag
	ccatcaagag	ctcaccctgg	ctgactggaa	actgagcttt	atctaccaca	cacctgtata

ttctcatctt ttgagaggag atgtgtacct agatagtacc aatgcttttt gctactgttt  
 tttgttttgt tttatttaaat cctaaacctc aacaaatgag gagctgggtct ttgatagtgt  
 tcctttcaat ttccctaaag ttactatgag aagtggggtg aggtgggcct ctcccagacc  
 agacacctgg cagccctgcc tcatatcaat ccctgtcata aaccaggcac cctggggaaa  
 5 cggcctggag gtgtgtgggc caggcctcca cgaggttcca tttgaaagtt gatttggaga  
 cataggtgtt tgactttgga gttcactcca atcatccagt ggtccctggc aatt

MEQRRPWPRALEVDSRSVLLSVVWVLLAPPAAGMPQFSTFHSENRDWTFNHLTVHQGTGAVYVGAINRVYK  
 10 LTGNLTIQVAHKTGPEEDNKSCYPPLIVQPCSEVLTLTNNVNKLLIIDYSENRLACGSLYQGVCCKLLRLDD  
 LFILVEPSHKKEHYLSSVNKTGTMYGVIVRSEGEDGKLFITAVDGKQDYFPTLSSRKLPRDPRESSAMLDYE  
 LHSDFVSSLIKIPSDTLALVSHFDIFYIYGFASSGFVYFLTVQPETPEGVAINSAGDLFYTSRIVRLCKDDP  
 KFHSYVSLPFGCTRAGVEYRLLQAAYLAKPGDSLQAQFNITSQDDVLFAlFSKGQKQYHHPDDLSALCAFPPI  
 RAINLQIKERLQSCYQGEENLELNWLLGKDQCTKAPVPIDDNFCGLDINQPLGGSTPVEGLTLYTTSRDRM  
 15 TSVASYVYNGYSVVFVGTKSGKLKKIRADGPPHGGVQYEMVSVLKDGPILRDMAFS IQCRYLYVMSEQRVT  
 RVPVESCEQYTTCEGLSSGDPHCGWCALHNMCSRRDKCQAWEPNRFASISQCVSLAVHPSSISVSEHSR  
 LLSLVSDAPDLSAGIACAFGNLTEVEGQVSGSQVICISPGPKDVPVPLDQDWFGLELQLRSKETGKIFVS  
 TEFKFYNCSAHQLCLSCVNSAFRCHWCKYRNLCTHDPTTCSFQEGRINISEDQPLVPTEEILIPVGEVKPI  
 TLKARNLPQPQSGQRGYECVLNIQGAHRVPALRFNSSSVQCQNSSYQYDGMDISNLAVDFAVWNGNFIID  
 20 NPQDLKVHLYKCAAQRESCGLCLKADRKFECGWCSGERRCTLHQHCTSPSSPWLDWSSHNVKCSNPQITEIL  
 TVSGPPEGGRVTVTHGVNLGLDFSEIAHHVQVAGVPCTPLPGEYIIAEQIVCEMGHALVGTTSGPVRLCIGE  
 CKPEFTKSHQYTFVNPVSVLSLNPIRGPESGGTMVTITGHYLGAGSSVAVYLGNGTCEFYGRSMSEIVCVS  
 PPSNGLGPVPVSVSVVDRAHVDSNLQFEYIDDPVRQRIEPEWSIASGHTPLTITGFNLDVIEPRIRVKFNG  
 KESVNVCKVNTTTLTCLAPSLTDDYRPLDTPVERPDEFGFVFNQVSLIYNDTKFIYYPNPTFELLSPTG  
 25 VLDQKPGSPIILKGKNLCPASGGAKLNYTVLIGETPCAVTVSETQLLCEPPNLTGQHKVMVHVGGMVFSFG  
 SVSVISDSLLTLPALVSIAGGSLLLIIVIIIVLIAKRSRENDLTLKRLQMMDNLESRVALECKEAFEL  
 QTDINELTSDLDMSGIPYLDYRTYAMRVLPFGIEDHPVLRLELVQNGQGHVEKALKLFAQLINNKFVLLTF  
 IRTLELQRSFSMRDRGNVASLIMTGLQGRLEYATDVLKQLSDLDIDKNLENKNHPKLLLRRTESVAEKMLTN  
 WFAFLHKLKFLKECAGEPLFMYCAIKQOMEKGPIDAITGEARYSLSEDKLIRQIEYKTLILNCVNPDNENS  
 30 PEIPVKVLNCDTITQVKEKILDVYKKNVPYSQRPRAVMDLEWRQGRIRVVLQDEDITTKIEGDWKRNLTL  
 MHYQVSDRSVVALVPKQTSSYNIPASASISRTSISRYGDSFFRYTGSPDSLRSRAPMITPDLESQVGVVHLV  
 KNHDHGDQKEGDRGSKMVSEIYLRLLATKGTLOKFVDDLFETLSTVHRGSALPLAIKYMFDLDEQADRH  
 SIHDTDVRHTWKSNCPLPLRFVWNVIKNPQFVFDIHKGSITDACLSVVAQTFMDSCTSEHRLGKDSPSNKL  
 YAKDIPSYKSWERYYADIAKLPAISDQDMNAYLAEQSRLHAVEFNMLSALNEIYSYVSKYSEELIGALEQD  
 EQARRQLAYKVEQLINAMSIES

35  
 deoxycytidylate deaminase

atgagtgaag tttcctgcaa gaaacggggac gactatttgg aatggccaga gtattttatg  
 40 gctgtggcct tcttatcagc acagagaagc aaagatccaa attcccaggt cggcgccctgc  
 atcgtgaatt cagaaaacaa gattgtcggg attgggtaca atgggatgcc aaatgggtgc  
 agtgatgacg tgttgccctg gagaaggaca gcagagaata agctggacac caaatacccg  
 tacgtgtgcc atgcggagct gaatgccatc atgaacaaaa attcgaccga tgtgaaaggc  
 tgtagtatgt atgtgcctt gttcccttgt aatgaatgcg ctaagctcat catccaggca  
 45 ggtataaaaag aagtgatttt cacgtctgat aaataccatg atagtacga ggcaactgct  
 gcgaggctcc tgtttaatat ggccgggggtg acattccgga aattcatacc gaagtgcagc  
 aagattgtca ttgactttga ttcaattaac agcagaccga gtcaaaagct tcagtgaagt  
 acatctcatt caatctccag aagattggga ttatcgtctt ctaagagggt gctaattgcct  
 ttcattctga agttacacat aacttcttat tagccagtat ggcaaaagta ggcactaaa  
 50 gaatataaag cctcaaatct tccttactgt ctctcttgtc acatggaatc tacatgtgtt  
 tgaactattg ctttaggatt taaaataggg gagcctgtgg tggcctgggtg cacagggtca  
 gaacgagagt gcctcccctt cttgtgtcct ggctggctgg gatgctgggtg gctcttcaga  
 ggagcatcag ctgtctgtca tctgtgcga tccggcagcc tctcttctact gctacatgtg  
 ctggaaggac aaataaataa ttgtggttgt gttcttaatg gggacgagca gacacactga  
 tctgaacatc tggcccaagt gaagcatggc atatagtgcc cttggaagaa aattaggcct  
 55 caaatgacag tagcattgaa gtgtttgtct cagagttgag ggaaaccccc agccaccctc  
 ccggaatccg agatagggtg gcacatctgt cctgacagac gaggagtgt actgaaccag  
 gaatatttcc tccattcctg ctctcccact gcacacaggg tgggtggcaca ttatccctct  
 ggggggtggg gacgcctgtt gttttggctc aatttgggtt tgttgggtcac atggagctct  
 tccatttctg ttagctgaat aatgagttgt tcctagagga gacagcctgt ctctccttgt  
 60 tgcccccaaa gcccatgccc tgccgtgggt gcagctgggg ctgtggatgg gaggggtccc  
 caacatggat gtgttgcccc tcctccgcat gccaacgcag ttcattgtaca agggccctct

5 gcaactggag agaaaattaa ttcctatccc gtgagtggat tgtgagaaat tccacccacg  
tggagacagc ttactgcagc actggttggtg ttcggagctc ttctgtgccc tggctccatg  
ctttcaccta cacaagcatc accttcctaa tcaccgcggg gcggggagcg tgtggctgtg  
cccccttctt ttaatctcat ttaattttta ttaaaccatgc tcagtacctg tgttgagaaa  
aggctttctt tatcctaaag attattacct ttttaaagtg ctcttatatt ttcattgagtt  
tttattttgt ctctgagatt ttgtattcca cattctaggg tattctgtaa tttggctcct  
taccaatatt attaaaatct tattaaaatc t

10 MSEVSCKKRDDYLEWPEYFMAVAFLSAQRSKDPNSQVGACIVNSENKIVGIGYNGMPNGCSDDVLPWRRTAE  
NKLDTKYPYVCHAE LNAIMNKNSTDVKGCSMYVALFPCNECAKLIIQAGIKEVIFTSDKYHDSDEATAARLL  
FNMAGVTFRKFIPKCSKIVIDFDSINSRPSQKLQ